

Pesticide Residues

Results from the period 2004-2011



Pesticide Residues

Results from the period 2004 – 2011

Prepared by National Food Institute, Technical University of Denmark

Division of Food Chemistry:

Annette Petersen

Bodil Hamborg Jensen

Jens Hinge Andersen

Mette Erecius Poulsen

Division of Nutrition:

Tue Christensen

Division of Toxicology and Risk Assessment

Elsa Nielsen

Acknowledgement: The authors wish to thank all that have participated in the analysis and gaining of the results at the Regional laboratory in Ringsted

Pesticide Residues
Results from the period 2004-2011

1. edition, June 2013

Copyright: National Food Institute, Technical University of Denmark

Photo: Colourbox

ISBN: 978-87-92763-78-5

This report is available at

www.food.dtu.dk

National Food Institute
Technical University of Denmark
Mørkhøj Bygade 19
DK-2860 Søborg

Tel: +45 35 88 70 00

Fax: +45 35 88 70 01

Table of contents

| | | |
|-----|--|-----|
| 1 | Preface | 4 |
| 2 | Sammenfatning og konklusion | 5 |
| 3 | Summary and conclusion..... | 6 |
| 4 | Pesticide residues and exposure..... | 9 |
| 4.1 | Monitoring programme | 9 |
| 4.2 | Residues | 11 |
| 4.3 | Exposure..... | 44 |
| 5 | References | 58 |
| 6 | Annexes | 61 |
| 6.1 | Exposure calculations | 61 |
| 6.2 | Correction for samples with non-detected residues | 65 |
| 7 | Appendices | 68 |
| 7.1 | Commodity groups not included in exposure calculations..... | 68 |
| 7.2 | Pesticides sought in fruit and vegetables resp. cereals in 2004 – 2011 and their frequency of detection in conventionally grown crops..... | 75 |
| 7.3 | Pesticides included in the monitoring and commodities where residues were found. | 81 |
| 7.4 | Consumption used for exposure calculations..... | 119 |
| 7.5 | ADIs for pesticides included in the risk assessment..... | 122 |
| 7.6 | Reduction factors..... | 124 |
| 7.7 | Exposure and sum of Hazard Quotients (Hazard Index (HI)) for individual commodities (consumer group “Adults”). | 126 |

| | | |
|------------|--|------------|
| 7.8 | Exposure and sum of Hazard Quotients (Hazard Index (HI)) for individual pesticides (consumer group “Children”). | 129 |
| 7.9 | Exposure and Hazard Quotients for individual pesticides (consumer group “Adults”). | 132 |

1 Preface

The present report presents the results from the 2004-2011 period of the monitoring programmes conducted by The Danish Veterinary and Food Administration. The programmes included commodities of fruit, vegetable, cereals and animal origin using random sampling from food on the Danish market. Since the beginning of the 1960, Denmark has monitored fruit and vegetables for pesticides residues.

For the periods 1993-1997 and 1998-2003, results were collated and the dietary exposure was calculated. In this report data for the analyses carried out in the period 2004-2011 are reported as well as the exposure calculations performed on the background of the residues found. All the analyses have been carried out by the laboratory of the Danish Veterinary and Food Administration in Ringsted. The samples were collected by the food control offices.

The residue data have been combined with consumption data and the exposures for different consumer groups have been estimated.

2 Sammenfatning og konklusion

I denne rapport præsenteres resultaterne for kontrol for pesticidrester i fødevarer i Danmark i perioden 2004-2011. Antal stoffer varierer fra år til år, da der løbende blev inkluderet nye stoffer i analyseprogrammet. Der blev analyseret for omkring 249 pesticider, der dækker ca. 275 individuelle stoffer inklusiv metabolitter. Der blev i alt analyseret 17309 prøver af frugt, grønt, cerealier, kød, børnemad og andre forarbejdede fødevarer. Fordelingen mellem de forskellige typer af fødevarer kan ses i tabel 1.

Resultaterne viser, at der var langt flere fund af pesticider i frugt og grønt (se tabel 3) end i andre afgrøder. Sammenlignes frugt, grønt og cerealier indeholdt frugt langt flere pesticidrester (se figur 1-3). Der var generelt flere pesticidrester i udenlandske produkter i forhold til danske (se figur 1-3), og der blev hyppigere fundet flere pesticider i samme prøve i udenlandske prøver sammenlignet med prøver fra Danmark. Der blev generelt fundet overskridelser af MRL i ca. 2,6 % af prøverne. I prøver som appelsin, mandarin og banan blev der fundet pesticidrester i stort set alle prøver. Det skal dog understreges, at prøverne blev analyseret med skræl. For langt de fleste pesticider viser undersøgelser af fordelingen i den spiselige og ikke-spiselige del (f.eks. skræl), at størstedelen af de fundne indhold findes i den ikke-spiselige del.

Der er foretaget en sammenligning af den del af afgrøderne, der bidrager mest til indtaget af pesticider. Der er foretaget en sammenligning af prøver mellem de lande, hvorfra der har været udtaget mere end 10 prøver til kontrol i perioden 2004-2011. For afgrøder, der bliver dyrket både i Danmark og i udlandet, viser resultaterne generelt, at der var en mindre hyppighed af pesticidrester i danske afgrøder sammenlignet med udenlandske afgrøder. For enkelte afgrøder var hyppigheden af fund i danske prøver imidlertid ikke det laveste blandt alle lande. Det drejede sig om jordbær, blommer, gulerødder, agurk, spinat, hvedemel og hvedekerner.

Resultaterne fra analyseprogrammet er brugt til at beregne eksponeringen fra fødevarer ved at gange et gennemsnitsindhold med et gennemsnitskonsum. Der kan bruges forskellige modeller til at beregne en eksponering. Man kan inkludere, at restindhold kan forøges eller formindskes ved forarbejdning. Man kan også tage i betragtning, at prøver med indhold under rapporteringsgrænsen (LOR) kan have et indhold på 0 mg/kg eller et meget lavt indhold. Derfor er der regnet med og uden forarbejdning, samt at indholdet er 0, $\frac{1}{2}$ LOR og $\frac{1}{2}$ LOR med korrektion for indhold under LOR. I sidstnævnte beregning indgår en korrektion med en begrænsning på maksimalt en faktor 25. Dette er nærmere beskrevet i Annex 6.2.

Eksponeringen varierede mellem 44 og 144 $\mu\text{g}/\text{person}/\text{dag}$ for børn og 68 og 222 $\mu\text{g}/\text{person}/\text{dag}$ for voksne alt efter hvilken model, der blev anvendt. I de videre beregninger er der valgt en model, hvor der er taget hensyn til forarbejdning og brug af $\frac{1}{2}$ LOR med en korrektion. Det gav en eksponering på henholdsvis 98 $\mu\text{g}/\text{person}/\text{dag}$ for børn og 146 $\mu\text{g}/\text{person}/\text{dag}$ for voksne.

Risikovurderingen for et enkelt pesticid blev udført ved beregning af en såkaldt Hazard Quotient (HQ). HQ er forholdet mellem eksponeringen og det Acceptable Daglige Indtag (ADI) for pesticidet. HQ for de enkelte pesticider ligger mellem 0,00001 % og 2,35 % (de fleste under 1 %), hvilket indikerer, at der ikke er en sundhedsmæssig risiko ved indtag af de enkelte pesticider. Der er også udført en risikovurdering af det kumulative indtag af de fundne pesticider ved at summere alle HQ for de enkelte pesticider til et såkaldt Hazard Indeks (HI). HI varierer mellem 4 % og 49 % for voksne og 10 % til 124 % for børn alt efter hvilken model, der er brugt i beregningerne. Med den valgte model er HI beregnet til 18 % for voksne og

44 % for børn. Da HI metoden forudsætter samme type effekt for alle de fundne pesticider, er metoden relativt konservativ (dvs. 'på den forsigtige side'), da alle pesticider ikke har samme type af effekter. HI på 18 % for voksne og 44 % for børn indikerer således, at der ikke er en sundhedsmæssig risiko ved indtag af de fundne pesticider samtidigt.

Som tidligere nævnt blev der generelt fundet færre pesticidrester i danske afgrøder sammenlignet med afgrøder fra udlandet. Dette har også indflydelse på eksponeringen. Spiste man danske afgrøder, når det var muligt, nedsatte man både eksponering og HI. For børn faldt eksponeringen fra 98 µg/person/dag til 52 µg/person/dag, mens HI faldt fra 44 % til 20 %. For voksne er de tilsvarende tal, at eksponeringen faldt fra 146 µg/person/dag til 76 µg/person/dag, og at HI faldt fra 18 % til 8 %.

Myndighederne anbefaler voksne at spise mindst 600 g frugt og grøntsager om dagen. For mænd og kvinder er indtaget beregnet for dem, som spiser mere end 550 g frugt og grønt om dagen. Dette fik eksponeringen og HI til at stige med en faktor 1,5-2. HI var dog stadig mindre end 100 %.

Det er også beregnet hvilke pesticider og afgrøder, der bidrog mest til eksponeringen. For pesticiderne var dette fordelt mellem mange stoffer. Fordelingen var også vidt forskellig, om man så på eksponering eller HQ. For afgrøder derimod var det et mindre antal afgrøder, der bidrog både til eksponering og HI. Æbler var i begge tilfælde langt den største bidragsyder. I alt bidrog 25 forskellige afgrøder til ca. 95 % af eksponeringen og HI.

Resultaterne for perioden 2004-2011 viser lige som resultaterne for sidste periode (1998-2003), at det beregnede Hazard Index var under 100 % for både børn og voksne. Dette gælder også for voksne, der spiser mere end 550 g frugt og grønt om dagen.

Der ses en mindre stigning i den beregnede eksponering for voksne, og et fald i Hazard Index. På grund af usikkerhederne og de forskellige forudsætninger for beregningerne i de to perioder, er det ikke muligt at afklare, hvorvidt tallene er udtryk for reelle ændringer.

For børn er der beregnet for forskellige aldersgrupper i de to perioder, og derfor kan tallene ikke sammenlignes.

3 Summary and conclusion

In this report the results for the analyses of pesticide residues in foods on the Danish market are presented. The analytical programme included about 249 pesticides covering about 275 substances including metabolites. The number of substances varied from year to year due to the fact that more and more substances are included in the programme. In total 17309 samples have been analysed. The samples included fruit, vegetables, cereals, meat, baby food and other processed food. The distribution between the different kinds of commodities is shown in Table 1.

The results show that more residues were found in samples of foreign origin compared to samples of Danish origin (see Figure 1-3). Overall fruits and vegetables had higher frequencies of residues than the other groups of commodities; fruits had higher frequencies compared to vegetables. Also, samples with more than one residue were more frequently found in samples of foreign origin. Overall residues above the MRLs were found in 2.6 % of the samples, most frequently in fruit.

For some of the commodities that contributed most to the exposure the frequency of residues in samples have been compared between countries when the number of samples were higher than 10. The frequency of residues in commodities grown both in Denmark and abroad were, in general, higher in samples of foreign origin than in Danish samples. Also, samples with residues above the MRLs were more often in samples of foreign origin. However, for strawberries, plums, carrots, cucumbers, spinach, wheat flour and wheat the frequencies in Danish samples were higher compared to some of the other countries.

The results from the analytical programme have been used to calculate the exposure for the Danish population by multiplying a mean of the residues with a mean of the consumption. There is no common agreement in EU or internationally on how to calculate the exposure; e.g. if a processing factor shall be included or not or residues below the reporting limit (LOR) also called non-detects shall be included as zero or with a value, e.g. $\frac{1}{2}$ LOR. To investigate the influence from these factors the exposure for adults (15-75 years) and children (4-6 years) were calculated with several models. This includes using a reduction factor for peeling for some fruits as well as using $\frac{1}{2}$ LOR and $\frac{1}{2}$ LOR with correction for non-detects. The correction is used to restrict the difference of using $\frac{1}{2}$ LOR instead of 0 to a factor of 25.

The exposure estimated using the different models varied between 44 and 144 $\mu\text{g}/\text{person}/\text{day}$ for children and between 68 and 222 $\mu\text{g}/\text{person}/\text{day}$ for adults. It was decided that the most appropriate would be to continue with a model taking peeling into consideration as well as eventual residues below LOR. Therefore, $\frac{1}{2}$ LOR with correction for non-detects was used in the model together with peeling. The different models are further described in Annex 6.2. With this model, the exposure was estimated to be 98 $\mu\text{g}/\text{person}/\text{day}$ and 146 $\mu\text{g}/\text{person}/\text{day}$ for children and adults, respectively.

The risk assessment for a single pesticide is performed by estimation of the so-called Hazard Quotient (HQ). The HQ is calculated by dividing the exposure with the Acceptable Daily Intake (ADI) for the individual pesticide. The HQs for the individual pesticides ranged from 0.00001% to 2.35% with most of the HQs being below 1% indicating no risk of adverse effects following exposure to the individual pesticides. Risk assessment of the cumulative exposure to the detected pesticides has been performed by summing up the HQs for the individual pesticides to provide a so-called Hazard Index (HI). HI varied between 4% and 49% for adults and 10% and 124% for children. With the chosen model, the HI is 18% for adults and 44% for children. As the HI method assumes the same kind of adverse effect for all the detected pesticides it is a relatively conservative (i.e. precautionary) approach for cumulative risk assessment. Overall, the HI of 18% for adults and 44% for children is not considered to indicate a risk of adverse effects following a cumulative exposure to all the detected pesticides.

As mentioned above, commodities of Danish origin, generally, contained fewer pesticides compared to commodities of foreign origin. This can also have an influence on the exposure. If commodities of Danish origin were chosen whenever possible, the exposure and HI decreased. For children, the exposure decreased from 98 $\mu\text{g}/\text{person}/\text{day}$ to 52 $\mu\text{g}/\text{person}/\text{day}$ while HI decreased from 44% to 20%. For adults the exposure decreased from 146 $\mu\text{g}/\text{person}/\text{day}$ to 76 $\mu\text{g}/\text{person}/\text{day}$ while the HI decreased from 18% to 8%.

For men and women the exposure has been estimated for those who consumed more than 550 g of fruit and vegetables every day. The exposure calculations showed that their exposure was 1.5-2 times higher than that for the average consumer. However, the HI was below 100% in both cases.

It has also been estimated which commodities and pesticides that contributed most to the exposure and HI. For the pesticides the contributions were distributed between several pesticides and there was a big difference in the pesticides that contributed most to the exposure and HI. However, only a limited number of commodities contributed most both to the exposure and HI. Apples were, under all circumstances, the greatest contributor. About 95% of the exposure and HI were accounted for by 25 different commodities.

The results from the period 2004-2001 show, as for the previous period (1998-2003) that the calculated Hazard Index was below 100% for both adults and children. This was also the case for adults consuming more than 550 g of fruit and vegetables per day.

For adults a minor increase in the calculated exposure and a decrease in Hazard Index were observed. However, due to the uncertainties and the different basis for the calculations in the two periods a clear conclusion on the changes cannot be drawn.

For children the calculations have been performed for two different age groups in the two periods, and therefore it is not possible to compare the results.

4 Pesticide residues and exposure

4.1 Monitoring programme

Over the years, the number of pesticides analysed has increased and was 249 in 2011. The number of substances including isomers and metabolites was approximately 275. The total number of samples in the period 2004-2011 has been relatively stable. However, the fruit, vegetable and cereal samples have increased and the number of samples of animal origin has decreased slightly (see **Figure 1**). The results have been published each year in the period 2004-2011 (Pesticidrester i fødevarer, 2004; Christensen et al., 2005; Christensen et al., 2006; Christensen et al., 2007; Petersen et al., 2008; Jensen et al., 2009; Jensen et al., 2010; Jensen et al., 2011).




Figure 1. Numbers of fruit, vegetable and cereal samples and samples of animal origin analysed from 2004-2011

Design of sampling plan

The Danish pesticide monitoring programme has two objectives. Firstly, the programme had to check compliance with the maximum residue levels laid down by the EU (EU Commission, 2005), and secondly to monitor the residue levels in foods to enable an evaluation of the exposure of the Danish population to pesticides.

The sampling plan for the period 2004 to 2006 was structured in the same manner as the period 1998-2003. A detailed description can be seen in Poulsen et al. (2005). The design of the sampling plan was changed in 2006 and has been kept the same way since then. The sampling plan for fruit and vegetables was designed in two parts. Part one included samples that can be used to estimate the exposure of pesticides and to see possible trends in pesticide residues between years. For 25 different fruit, vegetable and cereal commodities, a fixed number of 50 samples per year were collected. The commodities were chosen, based on their contribution to

the intake of pesticides for the Danish population calculated on the monitoring results from the period 1998-2003 (Poulsen et al., 2005). This included also processed commodities like wine. Likewise, 15 samples were collected of commodities included in the EU Multiannual Pesticide Control Programme (EU Commission, 2012). Part two included samples that contributed in minor degree to the intake of pesticides, but where the control was focussed on the compliance with MRLs or labelling of production method, e.g. organic grown, produced without growth regulator or surface treatment. Part one consisted of 70% of the fruit and vegetable and 15% of the cereal samples.

Sampling

Authorised personnel from regional food control units under the Danish Veterinary and Food Administration performed the sampling and collected the samples randomly within each commodity. The sampling procedure conformed to the EU directive on sampling for official control of pesticide residues (European Union, 2002). A total of 17309 samples were taken primarily at wholesalers, importers, slaughterhouses and at food processing companies (see **Table 1**). Most of the samples were conventionally grown fresh fruits and vegetables (70%), but also conventionally grown cereals (10%) and samples of animal origin (11%) were collected. In addition 6% samples of organically grown crops (fresh, frozen, processed) were collected as well as processed foods (e.g. wine) and samples of baby food. One third of the fruit and vegetable samples and two thirds of the cereals samples were of Danish origin. For meat more than 90% of the samples were of Danish origin. Almost 175 different fruits, vegetables and cereal commodities were sampled, of these 73 were also organically produced.

Sampling of meat and other products of animal origin are regulated by EU Directive 96/23/EC. The aim of this directive is to ensure that the Member States monitor primarily their own production of commodities of animal origin for different substances e.g. pesticides. However, imported samples from third countries shall also be monitored. Depending on the animal species the number of samples was between 0.03% and 0.15% of the production or import.

For fruits, vegetables, and cereals the aim has been to monitor the commodities sold on the Danish market. Consequently, more samples produced in EU Member States and third countries have been collected compared to samples of Danish origin.

Table 1. Number of samples analysed for the period 2004-2011, Danish and foreign origin, respectively.

| Foods | Danish | Foreign | Total |
|--|---------------|----------------|--------------|
| Fruit and vegetables (fresh, frozen, processed) | 2844 | 9182 | 12026 |
| Cereals (including processed) | 717 | 1060 | 1777 |
| Wine | 2 | 273 | 275 |
| Meat | 1589 | 358 | 1947 |
| Milk and honey | 146 | 0 | 146 |
| Baby food | 28 | 38 | 66 |
| Organically grown fruit, vegetables and cereals (fresh, frozen, processed) | 358 | 714 | 1072 |
| Total | 5684 | 11625 | 17309 |

Laboratories

Samples were primarily analysed at the Regional Food Laboratories. However, a few of the samples were analysed at DTU National Food Institute. All laboratories involved in the monitoring were accredited for pesticide analysis in accordance to ISO 17045 by the Danish body of accreditation, DANAK.

Analytical Programme

Analytical methods were developed and documented at the DTU National Food Institute.

Fruits and vegetables were analysed by up to five different analytical methods covering 149-238 pesticides (see **Table 2**) and including isomers and metabolites the number of substances were approximately 275. Cereals were analysed by three different methods and meat by one method. The number of analytical methods used for other commodities differs depending on the matrices.

Table 2. Number of pesticides, analysed for in the period 2004-2011 in different types of foods. Isomers and metabolites are not included

| Foods/Year | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 |
|----------------------|------|------|------|------|------|------|------|------|
| Fruit and vegetables | 149 | 158 | 164 | 176 | 176 | 222 | 238 | 230 |
| Cereals | 105 | 114 | 112 | 112 | 170 | 165 | 165 | 166 |
| Meat | 30 | 28 | 28 | 28 | 28 | 28 | 28 | 44 |
| Baby/infant food | 153 | 0 | 166 | 178 | 207 | 215 | 240 | 238 |

The pesticide profile is shown in Appendix 7.2

4.2 Residues

In **Table 3** the average frequencies of samples with residues are shown. However, the figures have a large variation covering commodities with very low frequencies and others where, practically all samples contained residues. Among the fruit and vegetables commodities, nuts and some of the herbs had the lowest frequencies of sample with residues. Other commodities with low frequencies were Danish produced courgette, broccoli and leek, all with 5% samples with residues. Likewise, foreign produced Chinese cabbage, cauliflowers/broccoli and sweat corn contain 0%, 2% and 5% samples, respectively, with residues. On the other end of the scale, were especially the foreign produced commodities from the citrus group, peas with pods, and banana/papaya with high frequencies of positive samples, namely 98%, 95% and 85%, respectively. No Danish produced fruit and vegetables had that high frequency of samples with residues.

Although the commodities in the group of processed fruit and vegetable are more limited there is also some variation in the frequencies from mixed juice with 0% to orange marmalade with 52% samples with residues.

The cereals covers less commodities and the cultivation is, likely, more comparable resulting in less variation in the frequencies.

However, in general the exposure of pesticides differs from commodity to commodity. This is described in Section 4.3. The frequencies listed above in **Table 3** have to be considered as the lowest possible frequency, since the pesticide profile in the analytical methods did not cover

all pesticides used in Denmark or in the countries exporting to Denmark. Furthermore, it is supposed that there will be residues below the detection limits.

Table 3. *Frequency of samples with residues*

| Foodstuff | Frequency of samples with residues¹ | Frequency of samples above MRL |
|---|---|---------------------------------------|
| Fruit and vegetables (fresh, frozen) | 53% | 3.8% |
| Cereals (including processed) | 27% | 0.1% |
| Meat | 0% | 0.0% |
| Baby food | 0% | 0.0% |
| Milk and honey | 0% | 0.0% |
| Processed fruit and vegetables | 42% | 0.2% |
| Processed cereals | 12% | 0.0% |
| Organically grown fruit, vegetables and cereals | 2% | 0.4% |
| Total | 40% | 2.6% |

1. Includes also samples above MRL

Comparison between Danish and foreign produced commodities

Figure 2 shows the frequencies of samples without detections, with detections below, the MRL at the MRL or above the MRL for fruit commodities produced in Denmark, the EU, and outside the EU.

In general, samples of fruit commodities produced in Denmark had lower frequencies of detections below MRL (38-67%) than fruit commodities produced outside Denmark (61-82%). However, the fruit commodities were not the same as many fruits, which cannot be grown in Denmark (e.g. oranges, pineapples). It seems as though the frequencies of samples without detection have increased throughout the years. No differences were seen between samples produced in the EU and outside the EU, except for samples with detection above MRL, where samples produced outside the EU, more frequently, had residues above MRL, namely, 4-7% and 0-3% respectively.




Figure 2. Pesticide residues detected in fruit produced in Denmark, the EU and outside the EU

Figure 3 shows frequencies of samples without detections, with detection below MRL and above MRLs for vegetable commodities produced in Denmark, the EU and outside the EU. In general, there were fewer samples of vegetable with residues compared to fruit. Furthermore, vegetables produced in Denmark had lower frequencies of detections below MRL (7-12%) than vegetables produced outside Denmark (22-42%). The type of vegetables produced in Denmark and outside was in general the same, except for commodities like sweet peppers and beans with pods. No differences were seen between samples produced in the EU and the outside EU, except for samples with detection above MRL, where samples produced outside the EU more frequently had residues over the MRL namely respectively 5-14% and 0-6%.




Figure 3. Pesticide residues detected in vegetables produced in Denmark, the EU, and outside the EU

Figure 4 shows frequencies of samples without detections, with detection below MRL and above MRLs for cereal commodities produced in Denmark, the EU, and outside the EU. In general, fewer samples of cereal commodities have residues compared to fruit and vegetables. Cereals produced in Denmark had lower frequencies of detections below MRL (8-26%) than cereals produced in the EU (39-67%), while cereals grown outside EU had residues frequencies at the same level as Denmark (9-30%). The type of cereals produced in Denmark and the EU was different from cereals produced outside the EU. The cereal samples produced outside the EU were mainly rice and the samples from the EU and consisted mainly of wheat, oat, rye and barley. Residues above MRLs are rarely found in cereals. For the period 2004-2011 only one Danish sample from 2010 exceeded the MRL (chlormequat in oat).




Figure 4. Pesticide residues detected in cereals produced in Denmark, the EU and outside the EU

Detailed evaluation of pesticide residue in commodities produced in different countries.

Although pesticide residues seldom exceed the maximum residue limits, consumer awareness on health issues, due to food contaminants, is high. Many consumers are anxious about the pesticides residues in their food and want to avoid them, as much as possible. The commodities consumed in Denmark are produced in many different countries and the monitoring data have been evaluated to see if there were differences in the frequencies of samples with residues produced in Denmark and outside Denmark, or between countries. For twelve fruit, eight vegetable and three cereal commodities, differences were observed in frequencies of detected pesticide residues. The results from these commodities are evaluated more closely, later in this paragraph.

Only commodities, where more than 10 samples from the same country were analysed, are included in this evaluation. Therefore, figures are only given for countries from where more than 10 samples were collected.

For bananas, grape fruit, lemon and oranges, no differences were seen as pesticides were detected in almost all samples

Bananas

Most samples taken originated from Columbia, Costa Rica, Ecuador (> 90%). In total, 425 banana samples were collected and analysed in the period 2004-2011. Most of them, 388, were exported from these four countries. The rest were imported to Denmark from 11 different countries (Bermuda, Brazil, Chile, Dominican Republic, Guatemala, Honduras, Israel, Mexico, Panama, Thailand, and the USA). In total, 85% of the samples contained 14 different pesticide residues; no residues were above the MRLs. Most of the frequencies were 80-70 and 50% of the samples contained more than one residue.

Mandarins and clementine

Most samples taken originated from Spain. In total, 411 samples of mandarins and clementine were collected and analysed in the period 2004-2011. Most of them, 322 (78%), were produced in Spain. The rest of the samples were produced in 12 different countries (Australia, Cyprus, Italy, Pakistan, Peru, the USA, Chile, Israel, Turkey, Morocco, South Africa and Uruguay). In total, 99% of the samples contained pesticide residues, 4% above the MRLs and 55 different pesticides were found. Almost all the samples had residues and 93% of the samples contained more than one residue.

Grapefruit

The main exporters to the Danish market were South Africa, Turkey, the USA and Israel. In total, 388 grapefruit samples were collected and analysed in the period 2004-2011. Most of them, 274, were produced in these four countries. The remaining samples were produced in 16 different countries (Argentina, Chile, Costa Rica, Cuba, Cyprus, Egypt, Greece, Honduras, Jamaica, Mexico, Morocco, Netherlands, Switzerland, Spain, Swaziland, and Zimbabwe). In total, 99% of the samples contained pesticide residues, 5% above the MRLs and 46 different pesticide residues were found. More than one residue was found in 86% of the samples.

Lemons

Most samples taken originated from Spain. In total, 381 lemon samples were collected and analysed in the period 2004-2011. Most of them, 243 (64%), were produced in Spain. The rest of the samples were produced in 8 other countries (Argentina, Israel, Italy, Morocco, South Africa, Turkey, the USA and Uruguay). In total, 97% of the samples contained pesticide residues, 1% above the MRLs and 39 different residues were found. Almost all the samples had residues and it was also found that 79% of the samples contained more than one residue.

Oranges

Most samples taken originated from Spain, Greece, South Africa and Morocco. In total, 491 orange samples were collected and analysed in the period 2004-2011. Most of them, 400, were produced in these four countries. The rest were produced in 12 other countries (Argentina, Australia, Brazil, Chile, Egypt, Israel, Italy, Portugal, Swaziland, Turkey, Uruguay and Zimbabwe). In total, 98% of the samples contained pesticide residues and 49 different pesticide residues were found. In 82% of the samples more than one residue was detected.

Apples

Apples are grown in Denmark and approximately 40% of total the number of samples in the period 2004-2011 was covered by Danish apples. Approximately, one third of the samples were produced in France and Italy and the rest were produced in 13 other countries (Argentina, Belgium, Brazil, Chile, China, Germany, Italy, New Zealand, Poland, Spain, South Africa, Uruguay and the USA). In total, 585 apple samples were collected and analysed

The Danish produced apples had residues of 12 different pesticides in 46% of the samples and 2% contained residues above the MRLs. The foreign produced apples had residues of 54 different pesticides in 80% of the samples and 3% contained residues above the MRLs. A major difference was seen between Danish apples and apples of foreign origin in respect to the number of pesticides found and the frequency of samples with residues. **Figure 5** shows the differences between apples samples. Most of the frequencies in foreign produced apples were between 58-100% and 39% of the foreign produced samples contained more than one residue, while only 9% of the Danish produced samples had multiple residues.




Figure 5. Frequencies of samples with residues for apples produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Table grapes

Most samples taken originated from South Africa, Italy, Chile, India and Spain. In total, 460 grape samples were collected and analysed in the period 2004-2011 and 354 (77%), were produced in the five mentioned countries. The rest were produced in 12 other countries (Argentina, Australia, Brazil, Egypt, Greece, Israel, Namibia, Netherlands, Peru, Saudi-Arabia, Turkey, and the USA). In total, 79% of the samples contained pesticide residues, 1% above MLRs and 54 different pesticide residues were found.

Table grapes from Brazil, Egypt, South Africa and Argentina contained residues in 50-75% of the samples while samples from the other countries with more than 10 samples showed higher frequencies. **Figure 6** shows the differences between table grapes samples. Most of the frequencies were 60-90%, and in 49% of the samples more than one residue was detected.




Figure 6. Frequencies of samples with residues for table grapes produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Melons

Most samples taken originated from Spain, Brazil and Costa Rica. In total, 312 melon samples were collected and analysed in the period 2004-2011 and 279 (89%) of the samples were produced in these three countries. The rest originated from 11 other countries (Croatia, Ecuador, France, Israel, Mauritania, Morocco, Netherlands, Panama, Thailand, Turkey, and the USA). In total, 54% of the foreign samples contained pesticide residues, 3% above MRLs and 38 different pesticide residues were found. **Figure 7** shows the differences between the five countries where more than 10 samples have been collected. Most of the frequencies were 40-94%, and 1% of the samples contained more than one residue.




Figure 7. Frequencies of samples with residues for melons produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Strawberries

Strawberries are grown in Denmark and approximately 40% of the samples were of Danish origin in the period 2004-2011. The main part of the samples was produced in Spain, Poland, Germany, and Belgium (45% of the samples). The rest of the samples (15%) were produced in 10 other countries (China, Egypt, Hong Kong, Israel, Italy, Morocco, Netherlands, Portugal, South Africa, and the USA). In total, 429 strawberry samples were collected and analysed.

The Danish produced strawberries contained residues of 11 different pesticides in 60% of the samples and none with residues above the MRLs. The foreign produced strawberries had residues of 45 different pesticides in 70% of the samples and 5% above the MRLs. Samples from China, Poland and Egypt had lower frequencies of samples with residues than the Danish produced samples (see **Figure 8**), while samples from other countries contained higher frequencies. Most of the frequencies were 50-85%, and 39% of the foreign produces samples contained more than one residue while 18% of the Danish produced samples had multiple residues.




Figure 8. Frequencies of samples with residues for strawberries produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Peaches and nectarines

Most samples taken, namely 342, originated from Spain, Italy and Chile. The rest, were produced in seven other countries (Argentina, Egypt, France, Greece, Morocco, Netherlands and South Africa). In total, 69% of the samples contained pesticide residues, 1% above the MRLs, and 50 different pesticide residues were found. **Figure 9** shows the differences between the countries. Most of the frequencies were 32-92%, and 38% of the samples contained more than one residue.




Figure 9. Frequencies of samples with residues for peaches and nectarines produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Pears

Pears are grown in Denmark, and approximately 30% of the collected samples in the period 2004-2011 were covered by Danish pears. Most samples originated from the Netherlands covering about one third of the samples taken. The rest of the samples originated from 10 other countries (Argentina, Belgium, Chile, China, France, Germany, Italy, Poland, Spain and South Africa). In total, 466 pear samples were collected and analysed.

The Danish produced pears had residues of six different pesticides in 55% of the samples and 1% above the MRLs. The samples of foreign origin had residues of 48 different pesticides in 88% of the samples (see **Figure 10**). Most of the frequencies were between 55-100% and 59% of the foreign produced samples contained more than one residue while 15% of the Danish produced samples had multiple residues.




Figure 10. Frequencies of samples with residues for pear produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Plums

Plums are grown in Denmark, and approximately 20% of the collected samples the period 2004-2011 were covered by Danish plums. Most of the samples originated from Spain, South Africa and Chile covering about one third of the market. The remaining samples were produced in 7 other countries (Argentina, Belgium, France, Italy, Portugal, Serbia-Montenegro, and Turkey). In total, 391 pear samples were collected and analysed.

The Danish produced plums had residues of 11 different pesticides in 33% of the samples and 4% above the MRLs. Samples of foreign origin contained residues of 31 different pesticides in 47% of the samples (see **Figure 11**). Samples from Argentina and France had lower frequencies of samples with residues than the Danish produced plums, while samples from other countries contained higher frequencies. Most of the frequencies were between 15-85% and 11% of the foreign produces samples contained more than one residue while 8% of the Danish produced samples had multiple residues.




Figure 11. Frequencies of samples with residues for plums produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Beans with pods

Only six samples produced in Denmark were analysed and no pesticide residues were found. Most of the samples originated from Kenya, Germany, Egypt, Netherlands and Morocco. In total, 311 bean samples were collected and analysed in the period 2004-2011 and 240 of them (79%), were produced in the five afore mentioned countries. The rest were originated from 16 other countries (Belgium, Ethiopia, France, Grenada, India, Israel, Italy, Malaysia, Senegal, South Africa, Spain, Sweden, Thailand, Turkey, Vietnam, and Zambia). In total, 51% of the samples of foreign origin contained pesticide residues, 10% above MRLs and 43 different pesticide residues were found.

Differences were observed between the countries and can be seen in **Figure 12**. Most of the frequencies were 50-75%, and 18% of the samples contained more than one residue.




Figure 12. Frequencies of samples with residues for beans with pods produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Carrots

Carrots are grown in Denmark, and approximately 70% of the collected samples in the period 2004-2011 were covered by Danish carrots. Most of the samples originated from Italy and Germany covering 20% of the market. The rest were produced in 7 other countries (Belgium, France, Netherlands, Poland, Portugal, Spain and Sweden). In total, 499 carrot samples were collected and analysed.

The Danish produced carrots had residues of 9 different pesticides in 16% of the samples (see **Figure 13**). Four of the pesticides were pollutants from earlier use and these residues were all above the MRLs. The samples of foreign origin had residues of 19 different pesticides in 33% of the samples. The exception was carrots from Belgium. Most of the frequencies were 16-43%, and 10% of the foreign produces samples contained more than one residue while only 1% of the Danish produced samples had multiple residues.




Figure 13. Frequencies of samples with residues for carrots produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Cucumbers

Cucumbers are grown in Denmark, and approximately 50% of the collected samples in the period 2004-2011 were covered by Danish cucumbers. Most of the samples originated from the Netherlands and Spain (50%). In total, 417 cucumbers samples were collected and analysed. Of these, 14 samples came from Egypt, Jordan, Macedonia, Poland, Sweden and Turkey.

The Danish produced cucumbers had residues of 6 different pesticides in 38% of the samples; none above the MRLs (see **Figure 14**). The samples of foreign origin had residues of 36 different pesticides in 58% of the samples, 3% above the MRLs. However, there is a major difference for the Netherlands and Spain. The frequency of positive samples was about the same in Denmark and the Netherlands while the frequency in samples from Spain is considerably higher; about 37% compared to 85% (see **Figure 14**). Most of the frequencies were 37-85%, and 6% of the foreign produced samples contained more than one residue while 7% of the Danish produced samples had multiple residues.




Figure 14. Frequencies of samples with residues for cucumbers produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Lettuce

Lettuce is grown in Denmark and approximately 50% of the collected samples in the period 2004-2011 were covered by Danish lettuce. One third of the samples originated from Spain and Germany while the rest were produced in 9 other countries (Albania, Belgium, France, Israel, Italy, Netherlands, Poland, Portugal, and Sweden). In total, 371 lettuce samples were collected and analysed.

The Danish produced lettuce had residues of 8 different pesticides in 13% of the samples (see **Figure 15**), 1% above the MRLs. The samples of foreign origin had residues of 36 different pesticides in 45% of the samples, 4% above the MRLs. A major difference was seen between the different countries but Danish produced lettuce has under all circumstances a much lower frequency of samples with residues compared to foreign samples. Most of the frequencies were 13-65%, and 25% of the foreign produces samples contained more than one residue while 2% of the Danish produced samples had multiple residues.




Figure 15. Frequencies of samples with residues for lettuce produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Sweet peppers

Only 2 samples produced in Denmark were analysed and no pesticide residues were found. Most of the samples originated from the Netherlands and Spain. In total, 387 sweet pepper samples were collected and analysed in the period 2004-2011 and 278 (72%) of them were produced in the two afore mentioned countries. The rest, originated from 11 other countries (Belgium, Colombia, Egypt, Germany, Greece, Hungary, Israel, Morocco, Poland, South Africa, and Turkey). In total, 38% of the samples of foreign origin contained pesticide residues, 2% above MRLs and 52 different pesticide residues were found.

Figure 16 shows the differences between the countries. Most of the frequencies were 11-80% and 17% of the samples contained more than one residue.




Figure 16. Frequencies of samples with residues for sweet peppers produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Potatoes

Potatoes are grown in Denmark, and approximately 85% of the collected samples in the period 2004-2011 were covered by Danish potatoes. Most of the samples originated from the United Kingdom and France. In total, 669 potato samples were collected and analysed. Almost 75% were produced in Denmark, 10% in the UK and France and 15% in 8 other countries (Cyprus, Egypt, Germany, Israel, Italy, Morocco, Spain, and the USA).

The Danish produced potatoes had residues of 4 different pesticides in only 2% of the samples (see **Figure 17**), none above the MRLs. One of the pesticides was pollutant from earlier uses. The foreign produced samples had residues of 7 different pesticides in 25% of the samples, none above the MRLs. One of the pesticides was pollutant from earlier use. Differences were seen between Danish and potatoes of foreign origin with respect to both the number of pesticides found and the frequency of samples with residues. Most of the frequencies were 2-54%, and 3% of the foreign produces samples contained more than one residue while none of the Danish produced samples had multiple residues.




Figure 17. Frequencies of samples with residues for potatoes produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Spinach

Spinach is grown in Denmark, and approximately 33% of the collected samples in the period 2004-2011 were covered by Danish spinach. Most of the samples originated from Italy and Germany (50%). The rest of the samples were produced in 7 other countries (Belgium, France, Netherlands, Poland, Spain, Sweden, and Thailand). In total, 244 spinach samples were collected and analysed.

The Danish produced spinach had residues of 9 different pesticides in 20% of the samples (see **Figure 18**). 7% was above the MRLs. One of the pesticides was a pollutant from earlier use. The foreign produced samples had residues of 25 different pesticides in 34% of the samples, 7% above the MRLs. One of the pesticides was a pollutant from earlier use. Danish and Belgian spinach samples had frequencies below 25%, while the other foreign produced spinach had up to 69%. Most of the frequencies were 17-69%, and 1% of the foreign produces samples contained more than one residue while 1% of the Danish produced samples had multiple residues.




Figure 18. Frequencies of samples with residues for spinach produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Tomatoes

Tomatoes are grown in Denmark, and approximately 45% of the collected samples in the period 2004-2011 were of Danish origin. Most of the samples originated from Spain and Netherlands (45%). The remaining samples were produced in 10 other countries (Belgium, Egypt, France, Israel, Italy, Morocco, Poland, Senegal, Thailand and Turkey). In total, 462 tomato samples were collected and analysed.

The Danish produced tomatoes had residues of six different pesticides in only 5% of the samples (see **Figure 19**) with none above the MRLs. The foreign produced had residues of 48 different pesticides in 59% of the samples, 1% above the MRLs. A major difference was seen between Danish and tomatoes of foreign origin with respect to the number of pesticides found and the frequency of samples with residues. However, also frequencies for the foreign samples varied greatly (see **Figure 19**). The foreign produced tomatoes had frequencies in the range of 33-82%, and 35% of the foreign produces samples contained more than one residue while only 1% of the Danish produced samples had multiple residues.




Figure 19. Frequencies of samples with residues for tomatoes produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Rolled oat

Oat is grown in Denmark, and approximately 40% of the samples of rolled oat were of Danish origin. The origin of 30% of the analysed samples was unknown, but samples from Germany, Poland, Sweden, and the United Kingdom were collected and analysed.

The Danish produced rolled oat had residues of 2 different pesticides in 4% (see **Figure 20**) of the samples, none above the MRLs. The unknown origin and the foreign produced samples had residues of 6 different pesticides in 39% of the samples; none above the MRLs. Samples originating from Denmark or Sweden and samples of unknown origin had frequencies below 18%, while rolled oat samples from UK and Germany had frequencies between 80-84%, and 5% of the foreign produces samples contained more than one residue while none of the Danish produced samples had multiple residues.




Figure 20. *Frequencies of samples with residues for rolled oat produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.*

Wheat flour

Wheat is grown in Denmark, and approximately 40% of samples of wheat flour in the period 2004-2011 were of Danish origin. The origin of 38% of the analysed samples was unknown, but samples from Germany, Hong Kong, Italy, Lithuania, Poland, Sweden, Thailand, the United Kingdom, and Hungary were collected and analysed.

The Danish produced wheat flour had residues of 5 different pesticides in 19% of the samples (see **Figure 21**), none above the MRLs. The unknown origin and the foreign produced samples had residues of 7 different pesticides in 39% of the samples, none above the MRLs. Samples from Denmark and unknown origin had frequencies below 25%, while wheat flour samples from Italy and Germany had frequencies between 67-88%, and 10% of the foreign produced samples contained more than one residue while only 2% of the Danish produced samples had multiple residues.




Figure 21. Frequencies of samples with residues for wheat flour produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Wheat kernels

Wheat is grown in Denmark, and approximately 50% of the collected samples of wheat kernels in the period 2004-2011 were covered by Danish wheat kernels. The origin of 11% of the analysed samples was unknown, but samples from Czech Republic, Estonia, France, Germany, India, Kazakhstan, Netherlands, Origin unknown, Poland, Sweden, Turkey and United Kingdom were collected and analysed.

The Danish produced wheat kernels had residues of 4 different pesticides in 23% (see **Figure 22**) of the samples, none above the MRLs. The unknown origin and the foreign produced samples had residues of 9 different pesticides in 26% of the samples; none above the MRLs. Frequencies can be seen in **Figure 22** and for samples of unknown origin, Sweden and Germany, the frequencies were between 9-41%. Independent of origin, 1% of the samples contained more than one residue.




Figure 22. Frequencies of samples with residues for wheat produced in different countries. Figures close to the bars are the number of samples analysed for the specific country.

Products with low frequencies of samples with residues

In addition to the commodities of fruit, vegetables and cereals mentioned above, several commodities with few residues have been analysed. This included meat, organic grown products, baby food, and processed food. No pesticide residues were found in any of the meat samples or other samples of animal origin. For baby food, 66 samples were analysed, and residues were not found. Processed food contained fewer residues than the raw material used, because of peeling, cooking, mixing, etc., can decrease the concentration. However, the commodities still reflected the situation of the detection frequencies of the raw materials. Consequently, commodities like orange juice, orange marmalade, wine and raisin had high frequencies compared to other processed foods.

In the organic grown products, residues were found in 3% of the samples of fruits and vegetables and in about 1% of the samples of cereals. No residues were found in baby food, food of animal origin or processed food, (see Appendix 7.1).

Multiple residues

Residues from several different pesticides, 2-10, were found in 28% of all samples and in 98 different commodities, see **Figure 23**. Especially, citrus fruits contained multiple residues, in more than 75% of the samples. The samples with the highest number of pesticides were chili peppers from Thailand, where 10 different pesticides were detected. Another chili sample from Thailand contained 9 different pesticides, and this was also the case for two table grape samples, from Italy and Chile. Samples with 8 detected pesticides were 3 chili peppers (Spain, Thailand and Vietnam), two lettuce samples (Belgium and France), and one apple sample (France), one pear sample from Belgium and one pea with pod sample from Kenya. It should be emphasised that it is not necessarily an individual fruit or vegetable that contained all the detected pesticides, since the analysed samples are composed of more than one fruit or vegetable, e.g. at least 10 individual fruits. The composite sample can also consist of commodities produced by different growers. **Table 4** shows the commodities with multiple residues and where more than 30 samples have been analysed for the period 2004-2011.




Figure 23. Number of samples with 2-10 residues per sample for the period 2004-2011.

Table 4. Percentage of samples with multiple residues. Only commodity types where more than 30 samples have been analysed for the period 2004-2011 are included.

| Commodities | Samples, multiple residues | Commodities | Samples, multiple residues | Commodities | Samples, multiple residues |
|-----------------|----------------------------|----------------------|----------------------------|---------------|----------------------------|
| Apricots | 52% | Lettuce | 15% | Potatoes | 1% |
| Apples | 39% | Limes | 50% | Rambutan | 26% |
| Aubergines | 11% | Mandarin, clementine | 93% | Raspberries | 31% |
| Bananas | 50% | Mangoes | 17% | Red currants | 51% |
| Beans with pods | 18% | Melons | 1% | Rice | 2% |
| Blackberry | 9% | Mushrooms, cult. | 8% | Rye flour | 6% |
| Blue berries | 20% | Oat kernels | 7% | Rye kernels | 2% |
| Carrots | 4% | Oranges | 82% | Spelt | 16% |
| Celery | 19% | Papaya | 61% | Spelt flour | 17% |
| Chilies | 48% | Parsley root | 5% | Spinach | 4% |
| Courgettes | 5% | Parsnip | 3% | Spring onions | 17% |
| Cucumbers | 17% | Passions fruits | 37% | Star fruit | 32% |
| Grapefruits | 86% | Peaches, nectarines | 38% | Strawberries | 37% |
| Grapes | 49% | Pears | 44% | Sweet peppers | 17% |
| Rolled oats | 3% | Peas with pods | 61% | Tea | 13% |
| Kakis | 7% | Peas without pods | 6% | Tomatoes | 19% |
| Kiwis | 10% | Pineapples | 23% | Water melons | 8% |
| Leeks | 2% | Plums | 11% | Wheat flour | 7% |
| Lemons | 79% | Pomelos | 61% | Wheat kernels | 6% |

Conclusion on residues and frequencies of the found pesticides

The overall conclusion on residues in the fruit and vegetables, responsible for the major part of the exposure to pesticides (in $\mu\text{g}/\text{day}$) is that Danish produced fruit, vegetables and cereals had lower frequencies of samples with pesticide residues, compared to products of foreign origin. It is estimated that the foreign produced commodities showed about 20% higher frequencies. Also, a smaller number of different pesticides were found in the Danish products. However, some of the foreign produced commodities had comparable detection frequencies with the Danish produced commodities, or even lower. This was the case for pears (China), plums (Argentina and France), strawberries (China, Egypt), carrots (Belgium), cucumber (Netherlands), spinach (Belgium) and rolled oat (Sweden). For other foreign produced commodities various differences between countries were observed. Residues from several different pesticides, 2-10, were found in 28% of all samples and in 98 different commodities

Pesticides found in fruit and vegetables

In Appendix 7.3 is shown which pesticides that have been found in fruit and vegetables in the period 2004-2011. **Figure 24** shows the frequencies of the most often found pesticides (> 1% of the samples).




Figure 24. Frequencies of pesticides found in fruit and vegetables higher than 1% in conventionally grown domestic, foreign or total samples. The pesticides were sought in approximately 12,500 samples, except for the following: a) dithiocarbamater: 7902, not analysed in samples with high background level of sulphur, like onion, cabbage etc., b) chlormequat: 1836, only analysed in specific commodities like fruiting vegetables, table grapes, pears, c) 2,4-D, pyraclostrobin and pyriproxyfen: 4936, included in scope from 2008, d) fenhexamid: 10461, included in scope from 2005.

As can be seen from **Figure 24**, imazalil was the pesticide most frequently found in foreign samples. Imazalil has been analysed in more than 12,000 samples and the frequency of 21% in foreign commodities is, therefore, calculated with high certainty. Imazalil is frequently used for post-harvest treatment of citrus fruits. Apart from imazalil, chlorpyrifos, thiabendazole, carbendazim (incl. benomyI), dithiocarbamates and chlormequat were found in more than 5% of the foreign samples. In Danish samples, azoxystrobin, chlormequat, dithiocarbamates, fenhexamid, propamocarb, pyraclostrobin, pyrimethanil, tolylfluanid and the sum of captan and folpet were found in more than 1% of the samples. The relatively high frequency of chlormequat detections was caused by 50 pear samples.

Pesticides found in cereals

The analyses of cereals covers barley (grit, malting, kernels), bulgur, corn flour, maize, millet, oat (bran, kernels, rolled oat), rice (brown, rice, white, wild, flour), rye (flour, bolted, kernels), spelt (flour, grain), wheat (bran, flour, kernels). In total, 1659 samples were analysed, 691 Danish produced and 998 samples of foreign origin. Pesticide residues were found in 34% of the samples.




Figure 25. *Frequencies of pesticides found in conventionally grown domestic and foreign samples of cereals. The pesticides were sought in approximately 1650 samples, except for the following: a) chlormequat and mepiquat: 1764. b) glyphosate: 1375 only sought in wheat, rye, oat and barley.*

As can be seen from **Figure 25**, chlormequat was the pesticide residue most often found in both Danish and foreign samples. Glyphosate was found frequently in Danish samples and more than one third of the Danish samples have residues of glyphosate. The pesticides, detection frequencies, and reporting limits are shown in Appendix 7.2.

Comparison of pesticides found in periods 1998-2004 with 2004-2011

The pesticides, detection frequencies, and reporting limits are shown in Appendix 7.2. Below is an evaluation of the findings and possible trends from the monitoring period 1998-2003 to this period.

The findings have been investigated to see whether there were any changes in the frequencies of pesticides from the previous monitoring period 1998-2003 and this period 2004-2011. Firstly, the pesticide profile has been expanded with 90 pesticides. This resulted in detection of 51 pesticides that was not included in the scope for the period 1998-2003). Additionally, 2 pesticides, atrazine and carbofuran, were analysed in both periods but only found in samples during the period 2004-2011. Thirty pesticides found in samples for the period 1998-2003 were not found in the period 2004-2011 (see **Table 5** and **Table 6**).

Table 7 shows the list of pesticides that were scarcely detected in the period 2004-2001 (>20 detection) and also scarcely detected in the period 1998-2003 and **Table 8** shows the pesticides that were not detected in the period 2004-2011.

Table 5. Pesticides detected in samples analysed in the period 2004-2011, but not in the period 1998-2003

| Pesticides analysed in both 1998-2003 and 2004-2011 | | |
|--|-------------------------|----------------|
| Atrazine | Carbosulfan | |
| Pesticides analysed only in 2004-2011 | | |
| Acetamiprid | Fenhexamid | Propamocarb |
| Aclonifen | Fluoxastrobin | Propoxur |
| Aldicarb | Fluroxypyr | Propyzamide |
| Benalaxyl | Flusilazole | Pymetrozine |
| Chlorthal-dimethyl | Flutriafol | Pyraclostrobin |
| Clofentezine | Hexaconazole | Pyridate |
| Cyromazine | Hexythiazox | Pyriproxyfen |
| Dichlorprop | Iprovalicarb | Quinoxifen |
| Diethofencarb | Linuron | Quizalofop |
| Diflufenican | Mecoprop | Spiroxamine |
| Dimethomorph | Methacrifos | Tebufenozide |
| Diniconazole | Methiocarb | Tebufenpyrad |
| Epoxiconazole | Methomyl and Thiodicarb | Tetraconazole |
| Ethoxyquin | Metribuzin | Triallate |
| Famoxadone | Oxamyl | Triflumuron |
| Fenazaquin | Oxydemeton-methyl | 2,4-D |
| Fenbuconazole | Pendimethalin | Pyridaben |

Table 6. Pesticides detected in samples analysed in the period 1998-2003, but not in the period 2004-2011

| Pesticides | | |
|-------------------|--------------|--------------------|
| Azinphos-ethyl | Etrimfos | Parathion |
| Biphenyl | Furathiocarb | Pentachloroanisole |
| Chinomethionat | Heptachlor | Pyrazophos |
| Chlorobenzilate | Isofenphos | Simazine |
| Chloropropylate | Mecarbam | Tetrasul |
| Demeton-S-methyl | Methoxychlor | |
| Dioxathion | Nuarimol | |

Table 7. Pesticides with less than 20 detection in the period 2004-2011 and similar detection frequencies in the period 1998-2003

| Pesticides | | |
|---------------------|-------------------|------------------|
| Aldrin and Dieldrin | Dichlorvos | Monocrotophos |
| Benfuracarb | Dicloran | Phorate |
| Binapacryl | Fenarimol | Propiconazole |
| Buprofezin | Fenpropathrin | Pyrazophos |
| Captafol | Fenpropimorph | Quinalphos |
| Carbaryl | Flucythrinate | Tecnazene |
| Carbofuran | Fluvalinate, tau- | Tolclofos-methyl |
| Chlorfenvinphos | HCH | Triazophos |
| Chlorpropham | Hexachlorobenzene | Trichlorfon |
| Cyfluthrin | Lindane | |
| DDT | Mevinphos | |

Table 8. Pesticides not detected in samples analysed in the period 2004-2011

| Pesticides | | |
|-----------------------------|-------------------------|---------------------------|
| 2-Naphthoxyacetic acid 1) | Endrin 1) | Nitrofen |
| 4-Chlorophenoxy acetic acid | Ethiofencarb | Nuarimol 2) |
| Acrinathrin | Etrimfos 2) | Ofurace |
| Azimsulfuron | Fenamiphos (sum) | Oxadixyl |
| Azinphos-ethyl 2) | Fenclorphos (sum) | Oxycarboxin |
| Bentazone (sum) | Fenpropidin 1) | Paclobutrazol |
| Bromophos 1) | Fluazifop-P-butyl (sum) | Parathion 2) |
| Bromophos-ethyl 1) | Flufenacet (sum) | Pentachloroanisole 2) |
| Bromoxynil | Flupyrsulfuron-methyl | Pentachlorothioanisole 1) |
| Bromuconazole (sum) | Fluquinconazole | Phenkapton 1) |
| Carbophenothion 1) | Flurtamone | Phosphamidon 1) |
| Carboxin | Flutolanil | Phoxim 1) |
| Chinomethionat 1) | Fonofos | Picolinafen |
| Chlorfenson 1) | Formothion 1) | Picoxystrobin |
| Chlorobenzilate 2) | Fuberidazole | Pirimiphos-Ethyl |
| Chloropropylate 2) | Furathiocarb | Propanil |
| Cinidon-ethyl (sum) | Heptachlor (sum) 2) | Propham |
| Clethodim | Heptenophos 1) | Proquinazid |
| Clodinafop-Propargyl | Iodosulfuron-methyl | Pyrazophos 2) |
| Cyproconazole | Isofenphos 2) | Pyridaphenthion |
| Demeton-S-Methyl 2) | Isofenphos-Methyl | Simazine 2) |
| Dialifos | Isoproturon | Sulfotep 1) |
| Dichlofenthion 1) | Jodfenphos 1) | TEPP 1) |
| Dimoxystrobin | Mecarbam 2) | Tetrasul 2) |
| Dinoterb | Methoxychlor 2) | Thiometon |

| Pesticides | | |
|-------------------|-------------|----------------|
| Dioxathion 2) | Metribuzin | Trifluralin |
| Ditalimfos 1) | Molinate | Triticonazole |
| DNOC | Monolinuron | Vamidothion 1) |

1) Analysed in the period 1998-2003, not detected

2) Analysed in the period 1998-2003 and detected

Table 9 shows increases or decreases of detection frequencies. It is not indicated if the frequencies have decreased or increased. This table includes pesticides that were often found in samples analysed in in the period 1998-2003 and/or the period 2004-2011. The table also includes data on the increase/decrease of the exposure in $\mu\text{g}/\text{person}/\text{day}$ for the pesticides. A significant increase/decrease in the frequency of a pesticide will not necessarily result in a significant increase/decrease in the exposure. This will be the effect if the detected levels of pesticides were low or high compared to the levels analysed earlier. The effect depends also on the commodities in which the pesticide is found.

Nine pesticides were found significantly more often in the period 2004-2011, namely acephate, azoxystrobin, bitertanol, bupirimate, chlorpyrifos-methyl, lambda-cyhalothrin, myclobutanil, tebuconazole and trifloxystrobin. The increase of detection of these pesticides resulted in an increase of the exposure, except for bitertanol. Also for difenoconazole, penconazole, prochloraz, profenofos and thiabendazole a clear increase in both frequencies and exposure were seen. For the rest of the pesticides in **Table 9**, the increases and decreases are not significant. However, the exposure for deltamethrin, kresoxim-methyl, pirimiphos-methyl, triadimefon+triadimenol and pyrimethanil was significantly higher (increased more than 200%) even though the frequencies were not.

Table 9. Trends in pesticides found and the resulting exposure from the period 1998-2003 to the period 2004-2011.

| Pesticide | Less than 20 detections in 1998-2003 | Less than 20 detections in 2004-2011 | Change in frequency of detection ¹⁾ | Change in exposure (µg/day) ¹⁾ |
|-------------------------|---|---|---|--|
| Acephate | * | | +++ | +++ |
| Azoxystrobin | * | | +++ | +++ |
| Bitertanol | * | | +++ | - |
| Bupirimate | * | | +++ | +++ |
| Chlorpyrifos-methyl | * | | +++ | ++ |
| Cyhalothrin, lambda- | * | | +++ | +++ |
| Myclobutanil | | | +++ | +++ |
| Tebuconazole | | | +++ | ++ |
| Trifloxystrobin | * | | +++ | +++ |
| Difenoconazole | * | | ++ | + |
| Penconazole | * | | ++ | ++ |
| Prochloraz | | | ++ | +++ |
| Profenofos | * | | ++ | + |
| Thiabendazole | | | ++ | + |
| Azinphos-methyl | | | + | ++ |
| Bifenthrin | | | + | - |
| Chlorpyrifos | | | + | + |
| Cypermethrin | | | + | + |
| Deltamethrin | | | + | +++ |
| Dicofol | | | + | - |
| Diphenylamine | | | + | + |
| Imazalil | | | + | - |
| Kresoxim-methyl | * | | + | +++ |
| Methamidophos | | | + | ++ |
| Pirimicarb | | | + | + |
| Pirimiphos-methyl | | | + | +++ |
| Triadimefon+triadimenol | | | + | +++ |
| Bromopropylate | | | - | - |
| Carbendazim and benomyl | | * | - | + |
| Chlormequat | | | - | - |
| Chlorothalonil | | | - | + |
| Cyprodinil | | | - | + |
| Diazinon | | | - | + |
| Dimethoate+omethoate | | | - | + |
| Dithiocarbamates | | | - | + |

| Pesticide | Less than 20 detections in 1998-2003 | Less than 20 detections in 2004-2011 | Change in frequency of detection ¹⁾ | Change in exposure (µg/day) ¹⁾ |
|------------------|---|---|---|--|
| Endosulfan | | | - | - |
| Ethion | | * | - | - |
| Fenitrothion | | | - | - |
| Fenthion | | * | - | - |
| Fludioxonil | | | - | + |
| Glyphosate | | | - | - |
| Iprodione | | | - | ++ |
| Malathion | | | - | - |
| Mepiquat | * | | - | + |
| Metalaxyl | | * | - | - |
| Methidathion | | | - | - |
| Parathion-methyl | | | - | - |
| Phosalone | | | - | - |
| Phosmet | | | - | + |
| Procymidone | | | - | - |
| Propargite | * | | - | - |
| Pyrimethanil | | | - | +++ |
| Quintozene | | | - | + |
| Tetradifon | | * | - | - |
| Tolyfluanid | | | - | - |
| Vinclozolin | | | - | - |

1) **+++** > 200%
++ 100-200%
+ 0-99%
- <0%

4.3 Exposure

Dietary exposure and risk assessment

The dietary exposure to pesticides has been calculated in order to assess the chronic (long-term) consumer health risk for the Danish population. To follow the trend in exposure over time, the exposure calculation has been calculated according to the approach of National Estimated Daily Intake given in “Guidelines for predicting dietary intakes of pesticides residue” (WHO, 1997).

The primary goal of the effort has been to assess whether the pesticide residues present in an average Danish diet is acceptable from a food safety point of view. For this reason, estimations that will give conservative (i.e. pessimistic) results were preferred to the contrary. If such estimates point to food safety problems, further refinements of the methods used must be performed to assess whether such problems are real, or just artefacts from a too simple model.

The dietary exposure to a pesticide residue in a given food was estimated by multiplying the residue level in the food by the amount of that food consumed. Residues were obtained from the Danish monitoring programme while consumption data were obtained from the Danish National Dietary Survey. The total dietary exposure to a given pesticide was estimated by summing the exposure for all food items containing residues of that pesticide. The exposure for each food item (i) is calculated, relative to the body weight (w) from the average residue concentration (C_i) in the food item, multiplied by the consumption (M_i) of the food item:

$$Exposure = \frac{C_1 * M_1 + C_2 * M_2 + C_3 * M_3 + \dots + C_i * M_i}{w}$$

In the risk assessment of the chronic exposure, the estimated total dietary exposure was compared with the toxicological reference value, Acceptable Daily Intake (ADI). ADI is the estimated maximum amount of a substance to which an individual in a (sub) population may be exposed daily over its lifetime without appreciable health risk (from WHO). A more detailed description of the exposure calculations can be found in Annex 6.1.

Consumption data

Exposure estimates were based on consumption data obtained from the Danish National Dietary Survey 2003-2008 (Pedersen et al., 2010). This cross-sectional survey included 2700 participants aged 4-75 years old drawn from the Danish Central Person Register. The participants can be characterised as close to representative for the Danish population. In this report, the consumer groups, children aged 4-6 years and adults, women and men aged 15-75 years, will be used.

The consumption data used for the exposure calculations for different consumer groups are shown in Appendix 7.4. It is not possible from the consumption data, to distinguish between consumption of commodities of Danish and foreign origin. Therefore, the distribution between domestic and foreign as well as the distribution between the foreign food commodities has been assumed to follow the distribution of samples in the monitoring programmes.

Residue data

As previously described, the residue data included a total of 17309 samples. The pesticides in the monitoring programme were found in about 130 different commodities.

An average content has been calculated for each combination of pesticide, commodity and origin (domestic or foreign). Only combinations of pesticide/commodity/origin with at least one detectable residue above LOR were included in the calculations of exposure.

In many circumstances, no detectable amount of pesticides was found, but this does not necessarily mean that the content is zero. The content may just be too low for detection with the available methods, or in other words, below the level of reporting. No exact method for estimation of the contribution from such non-detects exists. Different methods can be applied and it has been the most common to use zero for non-detects. In more and more circumstances, the exposure is, however, calculated using different assumptions e.g. zero and $\frac{1}{2}$ LOR for non-detects to indicate a lower bound and upper bound for the exposure (ANSES, 2011).

In some cases, correcting for samples without detected residues leads to a very high correction. This is, especially the case when many samples have been analysed, but only a few samples with residues were detected. Therefore a model was chosen where the difference between the dietary exposure with, and without, correction for non-detects is not allowed to be more than a factor of 25. This means that if the correction is more than a factor of 25 it is adjusted down to 25. The background for the correction factor of 25 is described in Annex 6.2.

To obtain the most realistic picture of the exposure to pesticides, it is also important to address processing factors. The pesticides included in the calculations were found in commodities such as citrus fruits, banana, water melon, and pineapples, for which processing factors for peel/pulp distribution are normally applied in a refined exposure calculation. For comparison two scenarios were performed: one where no processing factors were included and another, where processing factors were included. The processing factors used in this report were the same as those used in the report for the monitoring period 1998-2003 (Poulsen et al., 2004). For thiabendazole and the benomyl group a processing factor of 0.25 was used, while a factor of 0.1 was used for all the other pesticides (see Appendix 7.6).

Estimation of the cumulative exposure and risk assessment

There is no internationally agreed method for risk assessment of the cumulative exposure to multiple residues of pesticides. In the present report, assessments of exposure to a mixture of pesticides were performed by using the Hazard Index (HI) method as described by US EPA (1986a), Wilkinson et al. (2000), Reffstrup et al. (2010), and Kortenkamp et al. (2012). The HI is the sum of Hazard Quotients (HQ) for the individual pesticides. The HI method assumes that the effects following cumulative exposure can be predicted by the mathematical model of dose-addition (Wilkinson et al., 2000) and is designed for risk assessment for substances which have the same effect or common mode of action, e.g. the organophosphor pesticides or the triazole group. As the HI method assumes the same kind of adverse effect for all the detected pesticides, it is a relatively conservative (precautionary) approach for cumulative risk assessment. However, as described by Kortenkamp et al. (2012) and Reffstrup et al. (2010) the method can be used even when the substances have dissimilar toxicological endpoints. The results should then be interpreted with caution.

The Hazard Quotient (*HQ*) for a given pesticide is calculated from the total exposure from the diet to that pesticide divided by the ADI for that pesticide:

$$\text{Hazard Quotient (HQ)} = \frac{\text{Exposure}}{\text{ADI}}$$

The Hazard Index (HI) for a given diet is calculated by summing the Hazard Quotients (HQ) for each pesticide (p) in the diet:

$$\text{Hazard Index (HI)} = \text{HQ}_1 + \text{HQ}_2 + \text{HQ}_3 + \dots + \text{HQ}_p$$

Throughout the present report, the term HI will be used for the sum of HQs from the pesticides that the consumer is exposed to. If the HI exceeds 1 (or 100%), the mixture has exceeded the maximum acceptable level and there might thus be a risk (Reffstrup et al., 2010).

The ADIs used for calculation of the HQs for the individual pesticides are those accepted in the EU (Commission (COM) or EFSA) or by JMPR when available (see Appendix 7.5). For pesticides where an accepted ADI was not available, the ADI proposed by the Rapporteur Member State in the Draft Assessment Report (DAR) has been used for calculation of the HQ. No ADIs were available for about ten pesticides and therefore, these substances have not been included in the cumulative risk assessment based on HI.

Results and discussion of exposure calculations

In **Table 10** is shown the average exposure in $\mu\text{g}/\text{kg}$ bw/day and in $\mu\text{g}/\text{day}/\text{person}$ for the adult population using different assumptions in the calculations. The exposure for adults is between $0.9 \mu\text{g}/\text{kg}$ bw/day (correction for processing, no correction for non-detects) and $3.0 \mu\text{g}/\text{kg}$ bw/day (no correction for processing, all non-detects = $\frac{1}{2}$ LOR). For children, the same figures are between 2.0 and $6.6 \mu\text{g}/\text{kg}$ bw/day.

Table 10. Average exposure for the consumer groups “Adults” (15-75 years) and “Children” (4-6 years) using different models (LOR: Limit of reporting)

| Correction for undetected residues: | No correction | Exposure | |
|-------------------------------------|---------------|--|-----------------------------------|
| | | $\frac{1}{2}$ LOR for non-detects; Correction factor limited to 25 ($\mu\text{g}/\text{kg}$ bw/day) | $\frac{1}{2}$ LOR for non-detects |
| Adults, no reduction for peeling | 1.4 | 2.6 | 3.0 |
| Children, no reduction for peeling | 2.9 | 5.7 | 6.6 |
| Adults | 0.9 | 1.9 | 2.3 |
| Children | 2.0 | 4.5 | 5.3 |
| ($\mu\text{g}/\text{day}$) | | | |
| Adults, no reduction for peeling | 105 | 192 | 222 |
| Children, no reduction for peeling | 62 | 124 | 144 |
| Adults | 68 | 146 | 171 |
| Children | 44 | 98 | 116 |

In **Table 11** is shown the HI for adults (15-75 years) and children (4-6 years) using different assumptions in the calculations. For adults, the HI is calculated to be between 4% and 44% and for children, the HI is between 10% and 124%. As can be seen from **Table 10**, using

½LOR in the calculations has a very big impact on the exposure, as well as on HI. It can also be seen that using ½LOR has a higher impact on the exposure and HI, than including processing factors.

For children, the worst case calculation leads to a HI above 100%. However, as mentioned above, an examination of the details leading to this result shows clear indications of over-correction for non-detected residues (see Annexes 6.1 and 6.2). Using Model 3 in Annex 6.1 (with limiting factor 25 on corrections for non-detects), HI for children has been calculated to be 56% (44% with reduction for processing), which is below the critical value of 100%.

Table 11. Hazard Index (HI) for the consumer groups “Adults” (15-75 years) and “Children” (4-6 year) using different models (LOR: Limit of reporting)

| Correction for undetected residues: | No correction | Hazard Index | |
|-------------------------------------|---------------|---|----------------------|
| | | ½LOR for non-detects; Correction factor limited to 25 | ½LOR for non-detects |
| Adults, no reduction for peeling | 7% | 23% | 49% |
| Children, no reduction for peeling | 14% | 56% | 124% |
| Adults | 4% | 18% | 42% |
| Children | 10% | 44% | 108% |

As mentioned above it was evaluated that if ½LOR was used in the calculations, the most suitable model for this was the model where corrections for non-detects were limited to a factor of 25. In this case, it was taken into consideration that levels below the LOR could be higher than zero. At the same time, it was taken into consideration that a few positives will not have an unrealistic high impact on the exposure; e.g. that 2-3 residues above the LOR out of maybe 400 samples would mean that 397-398 samples were calculated as having contents of ½LOR.

Furthermore, it has been evaluated that the most realistic exposure calculation was obtained by using processing factors in the calculation. A processing factor of 0.1 has been used for most of the relevant pesticide/crop combinations. This represents some uncertainty because processing factors will be different depending on the pesticide/crop combination.

Unless otherwise stated, the calculations in this report have been performed using the model: corrections for non-detects were limited to a factor of 25 and processing factors were included.

In **Table 12** is shown the average exposure in µg/kg bw/day and in µg/person/day and the Hazard Index for the consumer groups ‘children, adults, men, and women’ using the chosen model. The results show that children have the highest exposure per kg bw followed by women, adults and men. The reason for the highest exposure for children is that they consume relatively more per kg bodyweight, compared to adults. Due to the fact that women consume more fruit and vegetables than men (see Appendix 7.4) they have a higher exposure than men. Since the exposure per kg bw is highest for children, the HI is also highest for children. It is seen from **Table 12** that the HI for children is more than twice as high as for women and about 3 times higher than for men, but still well below 100%.

Table 12. Average exposure and Hazard Index (HI) for adults, men, women and children (Correction for non-detected residues; correction factor limited to 25).

| | Exposure (µg/kg bw/day) | Exposure (µg/person/day) | Hazard Index |
|---------------------|----------------------------|-----------------------------|--------------|
| Adults, 15-75 years | 1.9 | 146 | 18% |
| Men, 15-75 years | 1.6 | 134 | 14% |
| Women, 15-75 years | 2.2 | 151 | 20% |
| Children, 4-6 years | 4.5 | 98 | 44% |

Commodities that contributes most to the exposure

The contribution of each commodity to the exposure as well as the HI has been calculated for the consumer group “Adults” and “Children”. In Appendix 7.7 and 7.8 details are shown for individual commodities. In **Table 13**, the exposure and HI is shown for the 25 commodities that contribute most to the exposure for the consumer group “Adults”. The exposure is shown both in µg/kg bw/day and in µg/day. As can be seen, the sum of HI constitutes about 97% of the total HI and about 95% of the total exposure. It should be noted that the consumption of a specific commodity can have a big impact on the HI. Thus, the HI can be high for a commodity consumed in high amounts even though the concentrations of the pesticides in this commodity are lower, compared to the concentrations of the pesticides found in other commodities. For example, for the consumer group “Adults”, table grapes contribute to the HI by 0.74%, which is about 2.7 times more than the contribution from kiwi (0.27%). The consumption of table grapes is higher than for kiwis. If the consumption of table grapes and kiwis was the same then kiwis would contribute about 50% more to the HI than table grapes.

Table 13. Exposure and Hazard Index (HI) for the group "Adults" and the 25 commodities that contribute most to the Hazard Index.

| Commodity | Exposure (µg/kg bw/day) | Exposure (µg/person/day) | Hazard Index |
|-------------------|----------------------------|-----------------------------|--------------|
| Apples | 0.57 | 43 | 6.9% |
| Wheat flour | 0.22 | 16 | 1.3% |
| Tomatoes | 0.15 | 11 | 1.2% |
| Pears | 0.14 | 11 | 1.15% |
| Carrots | 0.037 | 2.7 | 1.0% |
| Table grapes | 0.1 | 8.4 | 0.74% |
| Cucumbers | 0.053 | 4.1 | 0.69% |
| Wine, red | 0.11 | 8.4 | 0.48% |
| Rye flour | 0.032 | 2.4 | 0.44% |
| Peppers, sweet | 0.033 | 2.5 | 0.31% |
| Nectarines | 0.031 | 2.3 | 0.30% |
| Peaches | 0.032 | 2.4 | 0.28% |
| Oranges | 0.030 | 2.2 | 0.28% |
| Kiwi | 0.044 | 3.3 | 0.27% |
| Lettuce | 0.063 | 4.8 | 0.26% |
| Pineapples | 0.020 | 1.5 | 0.24% |
| Peas without pods | 0.0084 | 0.63 | 0.24% |

| Commodity | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | Exposure ($\mu\text{g}/\text{person}/\text{day}$) | Hazard Index |
|------------------------|---|--|--------------|
| Mandarins, clementines | 0.030 | 2.2 | 0.21% |
| Potatoes | 0.055 | 4.1 | 0.17% |
| Plums | 0.017 | 1.3 | 0.12% |
| Beans with pods | 0.0052 | 0.39 | 0.12% |
| Teas | 0.0047 | 0.35 | 0.12% |
| Strawberries | 0.023 | 1.8 | 0.11% |
| Melons | 0.0051 | 0.39 | 0.08% |
| Oranges, juice | 0.016 | 1.2 | 0.07% |
| Sum | 1.8 | 138 | 17% |
| Total | 1.9 | 146 | 18% |
| % of total | 95% | 95% | 97% |

Figure 26 shows the nine commodities that contribute most to the exposure for “Adults” together with the contributions from the rest of the commodities (“Others”).

Figure 27 shows the same for the commodities that contributes most to the HI for “Adults”. The seven commodities, apples, wheat flour, tomatoes, pear, table grapes, red wine and cucumber contribute most to both exposure and HI. Carrots and rye flour are in the ‘top nine’ when comparing HI, while lettuce and potatoes only are in the ‘top nine’ when comparing exposure. This difference reflects the different types of pesticides (with different ADIs) that were found in the different commodities. Apples contribute most to both the exposure and to HI.

Pesticides that contributes most to exposure

The contribution of each pesticide to the exposure, as well as the HI has been calculated using the chosen model. In Appendices 7.8 and 7.9, details are shown for individual pesticides. The HQs for the individual pesticides ranged from 0.00001% to 2.4% with most of the HQs being below 1% (see Appendix 7.9) indicating that there is no risk of adverse effects following exposure to the individual pesticides. Furthermore, the HI of 18% for adults and 44% for children is not considered to indicate a risk of adverse effects following a cumulative exposure to all the detected pesticides.




Figure 26: Relative contribution of *commodities* to total *exposure* to pesticide residues in the diet. Consumer group: Adults; estimated total exposure: 1.9 µg/kg bw/day. “Other” represents 78 different commodities.




Figure 27: Relative contribution of *commodities* to **Hazard Index** for pesticide residues in the diet. Consumer group: Adults; estimated diet Hazard Index: 18%. “Other” represents 78 different commodities.

In **Table 14**, the HQs and exposure for the 20 pesticides that have the highest HQ for the consumer group “Adults” are shown. The exposures are shown, both in $\mu\text{g}/\text{kg bw}/\text{day}$ and in $\mu\text{g}/\text{day}$. The sum of HQ constitutes about 72% of the total HI and about 33% of the total exposure. There is a big difference in ordering the pesticides according to exposure or HQ, which is due to differences in their ADI; e.g. as shown in **Table 14**, imazalil contributes most to the exposure, while diazinon has the highest HQ. For some pesticides, the exposure is calculated to be higher than for the pesticides shown in **Table 14**, but the HQ is so low that this pesticide is not among the 20 pesticides with highest HQ values; e.g. the exposure for pyrimethanil is $3.18 \mu\text{g}/\text{person}/\text{day}$, but the HQ is only 0.028%.

Table 14. Exposure and Hazard Quotients (HQ) for the group "Adults" and the 20 pesticides that contribute most to the Hazard Index.

| Pesticide name | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | Exposure ($\mu\text{g}/\text{person}/\text{day}$) | Hazard Quotient |
|-------------------------|---|--|-----------------|
| Diazinon | 0.0047 | 0.35 | 2.4% |
| Omethoate | 0.0039 | 0.29 | 1.3% |
| Pirimiphos-methyl | 0.043 | 3.2 | 1.1% |
| Dicofol (sum) | 0.017 | 1.3 | 0.86% |
| Phosmet (sum) | 0.021 | 1.6 | 0.69% |
| Procymidone | 0.018 | 1.34 | 0.64% |
| Dimethoate | 0.0063 | 0.48 | 0.63% |
| Propargite | 0.061 | 4.6 | 0.61% |
| Carbaryl | 0.043 | 3.2 | 0.57% |
| Chlorfenvinphos | 0.0028 | 0.21 | 0.57% |
| Azinphos-methyl | 0.022 | 1.7 | 0.44% |
| Carbendazim and benomyl | 0.087 | 6.5 | 0.43% |
| Dithiocarbamates | 0.21 | 16 | 0.42% |
| Linuron | 0.010 | 0.76 | 0.34% |
| Methomyl and thiodicarb | 0.0083 | 0.62 | 0.33% |
| Bitertanol | 0.0090 | 0.67 | 0.30% |
| Methamidophos | 0.0029 | 0.22 | 0.29% |
| Imazalil | 0.072 | 5.4 | 0.29% |
| Oxamyl | 0.0026 | 0.19 | 0.26% |
| Oxydemeton-methyl (sum) | 0.00074 | 0.055 | 0.25% |
| Sum ¹ | 0.65 | 48 | 13% |
| Total | 1.9 | 146 | 18% |
| % of total | 33% | 33% | 72% |

In **Figure 28**, is shown the 9 pesticides that contribute most to the exposure together with the contributions from the rest of the commodities (“Others”).

Figure 29 shows the same for the pesticides that contribute most to the HI.

¹ Summing has been performed using more decimals on individual exposures/HQs that shown in the table.




Figure 28: Relative contribution of pesticides to total exposure to pesticide residues in the diet. Consumer group: Adults; estimated total exposure: 1.9 µg/kg bw/day. “Other” represents 148 different pesticides.




Figure 29: Relative contribution of pesticides to Hazard Index for pesticide residues in the diet. Consumer group: Adults; estimated Hazard Index: 19%. “Other” represents 148 different pesticides.

In contrast to what was the case for commodities, the pesticide in the two ‘top nine’ groups are all different. The ‘top nine’ pesticides in both cases account for approximately half of the exposure (or HI), while the ‘top nine’ commodities accounted for approximately 75%.

Effect from origin of commodities on exposure to pesticide residues

Some commodities originate only from foreign countries e.g. oranges, while some commodities, e.g. apples, are produced both in Denmark and in foreign countries. As seen in **Figure 2** and **Figure 3**, fruit and vegetables from the EU and third countries have a higher frequency of pesticide residues compared to fruit and vegetables from Denmark. Therefore, the origin of the commodities can have an effect on the exposure to pesticides in $\mu\text{g}/\text{day}$, as well as the HI.

The exposure to pesticides has also been calculated assuming that the consumers eat commodities of Danish origin whenever possible, e.g. only Danish apples and pears, but oranges from other countries. The results of such calculation are shown in **Table 15**, both with regards to exposure and HI. The results show that both the average exposure per day and HI then decrease compared to eating commodities of both Danish and foreign origin. Since Danish commodities in general have lower contents and lower frequencies of pesticide residues compared to commodities of foreign origin, the average exposure can be expected to be lower when Danish commodities are eaten whenever possible. However, because of the use of $\frac{1}{2}\text{LOR}$ the correction for commodities of foreign origin may be overestimated in some cases. If a single residue was found in a commodity from a single country all residues of this commodity were corrected, independent of the country of origin.

Effect on exposure for consumers having a high consumption of fruit and vegetables

In Denmark, the Danish Food and Veterinary Administration recommend that everyone above 10 years of age, eat 600 g of fruit and vegetables per day. Calculations have been performed to investigate what effect a high consumption of fruits and vegetables has on the exposure to pesticides for the two consumer groups, men and women. The calculations were performed by taking all the participants in the food dietary survey who consumed more than 550 gram of fruits and vegetables (excluding potatoes) and calculate the exposure for these persons (high consumers). **Table 15** shows the exposure and HI for the high and the average consumers using the chosen model. It is observed that the exposure increases, about a factor of $1\frac{1}{2}$ -2 for both women and men. For the high consumers, men had the highest exposure, whereas, for the average consumers, it was women.

Table 15. Exposure and Hazard Index (HI) for normal consumers and for consumers with high consumption of fruit and vegetables (above 550 g/day excluding potatoes) as well as consumers that choose Danish grown commodities whenever possible^{a)}.

| Consumer group | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | Exposure ($\mu\text{g}/\text{day}$) | Hazard Index |
|---|---|--|--------------|
| Adults, average consumption | 1.9 | 146 | 18% |
| Adults, average consumption, domestic preferred | 1.0 | 76 | 8% |
| Men, average consumption | 1.6 | 134 | 14% |
| Men, average consumption, domestic preferred | 0.8 | 67 | 6% |
| Men, high consumption | 3.19 | 261 | 29% |
| Men, high consumption, domestic preferred | 1.5 | 125 | 13% |
| Women, average consumption | 2.2 | 151 | 20% |
| Women, average consumption, domestic preferred | 1.2 | 80 | 10% |
| Women, high consumption | 3.5 | 240 | 33% |

| Consumer group | Exposure (µg/kg bw/day) | Exposure (µg/day) | Hazard Index |
|---|----------------------------|----------------------|--------------|
| Women, high consumption, domestic preferred | 1.8 | 125 | 16% |
| Children, average consumption | 4.5 | 98 | 44% |
| Children, domestic preferred | 2.4 | 52 | 20% |

a) i.e. commodities where domestic samples have been analysed (see Appendix 7.4)

Comparison of exposure and Hazard Index for the two periods 1998–2003 and 2004–2011

The exposure has also been calculated in the reports for the two monitoring periods 1993–1997 and 1998–2003 (DVFA, 2000 and Poulsen et al., 2005), while the HI has only been calculated in the report for the period 1998–2003, and in the present report. The model for exposure calculation is the same in the present report as in the report for the previous period 1998–2003. Because the model for exposure calculation was not the same for the first period and, because of a large increase in the number of pesticides analysed, it was decided only to make a comparison for the exposure and HI for the two periods 2004–2011 and 1998–2003.

The model for exposure calculation was the same for the two periods, but the number of pesticides, reporting limits, commodities with consumer data and ADIs were not quite the same for the two periods:

Pesticide profile: The present study included and detected a higher number of pesticides. Since 2003, the number of pesticides in the monitoring programme has increased from 153 pesticides to about 275 pesticides including metabolites.

Reporting limits: In some cases reporting limits were different between the two periods. Some were lower, some were higher. This might influence the differences between the two periods.

Commodities with consumptions data: The present study included a higher number of commodities with consumption, about 130 commodities, while only about 90 commodities were included for the previous period. However, the 25 commodities that contributed most to the exposure were included in both periods. Consumption data from the survey performed in the period 2003–2008 were used in the present study while consumption data from the period 2000–2002 were used in the previous period.

ADIs: the ADIs for the substances have changed for some of the substances.

Age groups: Estimation of exposure was performed for children aged 4–14 in the report from the period 1998–2003, while the calculation was performed for children aged 4–6 in this study. Children eat relatively more per kg bodyweight compared to adults, hence children aged 4–6 have a higher exposure per kg bodyweight compared to children aged 4–14.

Table 16 shows the average exposure for men, women and children for both periods.

Table 16 Comparison of the average intake and HI for men, women and children using the chosen model for the two monitoring periods 1998-2003 and 2004-2011

| | Exposure (µg/kg bw/day) | Exposure (µg/day) | Hazard Index |
|----------------------|----------------------------|----------------------|--------------|
| Exposure, 2004-2011 | | | |
| Men, 15-75 years | 1.6 | 134 | 14% |
| Women, 15-75 years | 2.2 | 151 | 20% |
| Children, 4-6 years | 4.5 | 98 | 44% |
| Exposure, 1998-2003 | | | |
| Men, 15-75 years | | 124 | 19% |
| Women, 15-75 years | | 137 | 26% |
| Children, 4-14 years | | 103 | 35% |

For adults a minor increase in the calculated exposure and a decrease in Hazard Index were observed. However, due to the uncertainties and the different basis for the calculations in the two periods a clear conclusion on the changes cannot be drawn.

For children the calculations have been performed for two different age groups in the two periods, and therefore it is not possible to compare the results.

It should be noted that a detailed analysis has not been performed to clarify the observed differences.

However, in relation to these comparisons it should be emphasised that the HI was below 100% for all consumer groups in both periods

Table 17. Hazard Quotients (HQ) for the 20 pesticides that contribute most to the Hazard Index for the consumer group “Adults”, present vs. last periode.

| Pesticide name | Hazard Quotient 2004-2011 | Hazard Quotient ^{a)} 1998-2003 | Status under 1107/2009 |
|------------------------------|------------------------------|--|---------------------------|
| Diazinon | 2.4% | | Not approved (2007) |
| Omethoate | 1.3% | 0.49% | Not approved (2002) |
| Pirimiphos-methyl | 1.1% | | Approved |
| Dicofol (sum) | 0.86% | 2.7% | Not approved (2008) |
| Phosmet (sum) | 0.69% | 0.20% | Approved |
| Procymidone | 0.64% | | Not approved (2009) |
| Dimethoate | 0.63% | 0.49% | Approved |
| Propargite | 0.61% | 1.4% | Not approved (2008) |
| Carbaryl | 0.57% | 0.23% | Not approved (2007) |
| Chlorfenvinphos | 0.57% | 1.16% | Not approved (2002) |
| Azinphos-methyl | 0.44% | | Not approved (2005) |
| Carbendazim and beno- myl | 0.43% | 0.37% | Approved |
| Dithiocarbamates | 0.42% | 1.2% | Some approved |
| Linuron | 0.34% | | Approved |
| Methomyl and Thiodicarb | 0.33% | | Approved |
| Bitertanol | 0.30% | | Approved |

| Pesticide name | Hazard Quotient 2004-2011 | Hazard Quotient ^{a)} 1998-2003 | Status under 1107/2009 |
|-------------------------|------------------------------|--|---------------------------|
| Methamidophos | 0.29% | | Not approved (2008) |
| Imazalil | 0.29% | | Approved |
| Oxamyl | 0.26% | | Approved |
| Oxydemeton-methyl (sum) | 0.25% | | Not approved |

^{a)} Consumer group "All" (4-75 years)

Table 17 shows the 20 pesticides which had the highest HQ in the period 2004-2011. Some of the pesticides were also among the 20 pesticides that have the highest HQ in the period 1998-2003. For comparison the results from this period are also shown in the table. It is seen that the HQ for most of the non-approved substances is less in the present period, compared to the previous period. Dimethoate and omethoate are reported separately in this report, while they were reported as a sum in the previous report. Omethoate is not allowed to be used in the EU, but it is a metabolite of dimethoate, hence the relatively high HQ of omethoate is most likely due to the use of dimethoate. A large contribution from an eventual misuse of omethoate is not likely, because it is difficult to acquire the substance. Most of the substances, which have been "not-approved", quite recently, still contribute to the average exposure for the period 2004-2011. However, it is expected that the HQ for these substances will decrease further in the future.

Conclusion for exposure

The average exposure to pesticides for the group "Adults" was calculated to be 146 µg/day/person, using Model 3 (non-detects assumed to be ½LOR, but limiting correction to 25 times the value from Model 1 (non-detects assumed to be zero)). This model takes into account both processing factors and that non-detects could have contents above zero. Women have the highest average exposure (151 µg/day/person), compared to men and children (134 µg/day/person and 98 µg/day/person, respectively), as they eat more fruit and vegetables.

Consumers (men and women) eating more than 550 g of fruit and vegetables per day have an intake that was higher than the average exposure, namely 261 µg/day/person compared to 134 µg/day/person for men and 240 µg/day/person compared to 151 µg/day/person for women. The relative increase in exposure was independent of the model chosen; for both Model 1 and Model 3, the exposure as well as Hazard Index increased 155-170% for women and 190-210% for men, compared to the values for the average diets.

The choice of domestic vs. foreign grown commodities has a significant influence on the exposure. For all groups ("Adults", "Men", "Women", "Children", "High consumers" (male or female)), using either Model 1 or Model 3, the exposure as well as the Hazard Index was reduced to 45-55% of the mixed selection¹.

The risk assessment of the cumulative exposure was performed by the Hazard Index method. The Hazard Quotient was calculated for each pesticide and then summed into a so-called

¹ Exposure calculated using residue levels from domestic grown commodities for those commodities where domestic samples were analysed and residue levels for foreign samples otherwise vs. domestic and foreign grown commodities selected with the same distribution as the distribution of the analysed samples.

Hazard Index, (HI). The HQs for the individual pesticides range from 0.00001% to 2.4% with most of the HQs (98%) being below 1% (see Appendix 7.9) indicating that there is no risk of adverse effects following exposure to the individual pesticides. The HI for the group “Adults” was 18% with the chosen model. The HI was highest for children (44%), compared to women (20%) and men (14%). For consumers (men and women) eating more than 550 g fruit and vegetable per day, the HI increased from 14% to 29% for men, while for women the HI increased from 20% to 33%. The HI of 18% for adults and 44% for children is not considered to indicate a risk of adverse effects following a cumulative exposure to all the detected pesticides. The HI method assumes the same kind of adverse effect for all the detected pesticides and therefore it is a relatively conservative (precautionary) approach for cumulative risk assessment but the method was used here to give an indication of whether there is a risk with the cumulative exposure or not. Furthermore the method gives some indications of the commodities and pesticides that contribute most to the risk.

About 95% of the HI or exposure was accounted for by 25 different commodities. Regarding pesticides, the exposure and HI are distributed between many substances.

Summarising the results of the exposure and the HI it can be concluded that according to our present knowledge there is no reason to be concerned about pesticide exposure for the Danish population even for consumers (adults) who eat more than 550 g of fruit and vegetables each day.

On the other hand, the exposure to pesticide residues from the food should not be ignored and the basis for exposure calculations for the Danish consumers should be further improved by:

- 1) Expanding the number of pesticides in the monitoring programme.
- 2) Increasing the sensitivity of the analytical methods in order to minimize the numbers of samples with undetected residues.
- 3) Providing detailed dietary information for more commodities.

5 References

- Andersen J.H. 2011. Implementation of Electronic Transmission of Chemical Occurrence Data (CFP/EFSA/DATEX/2009/01) in Denmark. Scientific/technical report submitted to EFSA, Question No EFSA-Q-2009-00838. Available online: <http://www.efsa.europa.eu/en/supporting/pub/152e.htm>
- ANSES, 2011, Pesticide residues, additives, acrylamide and polycyclic aromatic hydrocarbons. Available from: <http://www.tds-exposure.eu/sites/default/files/WP1/RapportEAT2EN2.pdf>
- Christensen H.B., Petersen A., Poulsen, M., Grossmann A., Holm M., Pesticidrester i Fødevarer, (in Danish) 2005. Danish Veterinary and Food Administration. Available online: <http://www.foedevarestyrelsen.dk/Publikationer/Alle20publikationer/2006023.pdf>
- Christensen H.B., Hermann S.S., Petersen A., Poulsen, M., Lind R.R., Holm M., Cederberg D.L. Pesticidrester i Fødevarer, (in Danish) 2006. Danish Veterinary and Food Administration. Available online: <http://www.foedevarestyrelsen.dk/Publikationer/Alle20publikationer/2007016.pdf>
- Christensen H.B., Hermann S.S., Petersen A., Poulsen, M., Jensen B.H., Holm M. Pesticidrester i Fødevarer, (in Danish) 2007. Danish Veterinary and Food Administration. Available online: <http://www.foedevarestyrelsen.dk/Publikationer/Sider/publikationDetalje.aspx?pub=2008003>
- Danish Veterinary and Food Administration, Production aids, Monitoring system for foods, 1993-1997. Part 3. FødevareRapport 2000:03 (January 2000).
- European Commission, 2012. Commission Implementing Regulation (EU) No 788/2012 of 31 August 2012 concerning a coordinated multiannual control programme of the Union for 2013, 2014 and 2015 to ensure compliance with maximum residue levels of pesticides and to assess the consumer exposure to pesticide residues in and on food of plant and animal origin. EU Journal L235/8, Available online: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2012:235:0008:0027:EN:PDF>
- European Food Safety Authority (EFSA); Management of left-censored data in dietary exposure assessment of chemical substances. EFSA Journal 2010; 8(3) [96 pp.]. doi:10.2903/j.efsa.2010.1557. Available online: <http://www.efsa.europa.eu/en/efsajournal/pub/1557.htm>
- European Food Safety Authority; 2009 EU Report on Pesticide Residues. EFSA Journal 2011; 9(11):2430. [225 pp.]. doi:10.2903/j.efsa.2011.2430. Available online: <http://www.efsa.europa.eu/en/efsajournal/pub/2430.htm>
- European Union, 2002, Commission Directive 2002/63/EC of 11 July 2002 establishing Community methods of sampling for the official control of pesticide residues in and on

products of plant and animal origin and repealing Directive 79/700/EEC; EU Journal L187/30, published 16.7.2002

European Union, 2005, Regulation (EC) No 396/2005 of the European Parliament and of the Council of 23 February 2005 on maximum residue levels of pesticides in or on food and feed of plant and animal origin and amending Council Directive 91/414/EEC; EU Journal L70/1, published 16.3.2005

Environmental Protection Agency (EPA). 1986b. Guidelines for the Health Risk Assessment of Chemical Mixtures. 51 Federal Register 34014 (September 24, 1986).

Jensen B.H., Petersen A., Christensen H.B., Andersen J.H., Hermann S.S., Poulsen, M., Hilbert G., Grossmann A., Holm M., Pesticidrester i Fødevarer, (in Danish) 2009. Danish Veterinary and Food Administration. Available online:
<http://www.foedevarestyrelsen.dk/Publikationer/Alle20publikationer/2010003.pdf>

Jensen B.H., Christensen H.B., Andersen J.H., Petersen A., Hilbert G., Grossmann A., Madsen H.L. Pesticidrester i Fødevarer, (in Danish) 2010. Danish Veterinary and Food Administration. Available online:
http://www.foedevarestyrelsen.dk/Publikationer/Alle_publicationer/2011/002.htm

Jensen B.H., Petersen A., Andersen J.H., Hilbert G., Grossmann A., Holm M., Pesticidrester i Fødevarer, (in Danish) 2011. Danish Veterinary and Food Administration. Available online:
<http://www.foedevarestyrelsen.dk/Publikationer/Alle20publikationer/2012002.pdf>

Kortenkamp, A., Evans, R., Faust, M., Kalberlah, F., Scholze, M., and Schuhmacher-Wolz, U. Investigation of the state of the science on combined actions of chemicals in food through dissimilar modes of action and proposal for science-based approach for performing related cumulative risk assessment. Supporting Publications 2012:EN-232. [233 pp.]. Available online: www.efsa.europa.eu/publications

Pesticidrester i fødevarer – resultater for 2004. Danish Veterinary and Food Administration. Available online:
http://www.foedevarestyrelsen.dk/SiteCollectionDocuments/25_PDF_word_filer20til20download/05kontor/Pesticidresterifdevarer2004.pdf

Pedersen, A.N., Fagt, S., Groth, M.V., Christensen, T., Biloft-Jensen, A.P., Matthiessen, J., Andersen, N.L., Kørup, K., Hartkopp, H.B., Ygil, K.H., Hinsch, H-J., Saxholt, E., Trolle, .E, 2010, Danskernes kostvaner 2003 – 2008, Hovedresultater (Dietary habits in Denmark 2003-2008, Main results), National Food Institute, Technical University of Denmark

Petersen A., Jensen B.H., Christensen H.B., Andersen J.H., Hjort K., Poulsen, M., Holm M., Hilbert G. Pesticidrester i Fødevarer, (in Danish) 2008. Danish Veterinary and Food Administration. Available online:
<http://www.foedevarestyrelsen.dk/Publikationer/Alle20publikationer/2009006.pdf>

- Poulsen, M., Poulsen, Andersen J.H., Petersen A., Hartcopp H., 2005. *Food Monitoring Program 1998-2003, Part 2 Pesticides*. (Copenhagen; Danish Veterinary and Food Administration). Available online: http://www.dfvf.dk/Files/Filer/Pesticid/Monitoring_1998-2003.pdf
- Refstrup T. K., Larsen C. J., Meyer O., 2010. Risk assessment of mixtures of pesticides. Current approaches and future strategies. *Regulatory toxicology and Pharmacology* 56, 174-192.
- WHO, 1997. *Guidelines for Predicting Dietary Intake of Pesticide Residues*, 2nd Revised Edition, Document WHO/FSF/FOS/97.7, World Health Organization, Geneva
- Wilkinson C. F., Christoph G. R., Julien E., Kelley J. M., Kronenberg J., McCarthy J., and Reiss R., 2000: Assessing the risks of exposures to multiple chemicals with a common mechanism of toxicity: How to cumulate? *Regulatory Toxicology and Pharmacology*, 31, 30-43.

6 Annexes

6.1 Exposure calculations

Exposure to pesticide residues from food has been calculated from an estimate of the average content in foods of pesticides included in the monitoring programmes, combined with the estimated average consumption of Danish consumer groups.

As previously explained, most pesticides and all significant food items in the monitoring programmes have been included all eight years (2004-2011). All pesticides included in the monitoring programmes were used in the exposure calculations, but the available data for recently included pesticides do not cover all eight years.

The monitoring programmes primarily include fresh fruit and vegetables and cereals. For some commodities, a smaller part of the samples were sampled as frozen food. Results from these samples have been included in the results for fresh items.

In general, samples were analysed with their peels (conforming to the definitions of maximum residue levels), thus the measured content might include parts that are not normally consumed. In a few important cases, corrections have been made for the reduction of pesticide contents due to peeling. However, no other corrections have been made for the reduction of residues that may happen during preparation of the food.

Calculation of the average content of pesticides

Data have been extracted from the LIMS of the Danish Veterinary and Food Administration and transformed to the EFSA Standard Sample Description (Andersen, 2011), expanding the data on detected residues to include full information on the analytical profile for each sample, including the actual reporting limit (which may vary between the different years) for non-detected pesticides.

An average content has been calculated for each combination of pesticide, food item and origin.

Origin: As previously shown, the residue levels sometimes differ considerably between countries. Therefore calculations were performed separately for the two groups: Domestic or foreign origin.

Non-detected residues: Even though a pesticide has been used, not all pesticides residues are detected in all samples. Due to technical and economical limitations in the monitoring programme, some samples will contain residues not detected by the analytical procedures, either because the pesticides were not included in the programme or because the residue content was lower than the reporting limit used.

Due to the low detection frequency of most pesticides, it is difficult to set up a model for handling of left censored data (EFSA, 2010). Three different models have been used:

Model 1:

Content of pesticides not detected have been assumed 0 (zero).

In this model, the calculated results might be underestimated, since non-detected residues are ignored.

Model 2:

If a pesticide has not been detected in a commodity/origin combination, the average content of the pesticide is assumed to be 0 (zero) in that commodity/origin combination.

If a pesticide has been detected in a commodity/origin combination, content of pesticides not detected in a sample of that commodity/origin combination have been assumed to be 50% of the limit of detection (LOR).

In this model, the average content is over-estimated; because of sometimes very small frequencies of detection, the contribution from the 50% LOR-correction can be very high (in extreme cases up to nearly 3000 times more than the result using Model 0).

Model 3:

The average content is calculated in the same way as in Model 2. But the result from the 50%LOR-correction is limited to 25 times the result that has been calculated using Model 1. In this model extreme corrections from the 50% LOR-contribution are prevented. The background for this model is discussed in Annex6.2.

Processing factors: Detailed information on the actual processing performed by consumers as well as the effect on the residue levels is limited. On the other hand, intake from citrus fruits, bananas and melons contributes significantly to the total exposure of pesticides. As these food items mostly are consumed after peeling, corrections have been made for this process. Data has shown that approximately 90% of residues in these food items are found in the peel, and only 10% remains in the edible part – except for thiabendazole and pesticides from the benomyl group (carbendazim, thiophanat-methyl and benomyl), where about 25% remains in the edible parts (Appendix 7.6).

Calculation of consumption data

Consumption data have been provided by the Department of Nutrition at the Food Institute. The data used in the exposure calculations can be found in Appendix 7.4.

The data was collected as a part of DANSDA (DANish National Survey of Diet and physical Activity) in 2005-2008 and is a subset of the data reported in “Danskernes kostvaner 2003-08” (Pedersen et al. 2010). The subset was chosen as it matches the period for chemical analysis best and as it is the most recently reported dataset. The dataset covers exposure of food and beverages recorded for 7 consecutive days collected from a representative sample of 2700 Danish consumers aged 4 to 75 years. The individuals are drawn as simple random sample from the civil population registration system. DANSDA uses a 7 day pre-coded (semi-closed) food diary with answering categories for the most commonly consumed foods and dishes in the Danish diet. The questionnaire is organized according to the typical daily meal pattern. For food items not found in the pre-coded categories it is possible to note type and amount eaten. The amounts of food eaten were given in household measures and estimated from photos of different portion sizes. The information collected represents information about the cur-

rent dietary exposure in the population. The Danish National Centre for Social Research carried out an interview and the instruction in registration of the dietary exposure.

In this report the exposure is calculated in $\mu\text{g}/\text{kg bw}/\text{day}$ and then recalculated to $\mu\text{g}/\text{day}$. In the previous report the exposure was only calculated in $\mu\text{g}/\text{day}$.

For each participant consumption and body weight was combined giving the consumption in $\text{g}/\text{kg bw}/\text{day}$ (or $\text{mg}/\text{kg bw}/\text{day}$). For food the means was then calculated for the relevant consumer group, e.g. children 4-6 years of age.

To give the exposure in $\mu\text{g}/\text{person}/\text{day}$ the exposure in $\mu\text{g}/\text{kg bw}/\text{day}$ has been multiplied with the mean bodyweight of the relevant consumer group. Because of rounding of the figures multiplying an exposure in $\mu\text{g}/\text{kg bw}/\text{day}$ in Section 4.3 with a mean bodyweight will not necessarily give the value indicated in $\mu\text{g}/\text{day}$. It was shown that calculating in this way only give a reasonable result if the bodyweights do not differs considerable within the consumer group. Because of this the exposure has not be calculated for the consumer group "All" (4-75 years) but for "Adults" instead of.

It should be noted that consumption data are not available on raw commodities but on food as eaten, e.g. oranges without peel or prepared food, e.g. bread-roll. Consumption data has been included in the calculations as if they were given for raw commodities. If consumption data were available for both fresh and frozen food and there was residue data for the raw commodity, the consumptions for the two foods were added. The same has been the case for some commodities where consumptions were given for raw and canned but only residues in the raw commodity were analysed, e.g. raw and canned cherries. This have be done to include as many consumption data as possible.

Not all foods included in the dietary survey were included in the exposure calculations since they were not analysed in the monitoring programme. On the other hand not all food analysed were included in the dietary survey. If residue data were available a low consumption of 0.01 g/day was applied, radish and pomelos (see Appendix 7.4).

For bread consumption data have been recalculated to flour. In the dietary survey many data are available for bread, cakes etc. while in the monitoring programme kernels and flour have been analysed. For calculating the consumption for bread to consumption of flour a factor 0.5 and 0.7 was used for wheat while for rye a factor of 0.6 was.

Calculation of exposure

The contribution to exposure of pesticide residues for each combination of pesticide and food item and origin was calculated from the average content of that pesticide and the estimated average consumption of that food item within the target consumer group.

For each group of consumers, the individual contributions to exposure have been summed for food items, pesticides and origin.

Since the food surveys did not include information of the origin of the foods eaten, two models for the origin of food items have been used:

- a. The origins of food items consumed have the same distribution between domestic and foreign produce as the distribution between the samples analysed.
- b. As the previous model, except that the domestic produce is consumed whenever available (i.e. has been sampled in the monitoring programmes)

Effects of models and analysis of robustness

Calculated results for the exposure to pesticides are subject to some uncertainty partly caused by differences between the real world and the calculating models used, and partly because the data used in the modelling is sampled with some statistical uncertainty.

While the uncertainty of the residue contents in the single samples is well described, as determinations were performed by accredited methods (normally an analytical reproducibility standard deviation of 15-25% would be expected), the bias of the average content is not known due to the unknown contribution of non-detected residues and the origin of the food in the diet.

Average content: Calculations with different models for the compensation of non-detected residues reveals that differences can be quite high for some pesticides, while in other cases the compensation seems very reasonable.

The organically grown foods items have been excluded from calculations of exposure, as the consumption of the two types are expected to be very unevenly distributed between consumers. In some studies, no differentiation between organically grown and conventionally grown food items is made when calculating the exposure (EFSA, 2011). In the previous report on pesticide residues 1998-2003 (Poulsen et al., 2004) it was found that including the organically samples (4% of total samples) would reduce the exposure by approx. 6%. In the present report, organically grown samples (6% of total samples) are excluded from exposure calculations.

Consumption: The distribution of residues between samples of different origin could be expected to vary according to the pattern of pesticides used under different growing conditions and legislation. This expectation seems to be confirmed by the present study.

As previously mentioned, the food consumption study did not provide information on the origin of the consumed food. The sample plans from the monitoring programmes normally target samples to the expected distribution between consumption of domestic and foreign grown items, but the actual distribution might differ from this estimate.

In models that include corrections for non-detected residues (Model 2 and 3), samples have been separated in two groups for the correction: Domestically vs. foreign grown. Since the foreign grown samples may include samples from different countries using different types of pesticides, the algorithm used for corrections overestimate the contribution from non-detected residues in the foreign grown samples compared to the domestically grown. When comparing results for Danish vs. foreign grown samples, results for the relative difference calculated by Model 2 is 3-16% higher than results from Model 1.

Model used in the report

Unless otherwise stated, the following model has been used throughout the report:

Origin: The average content of pesticide residues (including compensation for non-detected residues) has been calculated separately for samples of domestic and foreign origin; origin of consumed food items is assumed to follow the same distribution between domestic and foreign samples as the distribution of samples in the monitoring programmes.

Non-detected residues: Results are calculated using Model 2, i.e. by compensating for non-detected residues using the $\frac{1}{2}$ LOR-model, but limiting the result to 25 times the result from Model 1 (which do not compensate for non-detected residues).

Processing factors: Corrections for the reduction of pesticide residues by peeling of citrus fruits, bananas and melons are included.

Organically grown samples are not included.

6.2 Correction for samples with non-detected residues

Comparing models

For children, using Model 2 (assuming that non-detected residues are 50% of the reporting limit) leads to a Hazard Index of 124%, i.e. above 100%, whereas using Model 1 (assuming that non-detected residues are zero) leads to a HI at 14% (not correcting for peeling in either case).

However, an examination of the details leading to these results shows clear indications of over-correction for non-detected residues using Model 2. Apples and pears contribute with 58% and 18% respectively. For apples, this is mainly caused by oxydemeton-methyl with an HQ of 35% (61% of HI for apples) based on one detected residue in 392 samples (correction factor¹: 2305), high values for the limit of reporting during some of the years (2006-2008: 0.36 mg/kg) and a low ADI of 0.0003 mg/kg bw/d. Without this single detection, HI would have been calculated to 23%. For pears a single detected residue of oxydemeton-methyl in 300 samples (correction factor: 2869) contributes with HQ=13% (68% of HI for pears).

Oxydemeton-methyl was detected once in each of four commodities in apples, pears, exotic fruit (218 samples analysed) and pineapples (55 samples analysed). In each case the HQ from oxydemeton-methyl contributed with 61% - 83% of the HI of that commodity (correction factors: 300 –2900).

For bananas (HI, Model 2 (with reduction for peeling) for children: 0.21%), linuron contributes with 25% of that value (HQ: 0.053%, correction factor: 203, one detected residue in 343 samples) and carbaryl contributes with 9% (HQ: 0.019%, correction factor: 408, one detected residue in 425 samples). Thus, one detection each of linuron and carbaryl contributes with 34% of the HI.

For cucumbers (HI, Model 2 for children: 8.2%), aldrin/dieldrin (sum) contributes with 36% of that value (HQ: 3.0%, correction factor: 54, one detected residue in 218 samples) and dichlorvos contributes with 33% of that value (HQ: 2.7%, correction factor: 92, one detected residue in 182 samples). Thus, one detection each of dichlorvos and aldrin/dieldrin (sum) contributes with 69% of the HI.

These (and similar examples) have been the background for setting up Model 3 using a limiting factor 25 on corrections for non-detected residues.

Using Model 3 (without peeling) HI for children has been calculated to 56% (45% of Model 2) as the contribution from oxydemeton-methyl is reduced from 35.5% to 0.38%.

Using Model 3 for (peeled) bananas, HI is reduced to 0,049% (65% of Model 2), mainly because the contribution from linuron and carbaryl is reduced from 0.072% to 0.008%.

¹ Correcting factor: (Result using Model 2 (with ½LOR-corrections)) / (Result using Model 1 (without correction))

Using Model 3 for cucumbers, HI is reduced to 4.2% (51% of Model 2), mainly because the contribution from linuron and carbaryl is reduced from 5.7% to 2.1%.

Limiting the correction

In the last multi-annual report on pesticide residues 1998-2003 (Poulsen et al., 2004), a limiting factor of 25 between the results calculated using Model 1 and Model 3 was used based on empirical evidence. Based on results from the present period, the same factor has been used in the present report.




Figure 30. Relative difference between average content calculated by Model 2 (non-detected residues assumed to be 50% of LOR) and Model 1 (non-detected residues assumed to be zero). Relative differences are shown using two different scales (0-3000 and 0-200, respectively).

(78): Phosalone in orange marmelade: 19 analysed, 1 detected (5.3%). Factor=78.1

(42): Carbaryl in apricots: 31 analysed, 1 detected (3.2%). Factor=41.9

(37): lambda-Cyhalothrin in currants: 20 analysed, 1 detected (5.0%). Factor=37.4

(32): lambda-Cyhalothrin in head cabbage: 54 analysed, 2 detected (3.7%). Factor=32.2

(26): Triadimefon (sum) in pomelos: 61 analysed, 2 detected (3.3%). Factor=25.5

(2869): Oxydemeton-methyl (sum) in pears: 300 analysed, 1 detected (0.3%). Factor=2869

(2305): Oxydemeton-methyl (sum) in apples: 392 analysed, 1 detected (0.3%). Factor=2305

(1147): Binapacryl in plums: 273 analysed, 1 detected (0.4%). Factor=1147

(788): Iprodione in lemons, lime: 432 analysed, 1 detected (0.2%). Factor=788

The relative difference between results using Model 2 (non-detected assumed to be 50% of LOR) and Model 1 (non-detected assumed to be zero) varies from close to 1 to up to nearly

3000 when looking at individual combinations of pesticide, commodity and origin (domestic or foreign) (**Figure 30**).

However, for combinations of pesticide, commodity and origin where the frequency of detection is higher than 3%, the relative difference is less than 25 for all but a few combinations.

7 Appendices

7.1 Commodity groups not included in exposure calculations

No residues were found in samples of animal origin¹, Therefore these samples have been excluded from exposure calculations.

Samples that have been cultivated by organic methods have not been included in exposure calculations. Although a few residues have been found, such samples would not contribute in any significant degree to the exposure of the general population. Also, estimation of their consumption would be uncertain.

Some commodity groups, not representing major parts of a normal diet, have been excluded when none or only a few residues have been found.

In some cases a commodity has been represented by analytical results from very few samples. In cases where the commodity does not represent major parts of a normal diet, these samples have been excluded from the exposure calculations.

The left side of the table below lists the excluded commodities. The number of samples (of Danish resp. foreign origin) analysed is listed together with the number of samples where residues were found (or not).

The right side of the table shows the pesticides that were found in the commodity in question, the number of samples of the commodity analysed for that pesticide and the number of samples where the pesticide was found. The range of reporting limits used are listed together with the average residue concentration in those samples where the pesticide was found.

The table lists the organically grown commodities followed by conventionally grown samples. Within each of these groups, fruit and vegetables, followed by cereals, products of animal origin and baby food are listed. Commodities are listed alphabetically within each group.

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quantifikation (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|---------------------------|---|-------------------------------|---------------------------|------------------------|--------------|-------------------------------|------------------------------------|--|--|
| Organically grown samples | | | | | | | | | |
| Apples (organic) | DK | 27 | 27 | | | | | | |
| Apples (organic) | F | 40 | 40 | | | | | | |
| Avocados (organic) | F | 5 | 5 | | | | | | |
| Bananas (organic) | F | 32 | 31 | 1 | Chlorpyrifos | 32 | 1 | 0.008-0.01 | 0.019 |

¹ Exposure to residues in samples of animal origin of pesticides from environmental sources have been reported in as part of Directive 96/23 and have not been included in the present report.

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quantification (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|----------------------------------|---|-------------------------------|---------------------------|------------------------|---|-------------------------------|------------------------------------|--|--|
| Basil (organic) | DK | 2 | 2 | | | | | | |
| Beans, kidney (organic) | F | 10 | 9 | 1 | Malathion (sum) | 10 | 1 | 0.01 | 0.04 |
| Beans, soy, dried (organic) | F | 1 | 1 | | | | | | |
| Broccoli (organic) | DK | 3 | 3 | | | | | | |
| Broccoli (organic) | F | 2 | 2 | | | | | | |
| Carrots (organic) | DK | 24 | 24 | | | | | | |
| Carrots (organic) | F | 21 | 21 | | | | | | |
| Cauliflowers (organic) | F | 1 | 1 | | | | | | |
| Chick pea (organic) | F | 7 | 7 | | | | | | |
| Chives (organic) | F | 1 | | 1 | Acetamiprid | 1 | 1 | 0.01 | 0.028 |
| Coffee beans, green (organic) | F | 4 | 4 | | | | | | |
| Courgettes (organic) | F | 1 | 1 | | | | | | |
| Cucumbers (organic) | DK | 10 | 10 | | | | | | |
| Cucumbers (organic) | F | 11 | 11 | | | | | | |
| Currants, black (organic) | F | 1 | 1 | | | | | | |
| Currants, red (organic) | DK | 1 | 1 | | | | | | |
| Ginger (organic) | F | 1 | 1 | | | | | | |
| Hazelnuts (organic) | F | 3 | 3 | | | | | | |
| Head cabbage, red (organic) | DK | 2 | 2 | | | | | | |
| Head cabbage, red (organic) | F | 1 | 1 | | | | | | |
| Head cabbage, spring (organic) | DK | 2 | 2 | | | | | | |
| Head cabbage, spring (organic) | F | 1 | 1 | | | | | | |
| Kiwi (organic) | F | 22 | 21 | 1 | Fenhexamid | 22 | 1 | 0.01-0.012 | 0.1 |
| Leek (organic) | DK | 5 | 5 | | | | | | |
| Lemons (organic) | F | 26 | 25 | 1 | Chlorpyrifos Propargite | 26 26 | 1 1 | 0.008-0.01 0.006-0.1 | 0.025 0.027 |
| Lentils (organic) | F | 10 | 10 | | | | | | |
| Lettuce (organic) | DK | 18 | 18 | | | | | | |
| Lettuce (organic) | F | 16 | 16 | | | | | | |
| Limes (organic) | F | 6 | 5 | 1 | Imazalil | 6 | 1 | 0.01 | 0.036 |
| Lingonberries (organic) | F | 1 | 1 | | | | | | |
| Mandarins, clementines (organic) | F | 9 | 9 | | | | | | |
| Melons (organic) | F | 2 | 2 | | | | | | |
| Onions (organic) | DK | 3 | 3 | | | | | | |
| Onions (organic) | F | 6 | 6 | | | | | | |
| Oranges (organic) | F | 31 | 30 | 1 | Imazalil Endosulfan (sum) | 30 3 | 1 1 | 0.01-0.05 0.005 | 0.13 0.009 |
| Parsley (organic) | DK | 3 | 2 | 1 | | | | | |
| Peaches (organic) | F | 1 | 1 | | | | | | |
| Pears (organic) | DK | 6 | 6 | | | | | | |
| Pears (organic) | F | 35 | 32 | 3 | Carbendazim and benomyl Chloromequat Cyhalothrin, lambda- | 35 34 35 | 1 2 1 | 0.007-0.05 0.006-0.01 0.002-0.05 | 0.25 0.027 0.007 |

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quantifikation (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|----------------------------|---|-------------------------------|---------------------------|------------------------|----------------------------|-------------------------------|------------------------------------|--|--|
| Peppers, sweet (organic) | DK | 7 | 7 | | Phosmet (sum) | 35 | 1 | 0.006-0.3 | 0.04 |
| Peppers, sweet (organic) | F | 26 | 25 | 1 | Pyrethrins | 26 | 1 | 0.003-0.04 | 0.14 |
| Persimmon (organic) | F | 1 | 1 | | | | | | |
| Pineapples (organic) | F | 8 | 8 | | | | | | |
| Plums (organic) | F | 1 | 1 | | | | | | |
| Pomegranate (organic) | F | 1 | 1 | | | | | | |
| Potatoes (organic) | DK | 31 | 31 | | | | | | |
| Potatoes (organic) | F | 12 | 11 | 1 | DDT (sum) | 12 | 1 | 0.008-0.05 | 0.012 |
| Raspberries (organic) | F | 2 | 2 | | | | | | |
| Ruccola (organic) | F | 4 | 4 | | | | | | |
| Sage (organic) | DK | 1 | 1 | | | | | | |
| Scarole (organic) | F | 1 | 1 | | | | | | |
| Strawberries (organic) | DK | 10 | 10 | | | | | | |
| Strawberries (organic) | F | 7 | 7 | | | | | | |
| Sugar pea (organic) | F | 1 | 1 | | | | | | |
| Sunflower seed (organic) | F | 5 | 5 | | | | | | |
| Table grapes (organic) | F | 31 | 30 | 1 | Fenhexamid | 29 | 1 | 0.01-0.012 | 0.03 |
| Tea, fruit (organic) | F | 1 | 1 | | | | | | |
| Tea, herbal (organic) | F | 3 | 2 | 1 | Carbendazim and benomyl | 3 | 1 | 0.01 | 0.012 |
| Teas (organic) | F | 11 | 7 | 4 | Bifenthrin | 3 | 2 | 0.006 | 0.21 |
| | | | | | Carbendazim and benomyl | 11 | 1 | 0.01 | 0.13 |
| | | | | | Clomazone | 11 | 1 | 0.015 | 0.05 |
| Tomatoes (organic) | DK | 21 | 21 | | | | | | |
| Tomatoes (organic) | F | 25 | 23 | 2 | Azoxystrobin | 25 | 1 | 0.004-0.04 | 0.008 |
| | | | | | Chlorothalonil | 25 | 1 | 0.006-0.08 | 0.009 |
| | | | | | Pyrethrins | 25 | 1 | 0.003-0.04 | 0.04 |
| Witloof (organic) | DK | 1 | 1 | | | | | | |
| SUM | DK | 177 | 176 | 1 | | | 1 | | |
| SUM | F | 448 | 428 | 20 | | | 24 | | |
| SUM | Total | 625 | 604 | 21 | | | 25 | | |
| Cereals (organic) | | | | | | | | | |
| Barley (malting) (organic) | DK | 1 | 1 | | | | | | |
| Barley grit (organic) | DK | 4 | 4 | | | | | | |
| Barley grit (organic) | F | 1 | 1 | | | | | | |
| Barley kernels (organic) | DK | 4 | 4 | | | | | | |
| Barley kernels (organic) | F | 5 | 5 | | | | | | |
| Millet (organic) | F | 3 | 3 | | | | | | |
| Oat bran (organic) | DK | 2 | 2 | | | | | | |
| Oat kernels (organic) | DK | 13 | 13 | | | | | | |
| Oat kernels (organic) | F | 7 | 7 | | | | | | |
| Rice flour (organic) | F | 1 | 1 | | | | | | |

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quantification (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|--|---|-------------------------------|---------------------------|------------------------|--------------|-------------------------------|------------------------------------|--|--|
| Rice, brown (organic) | F | 9 | 9 | | | | | | |
| Rice, white (organic) | F | 27 | 27 | | | | | | |
| Rice, wild (organic) | F | 2 | 2 | | | | | | |
| Rollled oat (organic) | DK | 25 | 25 | | | | | | |
| Rollled oat (organic) | F | 23 | 22 | 1 | Chloromequat | 23 | 1 | 0.01 | 0.11 |
| Rye flour (organic) | DK | 28 | 28 | | | | | | |
| Rye flour (organic) | F | 11 | 11 | | | | | | |
| Rye flour, bolted (organic) | DK | 1 | 1 | | | | | | |
| Rye kernels (organic) | DK | 19 | 19 | | | | | | |
| Rye kernels (organic) | F | 8 | 8 | | | | | | |
| Spelt (organic) | DK | 11 | 11 | | | | | | |
| Spelt (organic) | F | 6 | 5 | 1 | Chloromequat | 6 | 1 | 0.01 | 0.019 |
| Spelt, flour (organic) | DK | 5 | 5 | | | | | | |
| Spelt, flour (organic) | F | 16 | 16 | | | | | | |
| Spelt, grain (organic) | DK | 3 | 3 | | | | | | |
| Spelt, grain (organic) | F | 1 | 1 | | | | | | |
| Wheat bran (organic) | F | 2 | 2 | | | | | | |
| Wheat flour (organic) | DK | 26 | 25 | 1 | Chloromequat | 26 | 1 | 0.01 | 0.013 |
| Wheat flour (organic) | F | 28 | 27 | 1 | Chloromequat | 28 | 1 | 0.01 | 0.016 |
| Wheat germ (organic) | DK | 1 | 1 | | | | | | |
| Wheat kernels (organic) | DK | 22 | 22 | | | | | | |
| Wheat kernels (organic) | F | 16 | 16 | | | | | | |
| SUM | DK | 165 | 164 | 1 | | | 1 | | |
| SUM | F | 166 | 163 | 3 | | | 3 | | |
| SUM | Total | 331 | 327 | 4 | | | 4 | | |
| Animal products(organic) | | | | | | | | | |
| Pork meat (organic) | DK | 1 | 1 | | | | | | |
| Cow milk, 1 - 2.9% fat (semi-skimmed milk) (organic) | DK | 1 | 1 | | | | | | |
| Cow milk, raw (organic) | DK | 4 | 4 | | | | | | |
| Honey (organic) | DK | 1 | 1 | | | | | | |
| SUM | DK | 7 | 7 | | | | | | |
| SUM | Total | 7 | 7 | | | | | | |
| Other (organic) | | | | | | | | | |
| Marmelade, orange (organic) | F | 1 | 1 | | | | | | |
| Oranges, juice (organic) | F | 1 | 1 | | | | | | |
| Raisin (organic) | F | 3 | 3 | | | | | | |
| Wine, red (organic) | F | 4 | 4 | | | | | | |
| Pasta, dried (organic) | F | 2 | 2 | | | | | | |
| Rice cake (puffed) (organic) | F | 8 | 8 | | | | | | |
| SUM | F | 19 | 19 | | | | | | |
| SUM | Total | 19 | 19 | | | | | | |
| Baby food (organic) | | | | | | | | | |

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quantifikation (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|--|---|-------------------------------|---------------------------|------------------------|---|-------------------------------|------------------------------------|--|--|
| Babyfood, based on fruits and vegetables, canned (organic) | DK | 3 | 3 | | | | | | |
| Babyfood, based on fruits and vegetables, canned (organic) | F | 59 | 59 | | | | | | |
| Babyfood, cereal based, canned (organic) | F | 1 | 1 | | | | | | |
| Babyfood, cereal based, powder (organic) | DK | 6 | 6 | | | | | | |
| Babyfood, cereal based, powder (organic) | F | 21 | 21 | | | | | | |
| SUM | DK | 9 | 9 | | | | | | |
| SUM | F | 81 | 81 | | | | | | |
| SUM | Total | 90 | 90 | | | | | | |
| Fruit and vegetables (not organic) | | | | | | | | | |
| Bamboo shoots | F | 1 | 1 | | | | | | |
| Banana leaves | F | 1 | 1 | | | | | | |
| Brazil nuts | F | 11 | 11 | | | | | | |
| Cashew nut | F | 7 | 7 | | | | | | |
| Chestnuts | F | 4 | 4 | | | | | | |
| Chinese cabbage | DK | 9 | 9 | | | | | | |
| Chinese cabbage | F | 13 | 13 | | | | | | |
| Chinese radish | DK | 1 | | 1 | Trichloronat | 1 | 1 | 0.005 | 0.014 |
| Chinese radish | F | 3 | 3 | | | | | | |
| Globe artichokes | F | 2 | 1 | 1 | Bitertanol | 2 | 1 | 0.01 | 0.012 |
| Jerusalem artichokes | DK | 10 | 10 | | | | | | |
| Jerusalem artichokes | F | 1 | 1 | | | | | | |
| Lucerne sprouts | DK | 1 | 1 | | | | | | |
| Pak-choi | F | 2 | | 2 | Fenvalerat, esfenvalerat, RR- and SS- Fenvalerat, esfenvalerat, RS- and SR- Iprodione | 2 | 1 | 0.008 | 0.01 |
| | | | | | | 2 | 1 | 0.005- 0.008 | 0.005 |
| | | | | | | 2 | 2 | 0.006 | 1.03 |
| Pecans | F | 4 | 4 | | | | | | |
| Pistachios | F | 11 | 11 | | | | | | |
| Poppy seed | F | 5 | 4 | 1 | Chlormephos | 5 | 1 | 0.006 | 0.07 |
| Quince | F | 2 | | 2 | Carbendazim and benomyl | 2 | 2 | 0.007-0.01 | 0.02 |
| | | | | | Chlorpyrifos | 2 | 1 | 0.008-0.01 | 0.07 |
| | | | | | Thiophanate- methyl | 2 | 1 | 0.01-0.011 | 0.013 |
| Salsify | F | 3 | 2 | 1 | Aldrin and Dieldrin | 3 | 1 | 0.008 | 0.016 |
| Swedes | DK | 1 | 1 | | | | | | |
| Tea, herbal | F | 1 | | 1 | Acetamiprid | 1 | 1 | 0.01 | 0.034 |
| | | | | | Buprofezin | 1 | 1 | 0.01 | 0.014 |
| | | | | | Carbendazim and benomyl | 1 | 1 | 0.01 | 0.023 |

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quarifikation (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|-----------------------------------|---|-------------------------------|---------------------------|------------------------|--|-------------------------------|------------------------------------|---|--|
| Turnips | F | 2 | 1 | 1 | Methomyl and Thiodicarb | 1 | 1 | 0.01 | 0.011 |
| Water spinach | F | 3 | 3 | | Cyhalothrin, lambda- | 2 | 1 | 0.002- 0.005 | 0.006 |
| Witloof | DK | 2 | 1 | 1 | Bromide ion | 1 | 1 | 2.5 | 3.3 |
| Witloof | F | 3 | 1 | 2 | Chlorthal- dimethyl Cyfluthrin (sum) | 3 | 1 | 0.005 | 0.011 |
| | | | | | Permethrin (sum) | 3 | 1 | 0.005 | 0.015 |
| Yams | F | 1 | 1 | | | | | | |
| SUM | DK | 24 | 22 | 2 | | | 2 | | |
| SUM | F | 80 | 69 | 11 | | | 19 | | |
| SUM | Total | 104 | 91 | 13 | | | 21 | | |
| Cereals (not organic) | | | | | | | | | |
| Barley (malting) | DK | 1 | | 1 | Glyphosate | 1 | 1 | 0.15 | 1.7 |
| | | | | | Mepiquat | 1 | 1 | 0.01 | 0.027 |
| Millet | F | 2 | 2 | | | | | | |
| Oat bran | DK | 1 | 1 | | | | | | |
| Oat bran | F | 1 | 1 | | | | | | |
| Rye flour, bolted | F | 1 | 1 | | | | | | |
| SUM | DK | 2 | 1 | 1 | | | 2 | | |
| SUM | F | 4 | 4 | | | | | | |
| SUM | Total | 6 | 5 | 1 | | | 2 | | |
| Animal products (not organic) | | | | | | | | | |
| Beef liver | DK | 15 | 15 | | | | | | |
| Beef meat | DK | 195 | 195 | | | | | | |
| Beef meat | F | 196 | 196 | | | | | | |
| Chicken meat | DK | 95 | 95 | | | | | | |
| Chicken meat | F | 5 | 5 | | | | | | |
| Fat, deer (kidney fat) | DK | 30 | 30 | | | | | | |
| Horse meat | DK | 8 | 8 | | | | | | |
| Lambs meat | DK | 25 | 25 | | | | | | |
| Lambs meat | F | 146 | 146 | | | | | | |
| Pork meat | DK | 1097 | 1097 | | | | | | |
| Pork meat | F | 2 | 2 | | | | | | |
| Sheep meet | DK | 5 | 5 | | | | | | |
| Sheep meet | F | 4 | 4 | | | | | | |
| Veal meat | DK | 119 | 119 | | | | | | |
| Duck meat, steak (air or oil) | F | 5 | 5 | | | | | | |
| Cow milk | DK | 1 | 1 | | | | | | |
| Cow milk | DK | 4 | 4 | | | | | | |
| Cow milk, < 1% fat (skimmed milk) | DK | 1 | 1 | | | | | | |
| Cow milk, raw | DK | 11 | 11 | | | | | | |

| Commodity | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | Without detected residues | With detected residues | Pesticide | Number of samples analysed | Number of samples with residues | General limit of quarifikation (mg/kg) | Avg. content (in samples with residues) (mg/kg) |
|---|---|-------------------------------|---------------------------|------------------------|----------------------------|-------------------------------|------------------------------------|---|--|
| Honey | DK | 125 | 125 | | | | | | |
| Honey, monofloral | DK | 4 | 4 | | | | | | |
| SUM | DK | 1735 | 1735 | | | | | | |
| SUM | F | 358 | 358 | | | | | | |
| SUM | Total | 2093 | 2093 | | | | | | |
| Other (not organic) | | | | | | | | | |
| Beans with pods, canned | F | 1 | 1 | | | | | | |
| Cranberry, dried | F | 3 | 3 | | | | | | |
| Juice, grapefruit | F | 1 | | 1 | Orthophe- nylphenol | 1 | 1 | 0.03 | 0.1 |
| Juice, mixed fruit | F | 13 | 13 | | | | | | |
| Kiwi purée | F | 1 | 1 | | | | | | |
| Lemonade | F | 1 | 1 | | | | | | |
| Mangoes, candied | F | 1 | 1 | | | | | | |
| Marmelade, lemon | F | 1 | 1 | | | | | | |
| Marmelade, mixed citrus | F | 2 | 2 | | | | | | |
| Nuts, mixed, roasted | F | 3 | 3 | | | | | | |
| Oranges, juice, conc. | F | 2 | 1 | 1 | Carbendazim and benomyl | 2 | 1 | 0.007 | 0.017 |
| Corn flakes | F | 1 | 1 | | | | | | |
| Rice cake (puffed) | F | 6 | 6 | | | | | | |
| SUM | F | 36 | 34 | 2 | | | | | |
| SUM | Total | 36 | 34 | 2 | | | | | |
| Baby food (not organic) | | | | | | | | | |
| Babyfood, based on fruits and vegetables, canned | DK | 2 | 2 | | | | | | |
| Babyfood, based on fruits and vegetables, canned | F | 16 | 16 | | | | | | |
| Babyfood, based on fruits and vegetables, powder | F | 2 | 2 | | | | | | |
| Babyfood, cereal based, powder | DK | 26 | 26 | | | | | | |
| Babyfood, cereal based, powder | F | 20 | 20 | | | | | | |
| SUM | DK | 28 | 28 | | | | | | |
| SUM | F | 38 | 38 | | | | | | |
| SUM | Total | 66 | 66 | | | | | | |
| SUM | DK | 2147 | 2142 | 5 | | | 6 | | |
| SUM | F | 1230 | 1194 | 36 | | | 48 | | |
| SUM | Total | 3377 | 3336 | 41 | | | 54 | | |

7.2 Pesticides sought in fruit and vegetables resp. cereals in 2004 – 2011 and their frequency of detection in conventionally grown crops.

| Pesticide | Fruit and vegetables | | | Cereals | | |
|-----------------------------|----------------------------|--|-------------------------|----------------------------|--|-------------------------|
| | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) |
| 2,4-D (sum) | 4936 | 61 (1.2%) | 0.01-0.04 | 927 | 0 (0.0%) | 0.15 |
| 2-Naphthoxyacetic acid | 501 | 0 (0.0%) | 0.096 | 927 | 0 (0.0%) | 0.15 |
| 4-Chlorophenoxy acetic acid | 4936 | 0 (0.0%) | 0.023-0.085 | 927 | 0 (0.0%) | 0.33 |
| Acephate | 12477 | 24 (0.2%) | 0.005-0.01 | 1657 | 0 (0.0%) | 0.009-0.06 |
| Acetamiprid | 4936 | 40 (0.8%) | 0.01 | 927 | 0 (0.0%) | 0.06 |
| Aclonifen | 12447 | 1 (0.0%) | 0.007-0.025 | | | |
| Acrinathrin | 4936 | 0 (0.0%) | 0.095 | 927 | 0 (0.0%) | 0.04 |
| Aldicarb (sum) | 12477 | 12 (0.1%) | 0.008-0.033 | 27 | 0 (0.0%) | 0.008 |
| Aldrin and Dieldrin | 12447 | 2 (0.0%) | 0.008-0.01 | 1659 | 0 (0.0%) | 0.008 |
| Atrazine | 12482 | 5 (0.0%) | 0.009-0.01 | 1659 | 0 (0.0%) | 0.008 |
| Azimsulfuron | 3519 | 0 (0.0%) | 0.01 | | | |
| Azinphos-ethyl | 12447 | 0 (0.0%) | 0.014-0.1 | 1659 | 0 (0.0%) | 0.008 |
| Azinphos-methyl | 12484 | 128 (1.0%) | 0.005-0.114 | 1659 | 1 (0.1%) | 0.008 |
| Azoxystrobin | 12447 | 309 (2.5%) | 0.004-0.04 | 1659 | 2 (0.1%) | 0.006-0.008 |
| Benalaxyl (sum) | 8141 | 3 (0.0%) | 0.004 | | | |
| Benfuracarb | 7541 | 6 (0.1%) | 0.009-0.01 | 320 | 0 (0.0%) | 0.009-0.01 |
| Bentazone (sum) | 4936 | 0 (0.0%) | 0.01 | 728 | 0 (0.0%) | 0.03 |
| Bifenthrin | 12447 | 149 (1.2%) | 0.005-0.006 | 1659 | 0 (0.0%) | 0.008-0.05 |
| Binapacryl | 10347 | 1 (0.0%) | 0.007-0.059 | 1659 | 0 (0.0%) | 0.042-0.25 |
| Biphenyl | 10353 | 1 (0.0%) | 0.009-7 | | | |
| Bitertanol | 12486 | 94 (0.8%) | 0.006-0.01 | 1498 | 0 (0.0%) | 0.008 |
| Bromide ion | 37 | 11 (29.7%) | 2.5 | 15 | 3 (20.0%) | 2.5-5 |
| Bromophos | 12447 | 0 (0.0%) | 0.005-0.05 | 1498 | 0 (0.0%) | 0.008-0.083 |
| Bromophos-ethyl | 12447 | 0 (0.0%) | 0.01 | 1659 | 0 (0.0%) | 0.008 |
| Bromopropylate | 12447 | 44 (0.4%) | 0.008-0.025 | 1659 | 0 (0.0%) | 0.042-0.08 |
| Bromoxynil | 4936 | 0 (0.0%) | 0.01-0.018 | 927 | 0 (0.0%) | 0.04 |
| Bromuconazole (sum) | 2100 | 0 (0.0%) | 0.01 | | | |
| Bupirimate | 12487 | 23 (0.2%) | 0.01-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Buprofezin | 12489 | 18 (0.1%) | 0.008-0.01 | 927 | 1 (0.1%) | 0.03 |
| Captafol | 10347 | 1 (0.0%) | 0.004-0.5 | 1659 | 0 (0.0%) | 0.042-0.42 |
| Captan | 8030 | 6 (0.1%) | 0.004-0.3 | 1659 | 0 (0.0%) | 0.008-0.25 |
| Captan/Folpet (sum) | 2317 | 34 (1.5%) | 0.006-0.3 | | | |
| Carbaryl | 11978 | 78 (0.7%) | 0.006-0.05 | 954 | 0 (0.0%) | 0.006-0.33 |
| Carbendazim and benomyl | 12474 | 700 (5.6%) | 0.006-0.05 | 954 | 1 (0.1%) | 0.007-0.07 |
| Carbofuran (sum) | 12447 | 9 (0.1%) | 0.006-0.03 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Carbophenothion | 12447 | 0 (0.0%) | 0.01-0.3 | 1659 | 0 (0.0%) | 0.008-0.5 |
| Carbosulfan | 12482 | 2 (0.0%) | 0.008-0.05 | 1659 | 2 (0.1%) | 0.008-0.083 |
| Carboxin | 2100 | 0 (0.0%) | 0.01 | | | |
| Chinomethionat | 1338 | 0 (0.0%) | 0.006 | | | |
| Chlordane (sum) | | | | 1659 | 0 (0.0%) | 0.008 |
| Chlorfenson | 12447 | 0 (0.0%) | 0.01-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Chlorfenvinphos | 12447 | 13 (0.1%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008-0.042 |

| Pesticide | Fruit and vegetables | | | Cereals | | |
|----------------------|----------------------------|--|-------------------------|----------------------------|--|-------------------------|
| | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) |
| Chlormephos | 10347 | 1 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Chloromequat | 1836 | 139 (7.6%) | 0.006-0.1 | 1764 | 337 (19.1%) | 0.0027-0.01 |
| Chlorobenzilate | 12447 | 0 (0.0%) | 0.006-0.01 | 1659 | 0 (0.0%) | 0.008-0.4 |
| Chloropropylate | 12447 | 0 (0.0%) | 0.008-0.01 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Chlorothalonil | 12447 | 76 (0.6%) | 0.006-0.08 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Chlorpropham | 546 | 11 (2.0%) | 0.002-0.03 | | | |
| Chlorpropham (sum) | 11901 | 7 (0.1%) | 0.002-0.03 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Chlorpyrifos | 12447 | 1162 (9.3%) | 0.007-0.01 | 1659 | 4 (0.2%) | 0.01-0.042 |
| Chlorpyrifos-methyl | 12447 | 58 (0.5%) | 0.005-0.007 | 1659 | 10 (0.6%) | 0.01-0.042 |
| Chlorthal-dimethyl | 8141 | 3 (0.0%) | 0.005-0.006 | | | |
| Cinidon-ethyl (sum) | 3519 | 0 (0.0%) | 0.01 | | | |
| Clethodim | 4936 | 0 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.11 |
| Clodinafop-Propargyl | 3519 | 0 (0.0%) | 0.02 | | | |
| Clofentezine | 7825 | 5 (0.1%) | 0.023-0.06 | | | |
| Clomazone | 4936 | 1 (0.0%) | 0.01-0.015 | 927 | 0 (0.0%) | 0.05 |
| Cyfluthrin (sum) | 12447 | 9 (0.1%) | 0.005-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Cyhalothrin, lambda- | 12447 | 180 (1.4%) | 0.002-0.05 | | | |
| Cypermethrin (sum) | 12447 | 235 (1.9%) | 0.002-0.009 | 1659 | 0 (0.0%) | 0.008 |
| Cyproconazole | 2100 | 0 (0.0%) | 0.005 | | | |
| Cyprodinil | 12447 | 403 (3.2%) | 0.004-0.01 | | | |
| Cyromazine | 4936 | 17 (0.3%) | 0.026-0.2 | 927 | 0 (0.0%) | 0.14 |
| DDT (sum) | 12447 | 6 (0.0%) | 0.007-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Deltamethrin | 12447 | 80 (0.6%) | 0.005-0.008 | 1659 | 28 (1.7%) | 0.007-0.008 |
| Demeton-S-Methyl | 12487 | 0 (0.0%) | 0.002-0.3 | 927 | 0 (0.0%) | 0.33 |
| Dialifos | 12447 | 0 (0.0%) | 0.005-0.05 | | | |
| Diazinon | 12447 | 39 (0.3%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008 |
| Dichlofenthion | 8141 | 0 (0.0%) | 0.004-0.025 | | | |
| Dichlofluanid | 12447 | 2 (0.0%) | 0.005-0.06 | 1659 | 0 (0.0%) | 0.008-0.05 |
| Dichlorprop | 4936 | 2 (0.0%) | 0.01-0.025 | 927 | 0 (0.0%) | 0.05 |
| Dichlorvos | 10347 | 1 (0.0%) | 0.006 | 1498 | 0 (0.0%) | 0.008 |
| Dicloran | 12447 | 8 (0.1%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Dicofol (sum) | 12447 | 193 (1.6%) | 0.007-0.25 | 1498 | 0 (0.0%) | 0.042-0.083 |
| Diethofencarb | 8181 | 4 (0.0%) | 0.003-0.025 | 927 | 0 (0.0%) | 0.05 |
| Difenoconazole | 12447 | 70 (0.6%) | 0.004-0.03 | | | |
| Diflufenican | 12482 | 3 (0.0%) | 0.004-0.03 | | | |
| Dimethoate (sum) | 12487 | 88 (0.7%) | 0.007-0.01 | 927 | 2 (0.2%) | 0.04 |
| Dimethomorph | 4936 | 47 (1.0%) | 0.01 | 927 | 0 (0.0%) | 0.1 |
| Dimoxystrobin | 3519 | 0 (0.0%) | 0.01 | | | |
| Diniconazole | 2100 | 1 (0.0%) | 0.01 | | | |
| Dinoterb | 4936 | 0 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.07 |
| Dioxathion | 12447 | 0 (0.0%) | 0.005-0.025 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Diphenylamine | 12447 | 112 (0.9%) | 0.005-0.006 | 1659 | 0 (0.0%) | 0.008 |
| Ditalimfos | 12447 | 0 (0.0%) | 0.008-0.03 | 1366 | 0 (0.0%) | 0.008 |
| Dithiocarbamates | 7902 | 460 (5.8%) | 0.1 | | | |
| DNOC | 4936 | 0 (0.0%) | 0.01-0.046 | 927 | 0 (0.0%) | 0.08 |
| Endosulfan (sum) | 12447 | 110 (0.9%) | 0.005-0.008 | 1659 | 0 (0.0%) | 0.008 |
| Endrin | 12447 | 0 (0.0%) | 0.005-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Epoxiconazole | 4936 | 2 (0.0%) | 0.01-0.016 | 927 | 0 (0.0%) | 0.02 |
| Ethiofencarb | 10461 | 0 (0.0%) | 0.007-0.01 | 954 | 0 (0.0%) | 0.007-0.03 |
| Ethion | 12447 | 9 (0.1%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008-0.05 |

| Pesticide | Fruit and vegetables | | | Cereals | | |
|---------------------------------------|----------------------------|--|-------------------------|----------------------------|--|-------------------------|
| | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) |
| Ethoxyquin | 7628 | 1 (0.0%) | 0.05-0.25 | | | |
| Etrimfos | 12447 | 0 (0.0%) | 0.008-0.01 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Famoxadone | 3519 | 10 (0.3%) | 0.01 | | | |
| Fenamiphos (sum) | 1555 | 0 (0.0%) | 0.005 | | | |
| Fenarimol | 12447 | 15 (0.1%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008-0.8 |
| Fenazaquin | 4936 | 6 (0.1%) | 0.01 | 927 | 0 (0.0%) | 0.02 |
| Fenbuconazole | 2100 | 2 (0.1%) | 0.01 | | | |
| Fenchlorphos (sum) | 12447 | 0 (0.0%) | 0.008-0.01 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Fenhexamid | 10461 | 399 (3.8%) | 0.01-0.012 | 27 | 0 (0.0%) | 0.012 |
| Fenitrothion | 12447 | 28 (0.2%) | 0.005-0.04 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Fenoxaprop-P-Ethyl | 12490 | 1 (0.0%) | 0.005-0.04 | | | |
| Fenpropathrin | 12447 | 5 (0.0%) | 0.005-0.06 | 1659 | 0 (0.0%) | 0.083-0.4 |
| Fenpropidin | 12482 | 0 (0.0%) | 0.005-0.06 | | | |
| Fenpropimorph | 12447 | 6 (0.0%) | 0.005-0.03 | | | |
| Fenson | 12447 | 1 (0.0%) | 0.01-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Fenthion (sum) | 12482 | 40 (0.3%) | 0.006-0.05 | 1498 | 0 (0.0%) | 0.008 |
| Fenvalerat, esfenvalerat, RR- and SS- | 12447 | 14 (0.1%) | 0.008-0.05 | 1659 | 0 (0.0%) | 0.04 |
| Fenvalerat, esfenvalerat, RS- and SR- | 12447 | 9 (0.1%) | 0.005-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Fluazifop-P-butyl (sum) | 4936 | 0 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.14 |
| Flucythrinate | 12447 | 2 (0.0%) | 0.004-0.01 | | | |
| Flucythrinate (sum) | | | | 1659 | 0 (0.0%) | 0.008 |
| Fludioxonil | 12447 | 250 (2.0%) | 0.005-0.06 | | | |
| Flufenacet (sum) | 3519 | 0 (0.0%) | 0.01 | | | |
| Fluoxastrobin | 4936 | 2 (0.0%) | 0.01 | | | |
| Flupyrsulfuron-methyl | 4936 | 0 (0.0%) | 0.01-0.04 | | | |
| Fluquinconazole | 2100 | 0 (0.0%) | 0.04 | | | |
| Fluroxypyr (sum) | 4936 | 2 (0.0%) | 0.018-0.026 | | | |
| Flurtamone | 2100 | 0 (0.0%) | 0.005 | | | |
| Flusilazole | 9775 | 5 (0.1%) | 0.005-0.017 | | | |
| Flutolanil | 8141 | 0 (0.0%) | 0.02-0.051 | | | |
| Flutriafol | 2100 | 11 (0.5%) | 0.005 | | | |
| Fluvalinate, tau- | 4306 | 5 (0.1%) | 0.01 | | | |
| Folpet | 8030 | 0 (0.0%) | 0.005-0.044 | 1659 | 0 (0.0%) | 0.008 |
| Fonofos | 8141 | 0 (0.0%) | 0.01-0.048 | | | |
| Formothion | 11109 | 0 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Fuberidazole | 2100 | 0 (0.0%) | 0.005 | | | |
| Furathiocarb | 12447 | 0 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Glyphosate | 49 | 6 (12.2%) | 0.15 | 1375 | 21 (1.5%) | 0.07-0.83 |
| HCH (sum) | 12447 | 2 (0.0%) | 0.008-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Heptachlor (sum) | 12447 | 0 (0.0%) | 0.005-0.025 | 1498 | 0 (0.0%) | 0.008 |
| Heptenophos | 12482 | 0 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Hexachlorobenzene | 12447 | 4 (0.0%) | 0.007-0.01 | 1659 | 0 (0.0%) | 0.008 |
| Hexaconazole | 9751 | 8 (0.1%) | 0.005-0.01 | | | |
| Hexythiazox | 4936 | 17 (0.3%) | 0.01 | 927 | 0 (0.0%) | 0.06 |
| Imazalil | 12146 | 1993 (16.4%) | 0.01-0.05 | 954 | 0 (0.0%) | 0.011-0.04 |
| Iodosulfuron-methyl | 4936 | 0 (0.0%) | 0.01-0.04 | | | |
| Iprodione | 12447 | 442 (3.6%) | 0.006-0.1 | 1659 | 1 (0.1%) | 0.008 |
| Iprovalicarb | 4936 | 10 (0.2%) | 0.01 | | | |
| Isofenphos | 12447 | 0 (0.0%) | 0.01-0.06 | 1659 | 0 (0.0%) | 0.042-0.083 |

| Pesticide | Fruit and vegetables | | | Cereals | | |
|-------------------------|----------------------------|--|-------------------------|----------------------------|--|-------------------------|
| | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) |
| Isofenphos-Methyl | 8141 | 0 (0.0%) | 0.005-0.006 | | | |
| Isoproturon | 4936 | 0 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.008-0.03 |
| Jodfenphos | 12447 | 0 (0.0%) | 0.006-0.025 | 1498 | 0 (0.0%) | 0.008 |
| Kresoxim-methyl | 12447 | 43 (0.3%) | 0.005-0.007 | 1659 | 0 (0.0%) | 0.008 |
| Lindane | 12447 | 1 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Linuron | 10434 | 72 (0.7%) | 0.01-0.04 | 27 | 0 (0.0%) | 0.012 |
| Malathion (sum) | 12487 | 179 (1.4%) | 0.008-0.01 | 1659 | 39 (2.4%) | 0.008-0.08 |
| Mecarbam | 12482 | 0 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Mecoprop (sum) | 4936 | 1 (0.0%) | 0.01-0.022 | 927 | 0 (0.0%) | 0.04 |
| Mepiquat | 1353 | 4 (0.3%) | 0.01 | 1764 | 21 (1.2%) | 0.01 |
| Metalaxyl (sum) | 12487 | 89 (0.7%) | 0.005-0.03 | 1630 | 0 (0.0%) | 0.008-0.05 |
| Methacrifos | 10812 | 1 (0.0%) | 0.005-0.074 | 293 | 0 (0.0%) | 0.1 |
| Methamidophos | 12477 | 23 (0.2%) | 0.006-0.01 | 954 | 0 (0.0%) | 0.008-0.08 |
| Methidathion | 12447 | 96 (0.8%) | 0.01-0.3 | 293 | 0 (0.0%) | 0.5 |
| Methiocarb (sum) | 10461 | 11 (0.1%) | 0.01-0.011 | 755 | 0 (0.0%) | 0.011-0.04 |
| Methomyl and Thiodicarb | 10461 | 86 (0.8%) | 0.01 | 954 | 0 (0.0%) | 0.01-0.04 |
| Methoxychlor | 12447 | 0 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008 |
| Metribuzin | 3519 | 0 (0.0%) | 0.01 | | | |
| Mevinphos | 12482 | 1 (0.0%) | 0.005-0.01 | 1498 | 0 (0.0%) | 0.008 |
| Molinate | 8141 | 0 (0.0%) | 0.02-0.082 | | | |
| Monocrotophos | 12488 | 1 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Monolinuron | 4936 | 0 (0.0%) | 0.01-0.2 | 927 | 0 (0.0%) | 0.08 |
| Myclobutanil | 12447 | 130 (1.0%) | 0.006-0.025 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Nitrofen | 9751 | 0 (0.0%) | 0.004-0.005 | | | |
| Nuarimol | 12487 | 0 (0.0%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008 |
| Ofurace | 9791 | 0 (0.0%) | 0.005-0.01 | 927 | 0 (0.0%) | 0.13 |
| Orthophenylphenol | 12447 | 557 (4.5%) | 0.005-0.03 | | | |
| Oxadixyl | 8179 | 0 (0.0%) | 0.01-0.04 | 954 | 0 (0.0%) | 0.04 |
| Oxamyl | 10461 | 12 (0.1%) | 0.01 | 27 | 0 (0.0%) | 0.01 |
| Oxycarboxin | 9791 | 0 (0.0%) | 0.006-0.01 | 927 | 0 (0.0%) | 0.12 |
| Oxydemeton-methyl (sum) | 12435 | 4 (0.0%) | 0.008-0.36 | 927 | 0 (0.0%) | 0.04-0.05 |
| Paclbutrazol | 2100 | 0 (0.0%) | 0.01 | | | |
| Parathion | 12447 | 0 (0.0%) | 0.005-0.06 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Parathion-methyl (sum) | 12447 | 12 (0.1%) | 0.007-0.06 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Penconazole | 12447 | 34 (0.3%) | 0.007-0.01 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Pendimethalin | 8181 | 3 (0.0%) | 0.005-0.017 | 927 | 0 (0.0%) | 0.06 |
| Pentachloroanisole | 12447 | 0 (0.0%) | 0.008-0.01 | | | |
| Pentachlorobenzene | 10347 | 1 (0.0%) | 0.005-0.007 | | | |
| Pentachlorophenol | 12447 | 1 (0.0%) | 0.006-0.1 | 1659 | 0 (0.0%) | 0.008-0.4 |
| Pentachlorothioanisole | 12447 | 0 (0.0%) | 0.005-0.05 | | | |
| Permethrin (sum) | 12447 | 17 (0.1%) | 0.005-0.006 | 1659 | 2 (0.1%) | 0.008 |
| Phenkapton | 10347 | 0 (0.0%) | 0.009 | 1659 | 0 (0.0%) | 0.008 |
| Phenthoate | 12447 | 2 (0.0%) | 0.009-0.01 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Phorate (sum) | 12484 | 4 (0.0%) | 0.008-0.046 | | | |
| Phosalone | 12447 | 36 (0.3%) | 0.01-0.06 | 1659 | 0 (0.0%) | 0.042-0.08 |
| Phosmet (sum) | 12447 | 29 (0.2%) | 0.006-0.3 | 1659 | 0 (0.0%) | 0.008-0.5 |
| Phosphamidon | 12482 | 0 (0.0%) | 0.006-0.253 | | | |
| Phoxim | 12482 | 0 (0.0%) | 0.006-0.197 | 1498 | 0 (0.0%) | 0.008-0.042 |
| Picolinafen | 4936 | 0 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.04 |
| Picoxystrobin | 3519 | 0 (0.0%) | 0.01 | | | |

| Pesticide | Fruit and vegetables | | | Cereals | | |
|-----------------------|----------------------------|--|-------------------------|----------------------------|--|-------------------------|
| | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) |
| Pirimicarb (sum) | 12487 | 47 (0.4%) | 0.0055-0.01 | 1659 | 0 (0.0%) | 0.009 |
| Pirimiphos-Ethyl | 12447 | 0 (0.0%) | 0.005-0.008 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Pirimiphos-methyl | 12447 | 26 (0.2%) | 0.005-0.01 | 1659 | 86 (5.2%) | 0.008-0.042 |
| Prochloraz (sum) | 12447 | 191 (1.5%) | 0.004-0.06 | | | |
| Procyimdone | 12447 | 152 (1.2%) | 0.006-0.02 | 1659 | 0 (0.0%) | 0.008 |
| Profenofos | 12447 | 24 (0.2%) | 0.005-0.025 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Propamocarb (sum) | 4936 | 110 (2.2%) | 0.01 | 927 | 0 (0.0%) | 0.04 |
| Propanil | 2100 | 0 (0.0%) | 0.01 | | | |
| Propargite | 12447 | 64 (0.5%) | 0.006-0.3 | | | |
| Propham | 12447 | 0 (0.0%) | 0.004-0.3 | 1659 | 0 (0.0%) | 0.083-0.4 |
| Propiconazole | 12482 | 9 (0.1%) | 0.01-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Propoxur | 10461 | 1 (0.0%) | 0.007-0.01 | 954 | 0 (0.0%) | 0.007-0.04 |
| Propyzamide | 12447 | 6 (0.0%) | 0.007-0.03 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Proquinazid | 4936 | 0 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.02 |
| Prothiofos | 12447 | 9 (0.1%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008 |
| Pymetrozine | 4936 | 13 (0.3%) | 0.01 | 927 | 0 (0.0%) | 0.06 |
| Pyraclostrobin | 4936 | 213 (4.3%) | 0.01 | 927 | 0 (0.0%) | 0.03 |
| Pyrazophos | 12487 | 0 (0.0%) | 0.01 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Pyrethrins | 12447 | 2 (0.0%) | 0.003-0.04 | | | |
| Pyridaben | 8181 | 18 (0.2%) | 0.01-0.015 | 927 | 0 (0.0%) | 0.04 |
| Pyridaphenthion | 9791 | 0 (0.0%) | 0.002-0.01 | 927 | 0 (0.0%) | 0.05 |
| Pyridate (sum) | 4435 | 1 (0.0%) | 0.01 | 927 | 0 (0.0%) | 0.13 |
| Pyrimethanil | 11963 | 190 (1.6%) | 0.005-0.04 | | | |
| Pyriproxyfen | 4936 | 93 (1.9%) | 0.01 | 927 | 0 (0.0%) | 0.02 |
| Quinalphos | 12447 | 1 (0.0%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Quinoxifen | 2100 | 1 (0.0%) | 0.005 | | | |
| Quintozene (sum) | 12447 | 21 (0.2%) | 0.006-0.03 | 1659 | 0 (0.0%) | 0.008 |
| Quizalofop | 4936 | 3 (0.1%) | 0.016-0.04 | | | |
| Simazine | 12482 | 0 (0.0%) | 0.01-0.204 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Spiroxamine | 4936 | 11 (0.2%) | 0.01 | 927 | 0 (0.0%) | 0.03 |
| Sulfotep | 12447 | 0 (0.0%) | 0.005-0.025 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Tebuconazole | 12447 | 178 (1.4%) | 0.006-0.025 | 1659 | 1 (0.1%) | 0.019-0.042 |
| Tebufenozide | 3519 | 3 (0.1%) | 0.01 | | | |
| Tebufenpyrad | 8181 | 31 (0.4%) | 0.009-0.011 | 927 | 0 (0.0%) | 0.1 |
| Tecnazene | 12447 | 2 (0.0%) | 0.007-0.06 | 1659 | 0 (0.0%) | 0.008-0.08 |
| TEPP | 12482 | 0 (0.0%) | 0.01-0.06 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Tetrachlorvinphos | 12482 | 1 (0.0%) | 0.008-0.03 | 1659 | 0 (0.0%) | 0.05-0.083 |
| Tetraconazole | 2100 | 2 (0.1%) | 0.01 | | | |
| Tetradifon | 12447 | 13 (0.1%) | 0.01-0.05 | 1659 | 0 (0.0%) | 0.01-0.083 |
| Tetrasul | 12447 | 0 (0.0%) | 0.007-0.01 | 1659 | 0 (0.0%) | 0.008-0.05 |
| Thiabendazole | 12447 | 1036 (8.3%) | 0.01-0.05 | 954 | 0 (0.0%) | 0.015-0.05 |
| Thifensulfuron-methyl | | | | 927 | 0 (0.0%) | 0.14 |
| Thiometon | 12447 | 0 (0.0%) | 0.01-0.3 | | | |
| Thiophanate-methyl | 10461 | 58 (0.6%) | 0.01-0.011 | 27 | 0 (0.0%) | 0.011 |
| Tolclofos-methyl | 12483 | 9 (0.1%) | 0.006-0.019 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Tolyfluanid (sum) | 12447 | 172 (1.4%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.042 |
| Triadimefon (sum) | 12482 | 118 (0.9%) | 0.005-0.044 | 1659 | 0 (0.0%) | 0.008-0.083 |
| Triallate | 4936 | 2 (0.0%) | 0.01-0.1 | | | |
| Triasulfuron | | | | 927 | 0 (0.0%) | 0.04-0.15 |
| Triazophos | 12482 | 6 (0.0%) | 0.006-0.05 | 1659 | 0 (0.0%) | 0.008-0.083 |

| Pesticide | Fruit and vegetables | | | Cereals | | |
|-------------------|----------------------------|--|-------------------------|----------------------------|--|-------------------------|
| | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) | Number of samples analysed | Number of samples with detected residues | Reporting limit (mg/kg) |
| Trichlorfon | 12482 | 3 (0.0%) | 0.01-0.084 | | | |
| Trichloronat | 12447 | 1 (0.0%) | 0.005-0.03 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Trifloxystrobin | 12447 | 87 (0.7%) | 0.002-0.025 | 1659 | 0 (0.0%) | 0.008-0.042 |
| Triflumuron | 4936 | 21 (0.4%) | 0.01 | | | |
| Trifluralin | 2100 | 0 (0.0%) | 0.01 | | | |
| Triticonazole | 2100 | 0 (0.0%) | 0.01 | | | |
| Vamidothion | 12482 | 0 (0.0%) | 0.003-0.05 | | | |
| Vinclozolin (sum) | 12447 | 81 (0.7%) | 0.005-0.01 | 1659 | 0 (0.0%) | 0.008-0.042 |

7.3 Pesticides included in the monitoring and commodities where residues were found.

The left side of the table lists all pesticides found during the monitoring programmes 2004 – 2011 with representative sampling. The number of samples (of Danish resp. foreign origin) analysed for each pesticide is listed together with the number of samples where residues of that pesticide were found (or not). The full list of pesticides searched for can be found in Annex 6.2.

The right side of the table shows the commodities where the pesticide in question was found, the number of samples of the commodity that was analysed for the pesticide and the number of samples where the pesticide was found.

Commodities are listed alphabetically. The list includes all (conventional or organic) commodity groups.

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues | | | | | |
|-----------------|---|----------------------------|--|--|------------------------|----------------------------|--|-------|----|----------|-----|---|
| 2,4-D (sum) | F | 4516 | 4455 | 61 | Grapefruit | 155 | 19 | | | | | |
| | | | | | Lemons | 158 | 10 | | | | | |
| | | | | | Limes | 18 | 4 | | | | | |
| | | | | | Mandarins, clementines | 160 | 8 | | | | | |
| | | | | | Okra (Lady's fingers) | 5 | 1 | | | | | |
| | | | | | Oranges | 173 | 16 | | | | | |
| | | | | | Peppers, sweet | 170 | 1 | | | | | |
| | | | | | Pomelos | 34 | 2 | | | | | |
| | | | | | Acephate | F | 11068 | 11044 | 24 | Apples | 393 | 1 |
| | | | | | | | | | | Apricots | 31 | 1 |
| Beans with pods | 305 | 4 | | | | | | | | | | |
| Cape gooseberry | 2 | 1 | | | | | | | | | | |
| Chilli peppers | 42 | 2 | | | | | | | | | | |
| Grapefruit | 388 | 1 | | | | | | | | | | |
| Kiwi | 363 | 1 | | | | | | | | | | |
| Leek | 68 | 1 | | | | | | | | | | |
| Lentils | 13 | 1 | | | | | | | | | | |
| Lettuce | 208 | 3 | | | | | | | | | | |
| Limes | 52 | 3 | | | | | | | | | | |
| Melons | 311 | 1 | | | | | | | | | | |
| Plums | 318 | 1 | | | | | | | | | | |
| Strawberries | 265 | 1 | | | | | | | | | | |
| Tomatoes | 250 | 1 | | | | | | | | | | |
| Watermelon | 39 | 1 | | | | | | | | | | |
| Acetamiprid | DK | 1597 | 1594 | 3 | Apples | 69 | 2 | | | | | |
| | | | | | Lettuce | 60 | 1 | | | | | |
| Acetamiprid | F | 4516 | 4478 | 38 | Apples | 124 | 5 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|---------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| | | | | | Aubergines | 32 | 2 |
| | | | | | Chilli peppers | 9 | 1 |
| | | | | | Cucumbers | 89 | 2 |
| | | | | | Grapefruit | 155 | 5 |
| | | | | | Lemons | 158 | 1 |
| | | | | | Lettuce | 96 | 4 |
| | | | | | Nectarines | 90 | 1 |
| | | | | | Oranges | 173 | 1 |
| | | | | | Peaches | 64 | 2 |
| | | | | | Pears | 123 | 1 |
| | | | | | Peas with pods | 26 | 1 |
| | | | | | Peppers, sweet | 170 | 4 |
| | | | | | Tea, herbal | 1 | 1 |
| | | | | | Teas | 16 | 3 |
| | | | | | Tomatoes | 91 | 2 |
| | | | | | Gojiberries, dried | 1 | 1 |
| | | | | | Chives (organic) | 1 | 1 |
| Aclonifen | DK | 3179 | 3178 | 1 | Parsley Root | 33 | 1 |
| Aldicarb (sum) | F | 9973 | 9961 | 12 | Basil | 11 | 1 |
| | | | | | Carrots | 158 | 1 |
| | | | | | Ginger | 28 | 4 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Melons | 311 | 1 |
| | | | | | Nectarines | 211 | 1 |
| | | | | | Ruccola | 12 | 1 |
| | | | | | Table grapes | 458 | 1 |
| Aldrin and Dieldrin | F | 11309 | 11307 | 2 | Cucumbers | 218 | 1 |
| | | | | | Salsify | 3 | 1 |
| Atrazine | F | 11072 | 11067 | 5 | Lemons | 381 | 2 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Oranges | 491 | 1 |
| | | | | | Pomelos | 61 | 1 |
| Azinphos-methyl | DK | 3840 | 3837 | 3 | Strawberries | 163 | 2 |
| | | | | | Rye kernels | 100 | 1 |
| Azinphos-methyl | F | 11074 | 10947 | 127 | Apples | 394 | 45 |
| | | | | | Apricots | 31 | 3 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Nectarines | 212 | 22 |
| | | | | | Peaches | 158 | 12 |
| | | | | | Pears | 301 | 37 |
| | | | | | Peas with pods | 40 | 1 |
| | | | | | Plums | 319 | 3 |
| | | | | | Spinach | 163 | 1 |
| | | | | | Blueberry, dried | 1 | 1 |
| Azoxystrobin | DK | 3881 | 3803 | 78 | Carrots | 340 | 8 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues | | | | | |
|-------------|---|---------------------------------|---|---|-------------------------|---------------------------------|---|------|---|---------|-----|---|
| Azoxytrobin | F | 10951 | 10718 | 233 | Cucumbers | 199 | 36 | | | | | |
| | | | | | Kale | 9 | 3 | | | | | |
| | | | | | Lettuce | 163 | 5 | | | | | |
| | | | | | Parsley Root | 33 | 1 | | | | | |
| | | | | | Parsnips | 51 | 1 | | | | | |
| | | | | | Peas without pods | 10 | 1 | | | | | |
| | | | | | Radish | 3 | 1 | | | | | |
| | | | | | Ruccola | 8 | 1 | | | | | |
| | | | | | Spinach | 81 | 4 | | | | | |
| | | | | | Strawberries | 163 | 14 | | | | | |
| | | | | | Tomatoes | 212 | 2 | | | | | |
| | | | | | Rolled oat | 43 | 1 | | | | | |
| | | | | | Aubergines | 87 | 1 | | | | | |
| | | | | | Bananas | 425 | 28 | | | | | |
| | | | | | Beans with pods | 305 | 23 | | | | | |
| | | | | | Carambola | 103 | 6 | | | | | |
| | | | | | Carrots | 158 | 9 | | | | | |
| | | | | | Celery | 21 | 1 | | | | | |
| | | | | | Chilli peppers | 42 | 2 | | | | | |
| | | | | | Cucumbers | 218 | 17 | | | | | |
| | | | | | Grapefruit | 388 | 3 | | | | | |
| | | | | | Guava | 9 | 1 | | | | | |
| | | | | | Jambolan | 2 | 2 | | | | | |
| | | | | | Lemon grass | 16 | 1 | | | | | |
| | | | | | Lettuce | 208 | 7 | | | | | |
| | | | | | Limes | 52 | 1 | | | | | |
| | | | | | Mandarins, clementines | 411 | 1 | | | | | |
| | | | | | Melons | 311 | 9 | | | | | |
| | | | | | Oranges | 491 | 5 | | | | | |
| | | | | | Papayas | 89 | 1 | | | | | |
| | | | | | Parsnips | 10 | 1 | | | | | |
| | | | | | Peas with pods | 40 | 2 | | | | | |
| | | | | | Peas without pods | 43 | 5 | | | | | |
| | | | | | Peppers, sweet | 385 | 22 | | | | | |
| | | | | | Persimmon | 108 | 1 | | | | | |
| | | | | | Raspberries | 101 | 7 | | | | | |
| | | | | | Ruccola | 12 | 1 | | | | | |
| | | | | | Spinach | 163 | 1 | | | | | |
| | | | | | Spring onions | 32 | 1 | | | | | |
| | | | | | Strawberries | 266 | 19 | | | | | |
| | | | | | Table grapes | 459 | 46 | | | | | |
| | | | | | Tomatoes | 249 | 6 | | | | | |
| | | | | | Vegetables, unspecified | 3 | 1 | | | | | |
| | | | | | Rolled oat | 86 | 1 | | | | | |
| | | | | | Tomatoes (organic) | 25 | 1 | | | | | |
| | | | | | Benalaxyl (sum) | F | 6478 | 6475 | 3 | Lettuce | 153 | 1 |
| | | | | | | | | | | Melons | 232 | 2 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues |
|----------------|---|----------------------------|--|--|------------------------|----------------------------|--|
| Benfuracarb | F | 6237 | 6231 | 6 | Cucumbers | 129 | 1 |
| | | | | | Mandarins, clementines | 251 | 1 |
| | | | | | Nectarines | 121 | 1 |
| | | | | | Pears | 178 | 1 |
| | | | | | Peas without pods | 39 | 1 |
| | | | | | Teas | 24 | 1 |
| Bifenthrin | F | 10951 | 10802 | 149 | Apples | 394 | 9 |
| | | | | | Apricots | 31 | 1 |
| | | | | | Aubergines | 87 | 1 |
| | | | | | Bananas | 425 | 41 |
| | | | | | Beans with pods | 305 | 5 |
| | | | | | Blackberries | 35 | 3 |
| | | | | | Cherries | 24 | 1 |
| | | | | | Chilli peppers | 42 | 2 |
| | | | | | Cucumbers | 218 | 2 |
| | | | | | Garlics | 82 | 1 |
| | | | | | Lettuce | 208 | 5 |
| | | | | | Limes | 52 | 2 |
| | | | | | Melons | 311 | 10 |
| | | | | | Nectarines | 212 | 1 |
| | | | | | Papayas | 89 | 8 |
| | | | | | Peaches | 158 | 3 |
| | | | | | Pears | 301 | 2 |
| | | | | | Peppers, sweet | 385 | 6 |
| | | | | | Persimmon | 108 | 5 |
| | | | | | Plums | 319 | 5 |
| | | | | | Radish | 8 | 1 |
| | | | | | Raspberries | 101 | 1 |
| | | | | | Ruccola | 12 | 3 |
| | | | | | Spinach | 163 | 1 |
| | | | | | Strawberries | 266 | 6 |
| | | | | | Table grapes | 459 | 10 |
| Tomatoes | 249 | 12 | | | | | |
| Teas (organic) | 3 | 2 | | | | | |
| Binapacryl | F | 9620 | 9619 | 1 | Plums | 273 | 1 |
| Biphenyl | F | 8145 | 8144 | 1 | Apples | 349 | 1 |
| Bitertanol | DK | 3747 | 3666 | 81 | Apples | 214 | 37 |
| | | | | | Cherries | 3 | 1 |
| | | | | | Pears | 165 | 40 |
| Bitertanol | F | 10938 | 10925 | 13 | Plums | 72 | 3 |
| | | | | | Cucumbers | 218 | 1 |
| | | | | | Globe artichokes | 2 | 1 |
| | | | | | Nectarines | 212 | 5 |
| | | | | | Onions | 68 | 1 |
| | | | | | Peaches | 158 | 2 |
| Bromide ion | DK | 23 | 21 | 2 | Tomatoes | 250 | 3 |
| | | | | | Ruccola | 1 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|---------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Bromide ion | F | 29 | 17 | 12 | Witloof | 1 | 1 |
| | | | | | Spinach | 14 | 9 |
| | | | | | Rice, white | 7 | 3 |
| Bromopropylate | F | 10951 | 10907 | 44 | Apples | 394 | 6 |
| | | | | | Fruit, mixed | 5 | 1 |
| | | | | | Grapefruit | 388 | 17 |
| | | | | | Lemons | 380 | 8 |
| | | | | | Mandarins, clementines | 411 | 2 |
| | | | | | Oranges | 491 | 3 |
| | | | | | Pears | 301 | 3 |
| | | | | | Pomelos | 61 | 1 |
| | | | | | Table grapes | 459 | 1 |
| | | | | | Tomatoes | 249 | 2 |
| Bupirimate | F | 11077 | 11054 | 23 | Apples | 393 | 1 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Cucumbers | 218 | 3 |
| | | | | | Currants, black | 17 | 1 |
| | | | | | Gooseberries | 7 | 3 |
| | | | | | Melons | 311 | 3 |
| | | | | | Nectarines | 212 | 3 |
| | | | | | Peaches | 158 | 1 |
| | | | | | Peppers, sweet | 385 | 2 |
| | | | | | Strawberries | 266 | 4 |
| | | | | | Tomatoes | 250 | 1 |
| Buprofezin | F | 10496 | 10477 | 19 | Beans with pods | 305 | 2 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Grapefruit | 388 | 4 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Lemons | 381 | 1 |
| | | | | | Mandarins, clementines | 411 | 2 |
| | | | | | Oranges | 491 | 2 |
| | | | | | Pomelos | 61 | 1 |
| | | | | | Tea, herbal | 1 | 1 |
| | | | | | Teas | 40 | 1 |
| | | | | | Tomatoes | 250 | 2 |
| | | | | | Rice, short grained | 94 | 1 |
| Captafol | DK | 3443 | 3442 | 1 | Spinach | 63 | 1 |
| Captan | F | 7679 | 7673 | 6 | Apricots | 29 | 1 |
| | | | | | Lemons | 314 | 1 |
| | | | | | Passion fruits | 36 | 1 |
| | | | | | Table grapes | 401 | 3 |
| Captan/Folpet (sum) | DK | 745 | 727 | 18 | Apples | 192 | 16 |
| | | | | | Pears | 135 | 2 |
| Captan/Folpet (sum) | F | 1583 | 1567 | 16 | Apples | 349 | 9 |
| | | | | | Blackberries | 33 | 1 |
| | | | | | Currants, red | 27 | 3 |
| | | | | | Gooseberries | 7 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------------|---|---------------------------------|---|---|------------------------------|---------------------------------|---|
| Carbaryl | DK | 3497 | 3496 | 1 | Pears | 256 | 1 |
| | F | 10015 | 9930 | 85 | Raspberries | 82 | 1 |
| Carbaryl | | | | | Mushroom (Agaricus bisporus) | 41 | 1 |
| | | | | | Apples | 393 | 20 |
| | | | | | Apricots | 31 | 1 |
| | | | | | Aubergines | 87 | 1 |
| | | | | | Bananas | 425 | 1 |
| | | | | | Celery | 21 | 1 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Grapefruit | 283 | 6 |
| | | | | | Kiwi | 363 | 2 |
| | | | | | Mandarins, clementines | 303 | 1 |
| | | | | | Onions | 68 | 3 |
| | | | | | Oranges | 377 | 2 |
| | | | | | Peaches | 158 | 3 |
| | | | | | Pears | 301 | 10 |
| | | | | | Pineapples | 55 | 8 |
| | | | | | Plums | 319 | 1 |
| | | | | | Rambutan | 47 | 1 |
| | | | | | Raspberries | 101 | 1 |
| | | | | | Sesame seed | 19 | 2 |
| | | | | | Shallots | 12 | 1 |
| | | | | | Strawberries | 265 | 1 |
| | | | | | Table grapes | 458 | 10 |
| | | | | | Blueberry, dried | 1 | 1 |
| | | | | | Wine, red | 265 | 7 |
| Carbendazim and benomyl | DK | 3496 | 3492 | 4 | Broccoli | 21 | 1 |
| | | | | | Onions | 292 | 1 |
| | | | | | Plums | 72 | 1 |
| | | | | | Spinach | 81 | 1 |
| Carbendazim and benomyl | F | 10513 | 9727 | 786 | Apples | 393 | 81 |
| | | | | | Apricots | 33 | 9 |
| | | | | | Aubergines | 87 | 1 |
| | | | | | Basil | 11 | 1 |
| | | | | | Beans with pods | 305 | 19 |
| | | | | | Cape gooseberry | 2 | 1 |
| | | | | | Carambola | 103 | 25 |
| | | | | | Carrots | 159 | 1 |
| | | | | | Celery | 21 | 1 |
| | | | | | Cherries | 24 | 5 |
| | | | | | Chilli peppers | 42 | 8 |
| | | | | | Coriander, leaves | 9 | 2 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Cucumbers | 218 | 11 |
| | | | | | Currants, black | 17 | 7 |
| | | | | | Currants, red | 28 | 4 |
| | | | | | Fruit, mixed | 5 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------------|---|---------------------------------|---|---|------------------------------|---------------------------------|---|
| Carbendazin and benomyl | F | | | | Gooseberries | 7 | 3 |
| | | | | | Grapefruit | 388 | 25 |
| | | | | | Guava | 9 | 1 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Lemons | 381 | 57 |
| | | | | | Lentils | 13 | 1 |
| | | | | | Lettuce | 208 | 4 |
| | | | | | Limes | 52 | 11 |
| | | | | | Litchi | 8 | 1 |
| | | | | | Mandarins, clementines | 411 | 35 |
| | | | | | Mango | 58 | 5 |
| | | | | | Melons | 312 | 22 |
| | | | | | Mixed berries | 2 | 2 |
| | | | | | Mushroom (Agaricus bisporus) | 35 | 9 |
| | | | | | Nectarines | 210 | 23 |
| | | | | | Okra (Lady's fingers) | 15 | 1 |
| | | | | | Onions | 68 | 2 |
| | | | | | Oranges | 491 | 31 |
| | | | | | Papayas | 89 | 21 |
| | | | | | Passion fruits | 37 | 8 |
| | | | | | Peaches | 158 | 29 |
| | | | | | Pears | 301 | 85 |
| | | | | | Peas with pods | 40 | 6 |
| | | | | | Peas without pods | 43 | 1 |
| | | | | | Peppers, sweet | 385 | 7 |
| | | | | | Pineapples | 56 | 1 |
| | | | | | Pitaya | 19 | 6 |
| | | | | | Plums | 318 | 9 |
| | | | | | Pomegranate | 16 | 4 |
| | | | | | Pomelos | 61 | 3 |
| | | | | | Quince | 2 | 2 |
| | | | | | Rambutan | 47 | 18 |
| | | | | | Raspberries | 101 | 9 |
| | | | | | Shallots | 12 | 2 |
| | | | | | Spinach | 162 | 7 |
| | | | | | Spring onions | 32 | 1 |
| | | | | | Strawberries | 265 | 22 |
| | | | | | Table grapes | 458 | 22 |
| | | | | | Tamarillo | 20 | 5 |
| | | | | | Tea, herbal | 1 | 1 |
| Teas | 40 | 1 | | | | | |
| Tomatoes | 250 | 8 | | | | | |
| Vegetables, unspecified | 3 | 1 | | | | | |
| Watermelon | 39 | 2 | | | | | |
| Wheat kernels | 83 | 1 | | | | | |
| Apple, dried | 1 | 1 | | | | | |
| Gojiberries, dried | 1 | 1 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|------------------|---|---------------------------------|---|---|------------------------------|---------------------------------|---|
| Carbofuran (sum) | F | 10951 | 10942 | 9 | Oranges, juice | 44 | 3 |
| | | | | | Oranges, juice, conc. | 2 | 1 |
| | | | | | Raisin | 3 | 1 |
| | | | | | Wine, red | 265 | 80 |
| | | | | | Wine, white | 7 | 2 |
| | | | | | Pears (organic) | 35 | 1 |
| | | | | | Tea, herbal (organic) | 3 | 1 |
| | | | | | Teas (organic) | 11 | 1 |
| | | | | | Beans with pods | 305 | 1 |
| | | | | | Chilli peppers | 42 | 3 |
| Carbosulfan | DK | 3810 | 3809 | 1 | Grapefruit | 388 | 2 |
| | | | | | Melons | 311 | 1 |
| Carbosulfan | F | 11072 | 11068 | 4 | Oranges | 491 | 2 |
| Chlorfenvinphos | DK | 3881 | 3870 | 11 | Rye flour | 28 | 1 |
| | | | | | Pasta | 43 | 1 |
| Chlorfenvinphos | F | 10951 | 10949 | 2 | Chilli peppers | 42 | 1 |
| Chlormephos | F | 9262 | 9261 | 1 | Kiwi | 363 | 1 |
| Chlormequat | DK | 1313 | 1149 | 164 | Rye flour | 57 | 1 |
| | | | | | Pears | 165 | 50 |
| Chlormequat | F | 2479 | 2160 | 319 | Oat kernels | 10 | 4 |
| | | | | | Rolled oat | 71 | 2 |
| | | | | | Rye flour | 58 | 4 |
| | | | | | Rye kernels | 136 | 10 |
| | | | | | Spelt | 37 | 27 |
| | | | | | Spelt, flour | 19 | 8 |
| | | | | | Wheat bran | 2 | 2 |
| | | | | | Wheat flour | 137 | 19 |
| | | | | | Wheat kernels | 231 | 37 |
| | | | | | Wheat flour (organic) | 26 | 1 |
| | | | | | Aubergines | 35 | 1 |
| | | | | | Cucumbers | 113 | 1 |
| | | | | | Mushroom (Agaricus bisporus) | 19 | 7 |
| | | | | | Pears | 300 | 73 |
| | | | | | Peppers, sweet | 230 | 1 |
| | | | | | Table grapes | 89 | 4 |
| | | | | | Oat kernels | 20 | 1 |
| | | | | | Rolled oat | 113 | 42 |
| | | | | | Rye flour | 73 | 22 |
| | | | | | Rye kernels | 73 | 4 |
| | | | | | Spelt | 13 | 13 |
| | | | | | Spelt, flour | 22 | 18 |
| | | | | | Spelt, grain | 2 | 1 |
| Wheat bran | 13 | 5 | | | | | |
| Wheat flour | 223 | 72 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|--------------------|---|---------------------------------|---|---|----------------------------|---------------------------------|---|
| Chlorothalonil | DK | 3881 | 3880 | 1 | Wheat kernels | 185 | 42 |
| | F | 10951 | 10876 | 75 | Pear, canned | 9 | 4 |
| Chlorothalonil | | | | | Pasta | 42 | 1 |
| | | | | | Pasta, dried | 32 | 2 |
| | | | | | Pears (organic) | 34 | 2 |
| | | | | | Rolled oat (organic) | 23 | 1 |
| | | | | | Spelt (organic) | 6 | 1 |
| | | | | | Wheat flour (organic) | 28 | 1 |
| | | | | | Gooseberries | 4 | 1 |
| | | | | | Aubergines | 87 | 4 |
| | | | | | Carambola | 103 | 1 |
| | | | | | Celeriac | 7 | 1 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Cucumbers | 218 | 16 |
| | | | | | Lettuce | 208 | 1 |
| | | | | | Melons | 311 | 10 |
| | | | | | Nectarines | 212 | 5 |
| | | | | | Passion fruits | 38 | 4 |
| | | | | | Peaches | 158 | 6 |
| | | | | | Peas with pods | 40 | 2 |
| | | | | | Strawberries | 266 | 9 |
| | | | | | Table grapes | 459 | 1 |
| | | | | | Tomatoes | 249 | 11 |
| | | | | | Watermelon | 40 | 1 |
| | | | | | Tomatoes (organic) | 25 | 1 |
| Chlorpropham | F | 92 | 81 | 11 | Potatoes, new | 79 | 11 |
| Chlorpropham (sum) | F | 10859 | 10852 | 7 | Apples | 394 | 1 |
| | | | | | Kiwi | 363 | 2 |
| | | | | | Mandarins, clementines | 411 | 2 |
| | | | | | Spinach | 163 | 1 |
| | | | | | Tomatoes | 249 | 1 |
| Chlorpyrifos | DK | 5471 | 5468 | 3 | Radish | 3 | 3 |
| Chlorpyrifos | F | 11309 | 10138 | 1171 | Apples | 394 | 100 |
| | | | | | Apricots | 31 | 1 |
| | | | | | Asian cabbage,(unspecific) | 2 | 1 |
| | | | | | Asparagus, green | 38 | 2 |
| | | | | | Bananas | 425 | 39 |
| | | | | | Beans with pods | 305 | 3 |
| | | | | | Broccoli | 63 | 1 |
| | | | | | Carambola | 103 | 24 |
| | | | | | Carrots | 158 | 5 |
| | | | | | Celery | 21 | 2 |
| | | | | | Chilli peppers | 42 | 8 |
| | | | | | Chives | 6 | 1 |
| | | | | | Coriander, leaves | 9 | 1 |
| | | | | | Currants, black | 17 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|---------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| | | | | | Currants, red | 28 | 1 |
| | | | | | Ginger | 28 | 1 |
| | | | | | Grapefruit | 388 | 145 |
| | | | | | Herbs, unspecified | 3 | 1 |
| | | | | | Kiwano | 1 | 1 |
| | | | | | Kiwi | 363 | 15 |
| | | | | | Lemons | 380 | 136 |
| | | | | | Lettuce | 208 | 1 |
| | | | | | Mandarins, clementines | 411 | 260 |
| | | | | | Mango | 58 | 1 |
| | | | | | Mangosteens | 12 | 2 |
| | | | | | Melons | 311 | 3 |
| | | | | | Mineola | 3 | 2 |
| | | | | | Nectarines | 212 | 24 |
| | | | | | Onions | 68 | 1 |
| | | | | | Oranges | 491 | 213 |
| | | | | | Passion fruits | 38 | 4 |
| | | | | | Peaches | 158 | 30 |
| | | | | | Pears | 301 | 20 |
| | | | | | Peppers, sweet | 385 | 3 |
| | | | | | Persimmon | 108 | 8 |
| | | | | | Pineapples | 55 | 1 |
| | | | | | Plums | 319 | 12 |
| | | | | | Pomegranate | 16 | 1 |
| | | | | | Pomelos | 61 | 14 |
| | | | | | Quince | 2 | 1 |
| | | | | | Rambutan | 47 | 1 |
| | | | | | Raspberries | 101 | 1 |
| | | | | | Ruccola | 12 | 1 |
| | | | | | Sesame seed | 18 | 1 |
| | | | | | Spinach | 163 | 3 |
| | | | | | Spring onions | 32 | 1 |
| | | | | | Strawberries | 266 | 5 |
| | | | | | Table grapes | 459 | 53 |
| | | | | | Tomatoes | 249 | 1 |
| | | | | | Rice, white | 167 | 3 |
| | | | | | Wheat flour | 184 | 1 |
| | | | | | Marmelade, orange | 19 | 8 |
| | | | | | Bananas (organic) | 32 | 1 |
| | | | | | Lemons (organic) | 26 | 1 |
| Chlorpyrifos-methyl | DK | 5471 | 5467 | 4 | Spelt | 30 | 1 |
| | | | | | Spelt, flour | 16 | 1 |
| | | | | | Wheat flour | 78 | 2 |
| Chlorpyrifos-methyl | F | 11309 | 11245 | 64 | Apples | 394 | 7 |
| | | | | | Carrots | 158 | 2 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Kiwi | 363 | 3 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues | | | | | |
|------------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|-------|-----|-----------------|-----|----|
| | | | | | Lemons | 380 | 3 | | | | | |
| | | | | | Mandarins, clementines | 411 | 8 | | | | | |
| | | | | | Nectarines | 212 | 6 | | | | | |
| | | | | | Oranges | 491 | 7 | | | | | |
| | | | | | Peaches | 158 | 5 | | | | | |
| | | | | | Peppers, sweet | 385 | 4 | | | | | |
| | | | | | Strawberries | 266 | 1 | | | | | |
| | | | | | Table grapes | 459 | 11 | | | | | |
| | | | | | Rice, white | 167 | 1 | | | | | |
| | | | | | Wheat flour | 184 | 4 | | | | | |
| | | | | | Wheat kernels | 143 | 1 | | | | | |
| | | | | | Chlorthal-dimethyl | F | 6478 | 6475 | 3 | Lettuce | 153 | 1 |
| | | | | | | | | | | Pears | 202 | 1 |
| | | | | | | | | | | Witloof | 3 | 1 |
| | | | | | Clofentezine | F | 6235 | 6229 | 6 | Strawberries | 159 | 5 |
| Gojiberries, dried | 1 | 1 | | | | | | | | | | |
| Clomazone | F | 4516 | 4515 | 1 | Teas (organic) | 11 | 1 | | | | | |
| Cyfluthrin (sum) | F | 11309 | 11300 | 9 | Apples | 394 | 1 | | | | | |
| | | | | | Peas with pods | 40 | 1 | | | | | |
| | | | | | Plums | 319 | 1 | | | | | |
| | | | | | Ruccola | 12 | 1 | | | | | |
| | | | | | Table grapes | 459 | 3 | | | | | |
| | | | | | Tomatoes | 249 | 1 | | | | | |
| | | | | | Witloof | 3 | 1 | | | | | |
| | | | | | Cyhalothrin, lambda- | DK | 4739 | 4735 | 4 | Currants, black | 11 | 1 |
| | | | | | | | | | | Kale | 9 | 1 |
| | | | | | | | | | | Savoy cabbage | 6 | 2 |
| | | | | | Cyhalothrin, lambda- | F | 10185 | 10008 | 177 | Apples | 394 | 11 |
| | | | | | | | | | | Apricots | 31 | 1 |
| Beans with pods | 305 | 8 | | | | | | | | | | |
| Blueberries | 46 | 1 | | | | | | | | | | |
| Brussels sprouts | 15 | 1 | | | | | | | | | | |
| Carambola | 103 | 4 | | | | | | | | | | |
| Celery | 21 | 4 | | | | | | | | | | |
| Cherries | 24 | 2 | | | | | | | | | | |
| Currants, black | 17 | 1 | | | | | | | | | | |
| Kale | 2 | 1 | | | | | | | | | | |
| Lettuce | 208 | 14 | | | | | | | | | | |
| Mandarins, clementines | 411 | 14 | | | | | | | | | | |
| Mango | 58 | 1 | | | | | | | | | | |
| Mangosteens | 12 | 1 | | | | | | | | | | |
| Melons | 311 | 9 | | | | | | | | | | |
| Nectarines | 212 | 13 | | | | | | | | | | |
| Oranges | 491 | 12 | | | | | | | | | | |
| Papayas | 89 | 2 | | | | | | | | | | |
| Passion fruits | 38 | 2 | | | | | | | | | | |
| Peaches | 158 | 9 | | | | | | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|--------------------|---|---------------------------------|---|---|----------------------------|---------------------------------|---|
| Cypermethrin (sum) | DK | 5471 | 5448 | 23 | Pears | 301 | 4 |
| | | | | | Peas with pods | 40 | 2 |
| | | | | | Peppers, sweet | 385 | 4 |
| | | | | | Persimmon | 108 | 5 |
| | | | | | Plums | 319 | 1 |
| | | | | | Pomegranate | 16 | 1 |
| | | | | | Rambutan | 47 | 2 |
| | | | | | Spinach | 163 | 15 |
| | | | | | Spring onions | 32 | 2 |
| | | | | | Strawberries | 266 | 7 |
| | | | | | Table grapes | 459 | 17 |
| | | | | | Tamarillo | 20 | 1 |
| | | | | | Tomatoes | 249 | 2 |
| | | | | | Turnips | 2 | 1 |
| | | | | | Gojiberries, dried | 1 | 1 |
| | | | | | Pears (organic) | 35 | 1 |
| | | | | | Apples | 213 | 2 |
| | | | | | Celery | 11 | 1 |
| | | | | | Currants, black | 11 | 4 |
| | | | | | Currants, red | 9 | 1 |
| | | | | | Kale | 9 | 2 |
| | | | | | Lettuce | 163 | 2 |
| | | | | | Onions | 292 | 1 |
| Plums | 72 | 3 | | | | | |
| Ruccola | 8 | 2 | | | | | |
| Spinach | 81 | 5 | | | | | |
| Cypermethrin (sum) | F | 11309 | 11095 | 214 | Apples | 394 | 1 |
| | | | | | Apricots | 31 | 4 |
| | | | | | Asian cabbage,(unspecific) | 2 | 1 |
| | | | | | Aubergines | 87 | 4 |
| | | | | | Bananas | 425 | 1 |
| | | | | | Basil | 11 | 1 |
| | | | | | Basil (Ocimum sanctum) | 2 | 1 |
| | | | | | Beans with pods | 305 | 16 |
| | | | | | Blueberries | 46 | 2 |
| | | | | | Broccoli | 63 | 1 |
| | | | | | Carambola | 103 | 20 |
| | | | | | Celery | 21 | 1 |
| | | | | | Cherries | 24 | 4 |
| | | | | | Chilli peppers | 42 | 13 |
| | | | | | Chives | 6 | 1 |
| | | | | | Coriander, leaves | 9 | 1 |
| | | | | | Courgettes | 63 | 2 |
| | | | | | Cucumbers | 218 | 2 |
| | | | | | Currants, black | 17 | 2 |
| | | | | | Gooseberries | 7 | 1 |
| | | | | | Grapefruit | 388 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|------------|---|---------------------------------|---|---|-------------------------|---------------------------------|---|
| | | | | | Guava | 9 | 1 |
| | | | | | Jambolan | 2 | 1 |
| | | | | | Lemons | 380 | 3 |
| | | | | | Lettuce | 208 | 2 |
| | | | | | Limes | 52 | 2 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Mango | 58 | 1 |
| | | | | | Melons | 311 | 4 |
| | | | | | Nectarines | 212 | 6 |
| | | | | | Okra (Lady's fingers) | 15 | 3 |
| | | | | | Oranges | 491 | 5 |
| | | | | | Passion fruits | 38 | 12 |
| | | | | | Peaches | 158 | 5 |
| | | | | | Pears | 301 | 1 |
| | | | | | Peas with pods | 40 | 8 |
| | | | | | Peppers, sweet | 385 | 6 |
| | | | | | Pitaya | 19 | 6 |
| | | | | | Plums | 319 | 5 |
| | | | | | Pomegranate | 16 | 1 |
| | | | | | Pomelos | 61 | 5 |
| | | | | | Potatoes, new | 79 | 1 |
| | | | | | Rambutan | 47 | 20 |
| | | | | | Raspberries | 101 | 3 |
| | | | | | Rhubarbs | 8 | 1 |
| | | | | | Sapota | 2 | 1 |
| | | | | | Spinach | 163 | 5 |
| | | | | | Spring onions | 32 | 1 |
| | | | | | Strawberries | 266 | 1 |
| | | | | | Table grapes | 459 | 4 |
| | | | | | Tamarillo | 20 | 6 |
| | | | | | Teas | 24 | 3 |
| | | | | | Tomatoes | 249 | 7 |
| | | | | | Vegetables, unspecified | 3 | 1 |
| | | | | | Apricot, dried | 7 | 1 |
| | | | | | Gojiberries, dried | 1 | 1 |
| Cyprodinil | DK | 3027 | 3005 | 22 | Carrots | 340 | 1 |
| | | | | | Strawberries | 163 | 21 |
| Cyprodinil | F | 9827 | 9443 | 384 | Apples | 394 | 1 |
| | | | | | Apricots | 31 | 1 |
| | | | | | Aubergines | 87 | 7 |
| | | | | | Bananas | 425 | 1 |
| | | | | | Beans with pods | 305 | 34 |
| | | | | | Blackberries | 35 | 1 |
| | | | | | Blueberries | 46 | 5 |
| | | | | | Cherries | 24 | 1 |
| | | | | | Chilli peppers | 42 | 2 |
| | | | | | Cucumbers | 218 | 26 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|----------------|---|---------------------------------|---|---|--------------------|---------------------------------|---|
| | | | | | Currants, red | 28 | 3 |
| | | | | | Fennel | 11 | 1 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Lemons | 380 | 2 |
| | | | | | Lettuce | 208 | 18 |
| | | | | | Melons | 311 | 4 |
| | | | | | Mixed berries | 2 | 1 |
| | | | | | Nectarines | 212 | 12 |
| | | | | | Oranges | 491 | 1 |
| | | | | | Peaches | 158 | 4 |
| | | | | | Pears | 301 | 33 |
| | | | | | Peppers, sweet | 385 | 11 |
| | | | | | Plums | 319 | 6 |
| | | | | | Raspberries | 101 | 16 |
| | | | | | Spinach | 163 | 2 |
| | | | | | Spring onions | 32 | 3 |
| | | | | | Strawberries | 266 | 73 |
| | | | | | Table grapes | 459 | 79 |
| | | | | | Tomatoes | 249 | 32 |
| | | | | | Cyromazine | F | 4516 |
| Aubergines | 32 | 2 | | | | | |
| Celery | 5 | 1 | | | | | |
| Cucumbers | 89 | 1 | | | | | |
| Melons | 139 | 9 | | | | | |
| Peas with pods | 26 | 1 | | | | | |
| Tomatoes | 91 | 3 | | | | | |
| Carrots | 158 | 1 | | | | | |
| Passion fruits | 38 | 1 | | | | | |
| Peppers, sweet | 385 | 1 | | | | | |
| DDT (sum) | F | 11309 | 11303 | 6 | Potatoes, new | 79 | 1 |
| | | | | | Spinach | 163 | 1 |
| | | | | | Potatoes (organic) | 12 | 1 |
| | | | | | Spelt | 30 | 1 |
| | | | | | Wheat flour | 78 | 1 |
| | | | | | Apples | 394 | 2 |
| Deltamethrin | DK | 5471 | 5469 | 2 | Apricots | 31 | 3 |
| | | | | | Avocados | 39 | 1 |
| Deltamethrin | F | 11309 | 11203 | 106 | Beans with pods | 305 | 13 |
| | | | | | Blackberries | 35 | 1 |
| | | | | | Carambola | 103 | 1 |
| | | | | | Cucumbers | 218 | 1 |
| | | | | | Currants, red | 28 | 2 |
| | | | | | Figs, fresh | 13 | 1 |
| | | | | | Lettuce | 208 | 2 |
| | | | | | Mangosteens | 12 | 1 |
| | | | | | Nectarines | 212 | 1 |
| | | | | | Oranges | 491 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|------------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Diazinon | F | 10951 | 10912 | 39 | Peaches | 158 | 1 |
| | | | | | Peas with pods | 40 | 2 |
| | | | | | Peppers, sweet | 385 | 9 |
| | | | | | Ruccola | 12 | 4 |
| | | | | | Spinach | 163 | 14 |
| | | | | | Spring onions | 32 | 1 |
| | | | | | Strawberries | 266 | 4 |
| | | | | | Table grapes | 459 | 3 |
| | | | | | Teas | 24 | 1 |
| | | | | | Tomatoes | 249 | 11 |
| | | | | | Cornflour | 8 | 1 |
| | | | | | Maize | 41 | 18 |
| | | | | | Rice, white | 167 | 4 |
| | | | | | Rolled oat | 86 | 1 |
| | | | | | Wheat flour | 184 | 1 |
| | | | | | Wheat kernels | 143 | 1 |
| | | | | | Apples | 394 | 5 |
| | | | | | Cherries | 24 | 2 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Grapefruit | 388 | 10 |
| | | | | | Kiwi | 363 | 7 |
| | | | | | Lemons | 380 | 4 |
| Mandarins, clementines | 411 | 3 | | | | | |
| Melons | 311 | 1 | | | | | |
| Oranges | 491 | 4 | | | | | |
| Pears | 301 | 1 | | | | | |
| Pineapples | 55 | 1 | | | | | |
| Dichlofluanid | F | 10951 | 10949 | 2 | Pears | 301 | 1 |
| Dichlorprop | F | 4516 | 4514 | 2 | Raspberries | 101 | 1 |
| | | | | | Bananas | 157 | 1 |
| Dichlorvos | F | 9180 | 9179 | 1 | Oranges | 173 | 1 |
| | | | | | Cucumbers | 182 | 1 |
| Dicloran | F | 11309 | 11301 | 8 | Carrots | 158 | 3 |
| | | | | | Lettuce | 208 | 2 |
| Dicofol (sum) | F | 10814 | 10620 | 194 | Peaches | 158 | 1 |
| | | | | | Pears | 301 | 1 |
| | | | | | Table grapes | 459 | 1 |
| | | | | | Basil | 11 | 1 |
| | | | | | Beans with pods | 305 | 1 |
| | | | | | Chilli peppers | 42 | 5 |
| | | | | | Grapefruit | 388 | 4 |
| | | | | | Lemons | 380 | 68 |
| | | | | | Lettuce | 208 | 1 |
| | | | | | Mandarins, clementines | 411 | 55 |
| | | | | | Melons | 311 | 6 |
| Nectarines | 212 | 1 | | | | | |
| Oranges | 491 | 31 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|------------------|---|---------------------------------|---|---|-------------------|---------------------------------|---|
| Diethofencarb | F | 7145 | 7141 | 4 | Peas without pods | 43 | 1 |
| | | | | | Plums | 319 | 2 |
| Difenoconazole | DK | 3179 | 3178 | 1 | Pomelos | 61 | 6 |
| | | | | | Table grapes | 459 | 3 |
| Difenoconazole | F | 9827 | 9758 | 69 | Teas | 24 | 4 |
| | | | | | Tomatoes | 249 | 1 |
| Diflufenican | F | 9948 | 9945 | 3 | Watermelon | 40 | 3 |
| | | | | | Marmelade, orange | 19 | 1 |
| Dimethoate (sum) | DK | 3550 | 3531 | 19 | Pears | 202 | 3 |
| | | | | | Tomatoes | 145 | 1 |
| | | | | | Currants, red | 9 | 1 |
| | | | | | Apples | 394 | 2 |
| | | | | | Beans with pods | 305 | 2 |
| | | | | | Carambola | 103 | 4 |
| | | | | | Carrots | 158 | 5 |
| | | | | | Celeriac | 7 | 1 |
| | | | | | Celery | 21 | 7 |
| | | | | | Chervil | 3 | 1 |
| | | | | | Chilli peppers | 42 | 3 |
| | | | | | Coriander, leaves | 9 | 3 |
| | | | | | Dill | 3 | 2 |
| | | | | | Fennel | 11 | 3 |
| | | | | | Lettuce | 208 | 2 |
| | | | | | Melons | 311 | 2 |
| | | | | | Nectarines | 212 | 1 |
| | | | | | Papayas | 89 | 6 |
| | | | | | Parsley Root | 6 | 2 |
| | | | | | Passion fruits | 38 | 2 |
| | | | | | Peaches | 158 | 1 |
| | | | | | Pears | 301 | 6 |
| | | | | | Peas with pods | 40 | 4 |
| | | | | | Pitaya | 19 | 4 |
| | | | | | Plums | 319 | 1 |
| | | | | | Table grapes | 459 | 1 |
| | | | | | Tamarillo | 20 | 1 |
| | | | | | Tomatoes | 249 | 3 |
| | | | | | Peaches | 158 | 1 |
| | | | | | Pears | 301 | 1 |
| | | | | | Strawberries | 266 | 1 |
| | | | | | Apples | 214 | 5 |
| | | | | | Carrots | 340 | 1 |
| | | | | | Cauliflowers | 34 | 1 |
| | | | | | Head cabbage, red | 31 | 1 |
| | | | | | Lettuce | 163 | 5 |
| | | | | | Plums | 72 | 3 |
| | | | | | Spring onions | 3 | 1 |
| | | | | | Rye kernels | 59 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Dimethoate (sum) | F | 10403 | 10332 | 71 | Wheat kernels | 107 | 1 |
| | | | | | Apples | 393 | 7 |
| | | | | | Apricots | 31 | 1 |
| | | | | | Beans with pods | 305 | 10 |
| | | | | | Cherries | 24 | 2 |
| | | | | | Chilli peppers | 42 | 3 |
| | | | | | Coriander, leaves | 9 | 1 |
| | | | | | Currants, black | 17 | 1 |
| | | | | | Jambolan | 2 | 2 |
| | | | | | Lemons | 381 | 1 |
| | | | | | Lettuce | 208 | 11 |
| | | | | | Limes | 52 | 1 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Oranges | 491 | 4 |
| | | | | | Passion fruits | 38 | 1 |
| | | | | | Peas with pods | 40 | 17 |
| | | | | | Persimmon | 109 | 1 |
| | | | | | Sapota | 2 | 1 |
| | | | | | Spinach | 163 | 3 |
| | | | | | Spring onions | 32 | 2 |
| Vegetables, unspecified | 3 | 1 | | | | | |
| Dimethomorph | DK | 1597 | 1594 | 3 | Ruccola | 3 | 1 |
| | | | | | Spring onions | 1 | 1 |
| | | | | | Strawberries | 75 | 1 |
| Dimethomorph | F | 4516 | 4458 | 58 | Beans with pods | 121 | 1 |
| | | | | | Blackberries | 15 | 1 |
| | | | | | Chives | 4 | 1 |
| | | | | | Cucumbers | 89 | 8 |
| | | | | | Lettuce | 96 | 6 |
| | | | | | Melons | 139 | 3 |
| | | | | | Ruccola | 2 | 1 |
| | | | | | Spring onions | 15 | 3 |
| | | | | | Strawberries | 105 | 1 |
| | | | | | Table grapes | 193 | 16 |
| | | | | | Tomatoes | 91 | 3 |
| | | | | | Wine, red | 117 | 14 |
| Diniconazole | F | 1689 | 1688 | 1 | Table grapes | 58 | 1 |
| Diphenylamine | DK | 3881 | 3880 | 1 | Strawberries | 163 | 1 |
| Diphenylamine | F | 10951 | 10838 | 113 | Apples | 394 | 94 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Mandarins, clementines | 411 | 3 |
| | | | | | Oranges | 491 | 5 |
| | | | | | Pears | 301 | 7 |
| | | | | | Pomegranate | 16 | 1 |
| | | | | | Oranges, juice | 44 | 2 |
| | | | | | Dithiocarbamates | DK | 1880 |
| | | | | | Cucumbers | 194 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|------------------|---|---------------------------------|---|---|-------------------|---------------------------------|---|
| Dithiocarbamates | F | 6022 | 5604 | 418 | Currants, black | 5 | 2 |
| | | | | | Currants, red | 6 | 3 |
| | | | | | Gherkin | 3 | 1 |
| | | | | | Lettuce | 155 | 2 |
| | | | | | Pears | 165 | 15 |
| | | | | | Peas with pods | 5 | 1 |
| | | | | | Plums | 70 | 8 |
| | | | | | Spinach | 74 | 2 |
| | | | | | Apples | 370 | 39 |
| | | | | | Apricots | 31 | 9 |
| | | | | | Aubergines | 80 | 1 |
| | | | | | Bananas | 405 | 1 |
| | | | | | Basil | 4 | 1 |
| | | | | | Beans with pods | 266 | 11 |
| | | | | | Blueberries | 38 | 1 |
| | | | | | Celery | 18 | 1 |
| | | | | | Chilli peppers | 31 | 6 |
| | | | | | Courgettes | 60 | 4 |
| | | | | | Cucumbers | 211 | 30 |
| | | | | | Currants, red | 19 | 5 |
| | | | | | Figs, fresh | 12 | 3 |
| | | | | | Leek | 60 | 10 |
| | | | | | Lettuce | 198 | 24 |
| | | | | | Litchi | 5 | 1 |
| | | | | | Mango | 50 | 1 |
| | | | | | Melons | 293 | 5 |
| | | | | | Nectarines | 204 | 13 |
| | | | | | Papayas | 83 | 31 |
| | | | | | Passion fruits | 36 | 11 |
| | | | | | Peaches | 149 | 28 |
| | | | | | Pears | 283 | 85 |
| | | | | | Peas with pods | 37 | 24 |
| | | | | | Peppers, sweet | 363 | 6 |
| | | | | | Persimmon | 99 | 1 |
| | | | | | Pineapples | 53 | 1 |
| | | | | | Pitaya | 15 | 3 |
| | | | | | Plums | 307 | 5 |
| | | | | | Pomegranate | 14 | 1 |
| | | | | | Rambutan | 44 | 1 |
| | | | | | Raspberries | 25 | 2 |
| | | | | | Spinach | 107 | 2 |
| Strawberries | 165 | 2 | | | | | |
| Table grapes | 444 | 33 | | | | | |
| Tamarillo | 18 | 3 | | | | | |
| Tomatoes | 240 | 12 | | | | | |
| Watermelon | 35 | 1 | | | | | |
| Endosulfan (sum) | DK | 5471 | 5470 | 1 | Parsley (organic) | 3 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues |
|------------------|---|----------------------------|--|--|-------------------------------------|----------------------------|--|
| Endosulfan (sum) | F | 11309 | 11199 | 110 | Apples | 394 | 6 |
| | | | | | Apricots | 31 | 1 |
| | | | | | Avocados | 39 | 1 |
| | | | | | Beans with pods | 305 | 1 |
| | | | | | Beans, white | 23 | 1 |
| | | | | | Blackberries | 35 | 1 |
| | | | | | Blueberries | 46 | 1 |
| | | | | | Carrots | 158 | 1 |
| | | | | | Chilli peppers | 42 | 5 |
| | | | | | Chives | 6 | 1 |
| | | | | | Courgettes | 63 | 5 |
| | | | | | Cucumbers | 218 | 2 |
| | | | | | Currants, black | 17 | 1 |
| | | | | | Guava | 9 | 2 |
| | | | | | Lemons | 380 | 2 |
| | | | | | Lettuce | 208 | 1 |
| | | | | | Melons | 311 | 33 |
| | | | | | Nectarines | 212 | 1 |
| | | | | | Papayas | 89 | 1 |
| | | | | | Passion fruits | 38 | 1 |
| | | | | | Peaches | 158 | 1 |
| | | | | | Peppers, sweet | 385 | 13 |
| | | | | | Pineapples | 55 | 1 |
| | | | | | Pomegranate | 16 | 1 |
| | | | | | Radish | 8 | 1 |
| | | | | | Strawberries | 266 | 5 |
| | | | | | Table grapes | 459 | 1 |
| | | | | | Teas | 24 | 2 |
| | | | | | Tomatoes | 249 | 14 |
| | | | | | Watermelon | 40 | 2 |
| | | | | | Marmelade, orange | 19 | 1 |
| | | | | | Brussels sprouts | 13 | 1 |
| | | | | | Grapefruit | 155 | 1 |
| Ethion | F | 10951 | 10942 | 9 | Aubergines | 87 | 1 |
| | | | | | Chilli peppers | 42 | 2 |
| | | | | | Grapefruit | 388 | 2 |
| Ethoxyquin | F | 6057 | 6056 | 1 | Guava | 9 | 1 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Pomegranate | 16 | 1 |
| Famoxadone | F | 2886 | 2876 | 10 | Table grapes | 459 | 1 |
| | | | | | Pears | 201 | 1 |
| | | | | | Beans with pods | 81 | 1 |
| | | | | | Cantharelle (Cantharellus cibarius) | 4 | 1 |
| | | | | | Leek | 18 | 1 |
| | | | | | Papayas | 32 | 1 |
| Table grapes | 127 | 5 | | | | | |
| Tomatoes | 65 | 1 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues |
|------------------------|---|----------------------------|--|--|------------------------|----------------------------|--|
| Fenarimol | F | 10951 | 10936 | 15 | Currants, black | 17 | 1 |
| | | | | | Gooseberries | 7 | 2 |
| | | | | | Peaches | 158 | 1 |
| | | | | | Strawberries | 266 | 5 |
| | | | | | Table grapes | 459 | 3 |
| | | | | | Tomatoes | 249 | 3 |
| | | | | | | | |
| Fenazaquin | F | 4516 | 4510 | 6 | Apples | 124 | 3 |
| | | | | | Mandarins, clementines | 160 | 1 |
| | | | | | Nectarines | 90 | 1 |
| | | | | | Pears | 123 | 1 |
| Fenbuconazole | F | 1689 | 1687 | 2 | Grapefruit | 69 | 1 |
| | | | | | Nectarines | 33 | 1 |
| Fenhexamid | DK | 2487 | 2442 | 45 | Cherries | 2 | 1 |
| | | | | | Plums | 60 | 1 |
| | | | | | Strawberries | 149 | 40 |
| | | | | | Tomatoes | 164 | 1 |
| Fenhexamid | F | 8407 | 8012 | 395 | Wine, red | 2 | 2 |
| | | | | | Beans with pods | 285 | 1 |
| | | | | | Blackberries | 20 | 1 |
| | | | | | Blueberries | 46 | 7 |
| | | | | | Cucumbers | 177 | 4 |
| | | | | | Currants, black | 13 | 1 |
| | | | | | Currants, red | 17 | 5 |
| | | | | | Grapefruit | 354 | 1 |
| | | | | | Kiwi | 327 | 102 |
| | | | | | Lettuce | 184 | 4 |
| | | | | | Mixed berries | 2 | 1 |
| | | | | | Nectarines | 192 | 16 |
| | | | | | Oranges | 408 | 1 |
| | | | | | Parsley | 4 | 1 |
| | | | | | Peaches | 136 | 11 |
| | | | | | Peppers, sweet | 346 | 4 |
| | | | | | Pineapples | 27 | 1 |
| | | | | | Plums | 286 | 9 |
| | | | | | Raspberries | 82 | 18 |
| | | | | | Strawberries | 225 | 47 |
| | | | | | Table grapes | 423 | 108 |
| | | | | | Teas | 25 | 1 |
| | | | | | Tomatoes | 180 | 10 |
| Wine, red | 265 | 39 | | | | | |
| Kiwi (organic) | 22 | 1 | | | | | |
| Table grapes (organic) | 29 | 1 | | | | | |
| Fenitrothion | F | 10951 | 10923 | 28 | Apples | 394 | 4 |
| | | | | | Blueberries | 46 | 1 |
| | | | | | Celery | 21 | 2 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Currants, black | 17 | 2 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues | | | | | |
|---------------------------------------|---|----------------------------|--|--|---------------------------------------|----------------------------|--|-------|---|--------------|-----|---|
| Fenoxaprop-P-Ethyl | F | 9956 | 9955 | 1 | Currants, red | 28 | 1 | | | | | |
| | | | | | Kiwi | 363 | 1 | | | | | |
| | | | | | Mandarins, clementines | 411 | 2 | | | | | |
| | | | | | Nectarines | 212 | 2 | | | | | |
| | | | | | Oranges | 491 | 1 | | | | | |
| | | | | | Peaches | 158 | 2 | | | | | |
| | | | | | Pears | 301 | 1 | | | | | |
| | | | | | Plums | 319 | 2 | | | | | |
| | | | | | Spinach | 163 | 1 | | | | | |
| | | | | | Table grapes | 459 | 5 | | | | | |
| | | | | | Peppers, sweet | 385 | 1 | | | | | |
| | | | | | Fenpropathrin | F | 10951 | 10946 | 5 | Apricots | 31 | 1 |
| | | | | | | | | | | Grapefruit | 388 | 1 |
| | | | | | Fenpropimorph | F | 9827 | 9821 | 6 | Oranges | 491 | 2 |
| Peppers, sweet | 385 | 1 | | | | | | | | | | |
| Bananas | 425 | 4 | | | | | | | | | | |
| Fenson | F | 11309 | 11308 | 1 | Potatoes, new | 79 | 1 | | | | | |
| | | | | | Strawberries | 266 | 1 | | | | | |
| Fenthion (sum) | F | 10935 | 10893 | 42 | Lemons | 380 | 1 | | | | | |
| | | | | | Mandarins, clementines | 411 | 18 | | | | | |
| | | | | | Oranges | 491 | 10 | | | | | |
| | | | | | Peaches | 158 | 2 | | | | | |
| | | | | | Persimmon | 109 | 8 | | | | | |
| | | | | | Plums | 319 | 2 | | | | | |
| | | | | | Marmelade, orange | 19 | 2 | | | | | |
| Fenvalerat, esfenvalerat, RR- and SS- | F | 11309 | 11295 | 14 | Apples | 394 | 4 | | | | | |
| | | | | | Aubergines | 87 | 1 | | | | | |
| | | | | | Beans with pods | 305 | 1 | | | | | |
| | | | | | Blackberries | 35 | 1 | | | | | |
| | | | | | Carambola | 103 | 1 | | | | | |
| | | | | | Mandarins, clementines | 411 | 1 | | | | | |
| | | | | | Oranges | 491 | 1 | | | | | |
| | | | | | Pak-choi | 2 | 1 | | | | | |
| | | | | | Teas | 24 | 3 | | | | | |
| | | | | | Fenvalerat, esfenvalerat, RS- and SR- | F | 11309 | 11300 | 9 | Aubergines | 87 | 1 |
| Beans with pods | 305 | 1 | | | | | | | | | | |
| Carambola | 103 | 1 | | | | | | | | | | |
| Kiwi | 363 | 1 | | | | | | | | | | |
| Mandarins, clementines | 411 | 1 | | | | | | | | | | |
| Pak-choi | 2 | 1 | | | | | | | | | | |
| Teas | 24 | 3 | | | | | | | | | | |
| Flucythrinate | F | 9871 | 9869 | 2 | | | | | | Grapefruit | 388 | 1 |
| | | | | | | | | | | Table grapes | 459 | 1 |
| Fludioxonil | DK | 3179 | 3168 | 11 | Currants, red | 9 | 1 | | | | | |
| Fludioxonil | F | 9827 | 9587 | 240 | Strawberries | 163 | 10 | | | | | |
| | | | | | Apples | 394 | 10 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| | | | | | Apricots | 31 | 2 |
| | | | | | Aubergines | 87 | 2 |
| | | | | | Beans with pods | 305 | 3 |
| | | | | | Blackberries | 35 | 2 |
| | | | | | Blueberries | 46 | 4 |
| | | | | | Chilli peppers | 42 | 2 |
| | | | | | Cucumbers | 218 | 8 |
| | | | | | Currants, red | 28 | 3 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Lettuce | 208 | 5 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Nectarines | 212 | 8 |
| | | | | | Oranges | 491 | 4 |
| | | | | | Peaches | 158 | 2 |
| | | | | | Pears | 301 | 28 |
| | | | | | Peppers, sweet | 385 | 18 |
| | | | | | Plums | 319 | 4 |
| | | | | | Pomegranate | 16 | 2 |
| | | | | | Raspberries | 101 | 11 |
| | | | | | Spinach | 163 | 2 |
| | | | | | Strawberries | 266 | 50 |
| | | | | | Table grapes | 459 | 51 |
| | | | | | Tomatoes | 249 | 15 |
| | | | | | Wine, red | 181 | 1 |
| Fluoxastrobin | F | 3975 | 3973 | 2 | Mandarins, clementines | 160 | 2 |
| Fluroxypyr (sum) | F | 3975 | 3973 | 2 | Peaches | 64 | 1 |
| | | | | | Table grapes | 193 | 1 |
| Flusilazole | F | 7953 | 7948 | 5 | Beans with pods | 269 | 1 |
| | | | | | Currants, black | 9 | 1 |
| | | | | | Peppers, sweet | 336 | 1 |
| | | | | | Sweet corn (small) | 30 | 1 |
| | | | | | Table grapes | 409 | 1 |
| Flutriafol | F | 1689 | 1678 | 11 | Peppers, sweet | 62 | 11 |
| Fluvalinate, tau- | F | 3349 | 3344 | 5 | Apples | 180 | 1 |
| | | | | | Grapefruit | 96 | 2 |
| | | | | | Mandarins, clementines | 139 | 1 |
| | | | | | Peppers, sweet | 100 | 1 |
| Glyphosate | DK | 680 | 669 | 11 | Barley (malting) | 1 | 1 |
| | | | | | Barley kernels | 7 | 1 |
| | | | | | Oat kernels | 9 | 2 |
| | | | | | Spelt, flour | 16 | 1 |
| | | | | | Wheat kernels | 183 | 6 |
| Glyphosate | F | 864 | 848 | 16 | Chick pea | 7 | 1 |
| | | | | | Lentils | 15 | 5 |
| | | | | | Barley grit | 13 | 2 |
| | | | | | Rolled oat | 83 | 3 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues |
|-------------------|---|----------------------------|--|--|------------------------|----------------------------|--|
| HCH (sum) | DK | 3729 | 3728 | 1 | Spelt, flour | 22 | 1 |
| | | | | | Wheat flour | 181 | 2 |
| | | | | | Wheat kernels | 141 | 2 |
| HCH (sum) | F | 10951 | 10950 | 1 | Carrots | 340 | 1 |
| Hexachlorobenzene | DK | 3911 | 3908 | 3 | Ginger | 28 | 1 |
| Hexachlorobenzene | F | 10951 | 10950 | 1 | Carrots | 340 | 1 |
| | | | | | Courgettes | 20 | 1 |
| | | | | | Parsnips | 51 | 1 |
| Hexaconazole | F | 7843 | 7835 | 8 | Parsnips | 10 | 1 |
| Hexythiazox | F | 4516 | 4499 | 17 | Celery | 13 | 1 |
| | | | | | Chilli peppers | 20 | 2 |
| | | | | | Coriander, leaves | 6 | 2 |
| | | | | | Mint leaves | 1 | 1 |
| | | | | | Peppers, sweet | 336 | 1 |
| | | | | | Table grapes | 409 | 1 |
| | | | | | Blackberries | 15 | 1 |
| | | | | | Grapefruit | 155 | 1 |
| | | | | | Lemons | 158 | 4 |
| | | | | | Mandarins, clementines | 160 | 4 |
| | | | | | Peppers, sweet | 170 | 1 |
| | | | | | Raspberries | 43 | 3 |
| | | | | | Strawberries | 105 | 1 |
| | | | | | Table grapes | 193 | 1 |
| | | | | | Tomatoes | 91 | 1 |
| | | | | | Cucumbers | 199 | 11 |
| | | | | | Imazalil | DK | 3363 |
| Imazalil | F | 10310 | 8329 | 1981 | Apples | 392 | 5 |
| | | | | | Bananas | 425 | 309 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Grapefruit | 388 | 340 |
| | | | | | Kiwi | 361 | 3 |
| | | | | | Lemons | 381 | 321 |
| | | | | | Limes | 52 | 41 |
| | | | | | Mandarins, clementines | 410 | 386 |
| | | | | | Melons | 308 | 69 |
| | | | | | Mineola | 3 | 3 |
| | | | | | Oranges | 491 | 448 |
| | | | | | Pears | 299 | 13 |
| | | | | | Pineapples | 56 | 3 |
| | | | | | Plums | 317 | 2 |
| | | | | | Pomelos | 60 | 27 |
| | | | | | Strawberries | 265 | 1 |
| | | | | | Watermelon | 39 | 3 |
| Marmelade, orange | 19 | 3 | | | | | |
| Oranges, juice | 43 | 1 | | | | | |
| Limes (organic) | 6 | 1 | | | | | |
| Oranges (organic) | 30 | 1 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues |
|-----------------|---|----------------------------|--|--|------------------------|----------------------------|--|
| Iprodione | DK | 3729 | 3719 | 10 | Carrots | 340 | 7 |
| | | | | | Spinach | 81 | 1 |
| | | | | | Tomatoes | 212 | 2 |
| Iprodione | F | 10951 | 10510 | 441 | Apples | 394 | 18 |
| | | | | | Apricots | 31 | 4 |
| | | | | | Asparagus, green | 38 | 1 |
| | | | | | Beans with pods | 305 | 21 |
| | | | | | Blackberries | 35 | 2 |
| | | | | | Blueberries | 46 | 3 |
| | | | | | Boysenberries | 3 | 1 |
| | | | | | Carrots | 158 | 4 |
| | | | | | Cherries | 24 | 1 |
| | | | | | Chilli peppers | 42 | 3 |
| | | | | | Cucumbers | 218 | 10 |
| | | | | | Currants, red | 28 | 11 |
| | | | | | Head cabbage, red | 7 | 1 |
| | | | | | Kiwi | 363 | 37 |
| | | | | | Lemons | 380 | 1 |
| | | | | | Lettuce | 208 | 16 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Melons | 311 | 2 |
| | | | | | Nectarines | 212 | 27 |
| | | | | | Pak-choi | 2 | 2 |
| | | | | | Passion fruits | 38 | 1 |
| | | | | | Peaches | 158 | 11 |
| | | | | | Pears | 301 | 7 |
| | | | | | Peas without pods | 43 | 2 |
| | | | | | Peppers, sweet | 385 | 4 |
| | | | | | Pitaya | 19 | 2 |
| | | | | | Plums | 319 | 97 |
| | | | | | Pomelos | 61 | 1 |
| | | | | | Radish | 8 | 1 |
| | | | | | Raspberries | 101 | 7 |
| | | | | | Ruccola | 12 | 1 |
| | | | | | Spinach | 163 | 2 |
| | | | | | Spring onions | 32 | 1 |
| | | | | | Strawberries | 266 | 14 |
| | | | | | Table grapes | 459 | 91 |
| | | | | | Tomatoes | 249 | 24 |
| | | | | | Rice, white | 167 | 1 |
| | | | | | Wine, red | 181 | 8 |
| Iprovalicarb | F | 3975 | 3957 | 18 | Peas with pods | 26 | 1 |
| | | | | | Table grapes | 193 | 8 |
| | | | | | Tomatoes | 91 | 1 |
| | | | | | Wine, red | 117 | 8 |
| Kresoxim-methyl | DK | 3881 | 3877 | 4 | Apples | 213 | 1 |
| | | | | | Currants, black | 11 | 3 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues | | | | | |
|-----------------|---|----------------------------|--|--|------------------------|----------------------------|--|-------|----|-----------|-----|----|
| Kresoxim-methyl | F | 10951 | 10912 | 39 | Beans with pods | 305 | 1 | | | | | |
| | | | | | Currants, red | 28 | 5 | | | | | |
| | | | | | Passion fruits | 38 | 2 | | | | | |
| | | | | | Pears | 301 | 1 | | | | | |
| | | | | | Peppers, sweet | 385 | 5 | | | | | |
| | | | | | Strawberries | 266 | 14 | | | | | |
| | | | | | Table grapes | 459 | 11 | | | | | |
| | | | | | Lindane | F | 11309 | 11308 | 1 | Ginger | 28 | 1 |
| | | | | | Linuron | DK | 2485 | 2452 | 33 | Carrots | 281 | 22 |
| | | | | | | | | | | Celery | 11 | 1 |
| Parsley | 3 | 1 | | | | | | | | | | |
| Parsley Root | 20 | 3 | | | | | | | | | | |
| Parsnips | 36 | 4 | | | | | | | | | | |
| Spinach | 79 | 1 | | | | | | | | | | |
| Linuron | F | 8377 | 8338 | 39 | Spring onions | 3 | 1 | | | | | |
| | | | | | Apples | 276 | 1 | | | | | |
| | | | | | Bananas | 343 | 1 | | | | | |
| | | | | | Carrots | 108 | 27 | | | | | |
| | | | | | Celeriac | 5 | 1 | | | | | |
| | | | | | Celery | 14 | 1 | | | | | |
| | | | | | Kiwi | 326 | 1 | | | | | |
| | | | | | Lemons | 348 | 1 | | | | | |
| | | | | | Parsley Root | 5 | 1 | | | | | |
| | | | | | Parsnips | 8 | 1 | | | | | |
| | | | | | Pears | 242 | 1 | | | | | |
| | | | | | Raspberries | 82 | 2 | | | | | |
| | | | | | Spinach | 149 | 1 | | | | | |
| | | | | | Malathion (sum) | DK | 5371 | 5347 | 24 | Plums | 72 | 1 |
| | | | | | | | | | | Rye flour | 28 | 1 |
| Rye kernels | 100 | 4 | | | | | | | | | | |
| Spelt | 30 | 3 | | | | | | | | | | |
| Spelt, flour | 16 | 1 | | | | | | | | | | |
| Wheat flour | 78 | 3 | | | | | | | | | | |
| Wheat kernels | 187 | 11 | | | | | | | | | | |
| Malathion (sum) | F | 11435 | 11241 | 194 | Apples | 393 | 2 | | | | | |
| | | | | | Blackberries | 35 | 1 | | | | | |
| | | | | | Celery | 21 | 2 | | | | | |
| | | | | | Currants, red | 28 | 1 | | | | | |
| | | | | | Grapefruit | 388 | 9 | | | | | |
| | | | | | Kiwi | 363 | 2 | | | | | |
| | | | | | Kumquates | 14 | 5 | | | | | |
| | | | | | Lemons | 381 | 3 | | | | | |
| | | | | | Mandarins, clementines | 411 | 110 | | | | | |
| | | | | | Melons | 311 | 1 | | | | | |
| | | | | | Nectarines | 212 | 1 | | | | | |
| | | | | | Oranges | 491 | 30 | | | | | |
| | | | | | Peaches | 158 | 1 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-----------------|---|---------------------------------|---|---|------------------------------|---------------------------------|---|
| | | | | | Peppers, sweet | 385 | 5 |
| | | | | | Persimmon | 109 | 1 |
| | | | | | Strawberries | 266 | 1 |
| | | | | | Table grapes | 458 | 1 |
| | | | | | Tamarillo | 20 | 1 |
| | | | | | Cornflour | 8 | 2 |
| | | | | | Rice, white | 167 | 6 |
| | | | | | Rye flour | 57 | 1 |
| | | | | | Rye kernels | 47 | 1 |
| | | | | | Spelt, flour | 22 | 1 |
| | | | | | Wheat flour | 184 | 3 |
| | | | | | Wheat kernels | 143 | 2 |
| | | | | | Beans, kidney (organic) | 10 | 1 |
| Mecoprop (sum) | F | 4516 | 4515 | 1 | Pears | 123 | 1 |
| Mepiquat | DK | 1089 | 1087 | 2 | Barley (malting) | 1 | 1 |
| | | | | | Rye kernels | 136 | 1 |
| Mepiquat | F | 2198 | 2175 | 23 | Mushroom (Agaricus bisporus) | 12 | 2 |
| | | | | | Okra (Lady's fingers) | 4 | 1 |
| | | | | | Sweet corn (small) | 18 | 1 |
| | | | | | Rye flour | 73 | 3 |
| | | | | | Rye kernels | 73 | 1 |
| | | | | | Spelt | 13 | 3 |
| | | | | | Spelt, flour | 22 | 2 |
| | | | | | Wheat kernels | 185 | 10 |
| Metalaxyl (sum) | DK | 3802 | 3797 | 5 | Lettuce | 163 | 3 |
| | | | | | Potatoes | 424 | 1 |
| | | | | | Radish | 3 | 1 |
| Metalaxyl (sum) | F | 11103 | 11002 | 101 | Basil | 11 | 1 |
| | | | | | Basil (Ocimum sanctum) | 2 | 1 |
| | | | | | Beans with pods | 305 | 1 |
| | | | | | Chervil | 3 | 1 |
| | | | | | Chilli peppers | 42 | 5 |
| | | | | | Coriander, leaves | 9 | 2 |
| | | | | | Cucumbers | 218 | 21 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Lemons | 381 | 1 |
| | | | | | Lettuce | 208 | 14 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Melons | 311 | 1 |
| | | | | | Peas with pods | 40 | 3 |
| | | | | | Peppers, sweet | 385 | 2 |
| | | | | | Pitaya | 19 | 2 |
| | | | | | Potatoes, new | 79 | 5 |
| | | | | | Ruccola | 12 | 2 |
| | | | | | Strawberries | 266 | 2 |
| | | | | | Table grapes | 458 | 16 |
| | | | | | Tomatoes | 250 | 2 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues | | | | | |
|-------------------------|---|---------------------------------|---|---|-----------------|---------------------------------|---|-------|----|------------|-----|----|
| Methacrifos | F | 8849 | 8848 | 1 | Wine, red | 266 | 17 | | | | | |
| | | | | | Beans with pods | 292 | 1 | | | | | |
| Methamidophos | F | 10514 | 10491 | 23 | Apples | 393 | 1 | | | | | |
| | | | | | Beans with pods | 305 | 5 | | | | | |
| | | | | | Chilli peppers | 42 | 1 | | | | | |
| | | | | | Cucumbers | 218 | 2 | | | | | |
| | | | | | Figs, fresh | 13 | 1 | | | | | |
| | | | | | Melons | 311 | 2 | | | | | |
| | | | | | Nectarines | 211 | 1 | | | | | |
| | | | | | Oranges | 491 | 1 | | | | | |
| | | | | | Papayas | 89 | 1 | | | | | |
| | | | | | Peaches | 158 | 1 | | | | | |
| | | | | | Peas with pods | 40 | 1 | | | | | |
| | | | | | Peppers, sweet | 385 | 2 | | | | | |
| | | | | | Plums | 318 | 1 | | | | | |
| | | | | | Strawberries | 265 | 1 | | | | | |
| | | | | | Watermelon | 39 | 2 | | | | | |
| | | | | | Methidathion | F | 10424 | 10328 | 96 | Grapefruit | 388 | 16 |
| | | | | | | | | | | Guava | 9 | 1 |
| Lemons | 380 | 27 | | | | | | | | | | |
| Mandarins, clementines | 411 | 20 | | | | | | | | | | |
| Oranges | 491 | 28 | | | | | | | | | | |
| Pomelos | 61 | 4 | | | | | | | | | | |
| Onions | 52 | 1 | | | | | | | | | | |
| Methiocarb (sum) | F | 8845 | 8834 | 11 | Peppers, sweet | 346 | 8 | | | | | |
| | | | | | Plums | 286 | 1 | | | | | |
| | | | | | Table grapes | 423 | 1 | | | | | |
| Methomyl and Thiodicarb | F | 8948 | 8862 | 86 | Apples | 277 | 2 | | | | | |
| | | | | | Basil | 9 | 1 | | | | | |
| | | | | | Beans with pods | 285 | 7 | | | | | |
| | | | | | Boysenberries | 2 | 1 | | | | | |
| | | | | | Carambola | 97 | 18 | | | | | |
| | | | | | Chilli peppers | 26 | 1 | | | | | |
| | | | | | Cucumbers | 177 | 4 | | | | | |
| | | | | | Jambolan | 2 | 2 | | | | | |
| | | | | | Lettuce | 184 | 10 | | | | | |
| | | | | | Melons | 273 | 6 | | | | | |
| | | | | | Oranges | 408 | 1 | | | | | |
| | | | | | Papayas | 87 | 9 | | | | | |
| | | | | | Pears | 244 | 3 | | | | | |
| | | | | | Peas with pods | 40 | 2 | | | | | |
| | | | | | Peppers, sweet | 346 | 3 | | | | | |
| | | | | | Plums | 286 | 1 | | | | | |
| | | | | | Sapota | 2 | 1 | | | | | |
| | | | | | Spinach | 149 | 1 | | | | | |
| Strawberries | 225 | 4 | | | | | | | | | | |
| Table grapes | 423 | 6 | | | | | | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues | | | | | | | | | | |
|------------------------|---|---------------------------------|---|---|-------------------------|---------------------------------|---|-------|-----|------------------------|-----|------|------|---|-------------------|-----|------|
| Metribuzin | F | 2886 | 2884 | 2 | Tea, herbal | 1 | 1 | | | | | | | | | | |
| | | | | | Vegetables, unspecified | 3 | 1 | | | | | | | | | | |
| | | | | | Watermelon | 39 | 1 | | | | | | | | | | |
| | | | | | Wine, red | 91 | 2 | | | | | | | | | | |
| | | | | | Mevinphos | DK | 3762 | 3761 | 1 | Lettuce | 163 | 1 | | | | | |
| | | | | | | | | | | Cherries | 24 | 1 | | | | | |
| | | | | | Monocrotophos | F | 11078 | 11077 | 1 | Apples | 394 | 2 | | | | | |
| | | | | | | | | | | Bananas | 425 | 22 | | | | | |
| | | | | | Myclobutanil | F | 10951 | 10821 | 130 | Beans with pods | 305 | 2 | | | | | |
| | | | | | | | | | | Cherries | 24 | 4 | | | | | |
| | | | | | | | | | | Chilli peppers | 42 | 3 | | | | | |
| | | | | | | | | | | Cucumbers | 218 | 3 | | | | | |
| | | | | | | | | | | Lemons | 380 | 2 | | | | | |
| | | | | | | | | | | Mandarins, clementines | 411 | 7 | | | | | |
| | | | | | | | | | | Melons | 311 | 4 | | | | | |
| | | | | | | | | | | Nectarines | 212 | 2 | | | | | |
| | | | | | | | | | | Oranges | 491 | 7 | | | | | |
| | | | | | | | | | | Peaches | 158 | 1 | | | | | |
| | | | | | | | | | | Peppers, sweet | 385 | 8 | | | | | |
| | | | | | | | | | | Pomelos | 61 | 11 | | | | | |
| | | | | | | | | | | Strawberries | 266 | 18 | | | | | |
| | | | | | | | | | | Table grapes | 459 | 33 | | | | | |
| | | | | | | | | | | Tomatoes | 249 | 1 | | | | | |
| | | | | | | | | | | Ofurace | F | 8510 | 8509 | 1 | Wine, red | 266 | 1 |
| | | | | | | | | | | | | | | | Orthophenylphenol | F | 9827 |
| | | | | | | | | | | Beans with pods | 305 | 1 | | | | | |
| | | | | | | | | | | Grapefruit | 388 | 149 | | | | | |
| Kiwi | 363 | 1 | | | | | | | | | | | | | | | |
| Lemons | 380 | 87 | | | | | | | | | | | | | | | |
| Mandarins, clementines | 411 | 132 | | | | | | | | | | | | | | | |
| Melons | 311 | 5 | | | | | | | | | | | | | | | |
| Mineola | 3 | 2 | | | | | | | | | | | | | | | |
| Nectarines | 212 | 1 | | | | | | | | | | | | | | | |
| Oranges | 491 | 165 | | | | | | | | | | | | | | | |
| Peaches | 158 | 1 | | | | | | | | | | | | | | | |
| Peppers, sweet | 385 | 2 | | | | | | | | | | | | | | | |
| Pomelos | 61 | 3 | | | | | | | | | | | | | | | |
| Potatoes, new | 79 | 1 | | | | | | | | | | | | | | | |
| Raspberries | 101 | 3 | | | | | | | | | | | | | | | |
| Spinach | 163 | 1 | | | | | | | | | | | | | | | |
| Juice, grapefruit | 1 | 1 | | | | | | | | | | | | | | | |
| Marmelade, orange | 19 | 2 | | | | | | | | | | | | | | | |
| Pear, canned | 9 | 1 | | | | | | | | | | | | | | | |
| Oxamyl | F | 8407 | 8395 | 12 | Courgettes | 36 | 1 | | | | | | | | | | |
| | | | | | Cucumbers | 177 | 1 | | | | | | | | | | |
| | | | | | Limes | 46 | 3 | | | | | | | | | | |
| | | | | | Melons | 273 | 1 | | | | | | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Oxydemeton-methyl (sum) | F | 10448 | 10444 | 4 | Peppers, sweet | 346 | 2 |
| | | | | | Pineapples | 27 | 1 |
| | | | | | Tomatoes | 180 | 3 |
| | | | | | Apples | 392 | 1 |
| | | | | | Carambola | 103 | 1 |
| Parathion-methyl (sum) | F | 10951 | 10939 | 12 | Pears | 300 | 1 |
| | | | | | Pineapples | 55 | 1 |
| | | | | | Apples | 394 | 1 |
| | | | | | Basil | 11 | 1 |
| | | | | | Figs, fresh | 13 | 1 |
| | | | | | Grapefruit | 388 | 4 |
| | | | | | Guava | 9 | 1 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Nectarines | 212 | 1 |
| | | | | | Pomelos | 61 | 1 |
| Penconazole | F | 10951 | 10917 | 34 | Sesame seed | 18 | 1 |
| | | | | | Peaches | 158 | 1 |
| | | | | | Pears | 301 | 1 |
| | | | | | Peppers, sweet | 385 | 5 |
| | | | | | Strawberries | 266 | 4 |
| Pendimethalin | DK | 2404 | 2403 | 1 | Kale | 9 | 1 |
| Pendimethalin | F | 7145 | 7143 | 2 | Carrots | 85 | 1 |
| Pentachlorobenzene | F | 8496 | 8495 | 1 | Peppers, sweet | 285 | 1 |
| | | | | | Carrots | 132 | 1 |
| Pentachlorophenol | F | 10951 | 10950 | 1 | Grapefruit | 388 | 1 |
| Permethrin (sum) | F | 11309 | 11290 | 19 | Celery | 21 | 1 |
| | | | | | Coriander, leaves | 9 | 1 |
| | | | | | Ginger | 28 | 1 |
| | | | | | Lettuce | 208 | 1 |
| | | | | | Litchi | 7 | 1 |
| | | | | | Mango | 58 | 1 |
| | | | | | Melons | 311 | 6 |
| | | | | | Oranges | 491 | 1 |
| | | | | | Tamarillo | 20 | 2 |
| | | | | | Watermelon | 40 | 1 |
| | | | | | Witloof | 3 | 1 |
| | | | | | Rice, white | 167 | 1 |
| | | | | | Wheat kernels | 143 | 1 |
| | | | | | Phenthoate | F | 10951 |
| Phorate (sum) | F | 9950 | 9946 | 4 | Ginger | 28 | 2 |
| | | | | | Papayas | 89 | 1 |
| | | | | | Pears | 301 | 1 |
| Phosalone | DK | 3881 | 3863 | 18 | Apples | 213 | 17 |
| | | | | | Plums | 72 | 1 |
| Phosalone | F | 10951 | 10932 | 19 | Apples | 394 | 9 |
| | | | | | Gooseberries | 7 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples analysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples analysed | Number of samples with detected residues |
|-------------------|---|----------------------------|--|--|------------------------|----------------------------|--|
| Phosmet (sum) | F | 10951 | 10921 | 30 | Pears | 301 | 3 |
| | | | | | Plums | 319 | 4 |
| | | | | | Rambutan | 47 | 1 |
| | | | | | Marmelade, orange | 19 | 1 |
| | | | | | Apples | 394 | 9 |
| | | | | | Currants, red | 28 | 1 |
| | | | | | Mandarins, clementines | 411 | 4 |
| | | | | | Nectarines | 212 | 1 |
| | | | | | Oranges | 491 | 4 |
| | | | | | Peaches | 158 | 5 |
| | | | | | Pears | 301 | 1 |
| | | | | | Persimmon | 108 | 1 |
| | | | | | Plums | 319 | 2 |
| | | | | | Blueberry, dried | 1 | 1 |
| Pirimicarb (sum) | DK | 3812 | 3791 | 21 | Pears (organic) | 35 | 1 |
| | | | | | Apples | 214 | 8 |
| | | | | | Lettuce | 163 | 5 |
| | | | | | Plums | 72 | 2 |
| Pirimicarb (sum) | F | 11077 | 11051 | 26 | Strawberries | 163 | 6 |
| | | | | | Apples | 393 | 8 |
| | | | | | Beans with pods | 305 | 3 |
| | | | | | Brussels sprouts | 15 | 1 |
| | | | | | Celery | 21 | 1 |
| | | | | | Lemons | 381 | 1 |
| | | | | | Lettuce | 208 | 3 |
| | | | | | Mandarins, clementines | 411 | 2 |
| | | | | | Pears | 301 | 3 |
| | | | | | Peppers, sweet | 385 | 1 |
| Pirimiphos-methyl | DK | 5471 | 5463 | 8 | Strawberries | 266 | 3 |
| | | | | | Celeriac | 12 | 1 |
| | | | | | Maize | 1 | 1 |
| Pirimiphos-methyl | F | 11309 | 11198 | 111 | Wheat flour | 78 | 6 |
| | | | | | Chilli peppers | 42 | 2 |
| | | | | | Grapefruit | 388 | 2 |
| | | | | | Lemons | 380 | 3 |
| | | | | | Mandarins, clementines | 411 | 3 |
| | | | | | Melons | 311 | 1 |
| | | | | | Oranges | 491 | 7 |
| | | | | | Peppers, sweet | 385 | 5 |
| | | | | | Persimmon | 108 | 1 |
| | | | | | Sesame seed | 18 | 1 |
| | | | | | Cornflour | 8 | 2 |
| | | | | | Maize | 41 | 1 |
| | | | | | Rice, white | 167 | 13 |
| Rolled oat | 86 | 2 | | | | | |
| Rye flour | 57 | 6 | | | | | |
| Rye kernels | 47 | 7 | | | | | |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|------------------|---|---------------------------------|---|---|------------------------------|---------------------------------|---|
| Prochloraz (sum) | DK | 3179 | 3170 | 9 | Spelt | 13 | 1 |
| | F | 9827 | 9645 | 182 | Wheat bran | 13 | 2 |
| Prochloraz (sum) | | | | | Wheat flour | 184 | 30 |
| | | | | | Wheat kernels | 143 | 15 |
| Procymidone | | | | | Pasta | 43 | 5 |
| | | | | | Pasta, dried | 32 | 2 |
| Prochloraz (sum) | DK | 3179 | 3170 | 9 | Mushroom (Agaricus bisporus) | 41 | 9 |
| | F | 9827 | 9645 | 182 | Avocados | 39 | 1 |
| Procymidone | | | | | Grapefruit | 388 | 13 |
| | | | | | Lemons | 380 | 41 |
| Procymidone | | | | | Limes | 52 | 3 |
| | | | | | Mandarins, clementines | 411 | 35 |
| Procymidone | | | | | Mango | 58 | 18 |
| | | | | | Mushroom (Agaricus bisporus) | 35 | 8 |
| Procymidone | | | | | Oranges | 491 | 23 |
| | | | | | Papayas | 89 | 30 |
| Procymidone | | | | | Pineapples | 55 | 1 |
| | | | | | Pomelos | 61 | 9 |
| Procymidone | F | 10951 | 10799 | 152 | Apples | 394 | 1 |
| | | | | | Beans with pods | 305 | 2 |
| Procymidone | | | | | Blueberries | 46 | 1 |
| | | | | | Cauliflowers | 51 | 1 |
| Procymidone | | | | | Chilli peppers | 42 | 3 |
| | | | | | Courgettes | 63 | 2 |
| Procymidone | | | | | Cucumbers | 218 | 14 |
| | | | | | Kiwi | 363 | 3 |
| Procymidone | | | | | Lettuce | 208 | 9 |
| | | | | | Mandarins, clementines | 411 | 1 |
| Procymidone | | | | | Melons | 311 | 9 |
| | | | | | Nectarines | 212 | 1 |
| Procymidone | | | | | Peaches | 158 | 1 |
| | | | | | Pears | 301 | 1 |
| Procymidone | | | | | Peas without pods | 43 | 1 |
| | | | | | Peppers, sweet | 385 | 26 |
| Procymidone | | | | | Plums | 319 | 2 |
| | | | | | Raspberries | 101 | 8 |
| Procymidone | | | | | Spring onions | 32 | 1 |
| | | | | | Strawberries | 266 | 17 |
| Procymidone | | | | | Table grapes | 459 | 13 |
| | | | | | Tomatoes | 249 | 35 |
| Profenofos | F | 10951 | 10927 | 24 | Beans with pods | 305 | 2 |
| | | | | | Chilli peppers | 42 | 7 |
| Profenofos | | | | | Coriander, leaves | 9 | 1 |
| | | | | | Lemons | 380 | 1 |
| Profenofos | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Okra (Lady's fingers) | 15 | 1 |
| Profenofos | | | | | Oranges | 491 | 4 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Propamocarb (sum) | DK | 1597 | 1575 | 22 | Passion fruits | 38 | 3 |
| | | | | | Peas with pods | 40 | 2 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Pomelos | 61 | 1 |
| | | | | | Strawberries | 266 | 1 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Cucumbers | 77 | 21 |
| | | | | | Tomatoes | 75 | 1 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Aubergines | 32 | 2 |
| | | | | | Beans with pods | 121 | 1 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Carrots | 50 | 2 |
| | | | | | Cucumbers | 89 | 49 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Leek | 20 | 1 |
| | | | | | Lettuce | 96 | 5 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Melons | 139 | 1 |
| | | | | | Papayas | 48 | 2 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Peppers, sweet | 170 | 8 |
| | | | | | Potatoes | 40 | 3 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Radish | 3 | 1 |
| | | | | | Ruccola | 2 | 1 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Spinach | 66 | 3 |
| | | | | | Spring onions | 15 | 1 |
| Propamocarb (sum) | F | 4516 | 4428 | 88 | Strawberries | 105 | 1 |
| | | | | | Tomatoes | 91 | 7 |
| Propargite | F | 9827 | 9763 | 64 | Apples | 394 | 34 |
| | | | | | Apricots | 31 | 1 |
| Propargite | F | 9827 | 9763 | 64 | Beans with pods | 305 | 1 |
| | | | | | Grapefruit | 388 | 1 |
| Propargite | F | 9827 | 9763 | 64 | Lemons | 380 | 5 |
| | | | | | Mandarins, clementines | 411 | 3 |
| Propargite | F | 9827 | 9763 | 64 | Mixed berries | 2 | 1 |
| | | | | | Nectarines | 212 | 2 |
| Propargite | F | 9827 | 9763 | 64 | Okra (Lady's fingers) | 15 | 1 |
| | | | | | Oranges | 491 | 6 |
| Propargite | F | 9827 | 9763 | 64 | Peppers, sweet | 385 | 1 |
| | | | | | Plums | 319 | 5 |
| Propargite | F | 9827 | 9763 | 64 | Table grapes | 459 | 1 |
| | | | | | Tomatoes | 249 | 1 |
| Propiconazole | F | 11430 | 11421 | 9 | Lemons (organic) | 26 | 1 |
| | | | | | Chilli peppers | 42 | 1 |
| Propiconazole | F | 11430 | 11421 | 9 | Coriander, leaves | 9 | 2 |
| | | | | | Currants, black | 17 | 1 |
| Propiconazole | F | 11430 | 11421 | 9 | Mandarins, clementines | 411 | 1 |
| | | | | | Mango | 58 | 1 |
| Propiconazole | F | 11430 | 11421 | 9 | Peas with pods | 40 | 1 |
| | | | | | Rambutan | 47 | 1 |
| Propoxur | F | 8948 | 8947 | 1 | Tomatoes | 249 | 1 |
| | | | | | Tamarillo | 20 | 1 |
| Propyzamide | F | 10951 | 10945 | 6 | Beans with pods | 305 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|----------------|---|---------------------------------|---|---|--------------------------|---------------------------------|---|
| Prothiofos | F | 11309 | 11300 | 9 | Lettuce | 208 | 5 |
| | | | | | Aubergines | 87 | 1 |
| | | | | | Grapefruit | 388 | 2 |
| | | | | | Guava | 9 | 2 |
| Pymetrozine | F | 4516 | 4503 | 13 | Oranges | 491 | 4 |
| | | | | | Cucumbers | 89 | 6 |
| | | | | | Peppers, sweet | 170 | 4 |
| | | | | | Spinach | 66 | 1 |
| Pyraclostrobin | DK | 1597 | 1560 | 37 | Strawberries | 105 | 1 |
| | | | | | Tomatoes | 91 | 1 |
| | | | | | Apples | 69 | 9 |
| | | | | | Pears | 68 | 4 |
| Pyraclostrobin | F | 4516 | 4340 | 176 | Plums | 29 | 7 |
| | | | | | Spinach | 41 | 1 |
| | | | | | Strawberries | 75 | 16 |
| | | | | | Apples | 124 | 19 |
| | | | | | Beans with pods | 121 | 1 |
| | | | | | Blueberries | 41 | 1 |
| | | | | | Brussels sprouts | 7 | 3 |
| | | | | | Grapefruit | 155 | 24 |
| | | | | | Lemons | 158 | 4 |
| | | | | | Lettuce | 96 | 4 |
| | | | | | Limes | 18 | 1 |
| | | | | | Mandarins, clementines | 160 | 1 |
| | | | | | Nectarines | 90 | 2 |
| | | | | | Oranges | 173 | 7 |
| | | | | | Peaches | 64 | 3 |
| | | | | | Pears | 123 | 55 |
| | | | | | Peppers, sweet | 170 | 1 |
| Pomelos | 34 | 3 | | | | | |
| Raspberries | 43 | 1 | | | | | |
| Ruccola | 2 | 2 | | | | | |
| Strawberries | 105 | 18 | | | | | |
| Table grapes | 193 | 19 | | | | | |
| Tomatoes | 91 | 7 | | | | | |
| Pyrazophos | F | 11077 | 11076 | 1 | Wine, red | 266 | 1 |
| Pyrethrins | F | 9827 | 9825 | 2 | Peppers, sweet (organic) | 26 | 1 |
| | | | | | Tomatoes (organic) | 25 | 1 |
| Pyridaben | F | 7145 | 7126 | 19 | Grapefruit | 292 | 6 |
| | | | | | Lemons | 290 | 3 |
| | | | | | Mandarins, clementines | 272 | 3 |
| | | | | | Peaches | 111 | 1 |
| | | | | | Tomatoes | 145 | 5 |
| Pyridate (sum) | F | 4015 | 4014 | 1 | Gojiberries, dried | 1 | 1 |
| Pirimethanil | DK | 3091 | 3033 | 58 | Mandarins, clementines | 52 | 1 |
| | | | | | Apples | 214 | 1 |
| | | | | | Cucumbers | 199 | 20 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Pyrimethanil | F | 9429 | 9294 | 135 | Plums | 72 | 1 |
| | | | | | Strawberries | 163 | 31 |
| | | | | | Tomatoes | 212 | 5 |
| | | | | | Apples | 392 | 7 |
| | | | | | Aubergines | 86 | 1 |
| | | | | | Bananas | 425 | 2 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Cucumbers | 216 | 5 |
| | | | | | Lemons | 273 | 10 |
| | | | | | Limes | 36 | 1 |
| | | | | | Mandarins, clementines | 303 | 5 |
| | | | | | Oranges | 377 | 2 |
| | | | | | Papayas | 89 | 1 |
| | | | | | Pears | 299 | 10 |
| | | | | | Peppers, sweet | 382 | 10 |
| | | | | | Plums | 318 | 2 |
| | | | | | Raspberries | 101 | 15 |
| | | | | | Strawberries | 266 | 8 |
| | | | | | Table grapes | 457 | 30 |
| | | | | | Tomatoes | 250 | 22 |
| Marmelade, orange | 19 | 1 | | | | | |
| Wine, red | 266 | 2 | | | | | |
| Pyriproxyfen | F | 4516 | 4423 | 93 | Aubergines | 32 | 2 |
| | | | | | Grapefruit | 155 | 13 |
| | | | | | Lemons | 158 | 44 |
| | | | | | Mandarins, clementines | 160 | 17 |
| | | | | | Oranges | 173 | 7 |
| | | | | | Peppers, sweet | 170 | 2 |
| | | | | | Pomelos | 34 | 1 |
| | | | | | Tomatoes | 91 | 7 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Melons | 51 | 1 |
| Quinalphos | F | 10951 | 10950 | 1 | Kiwi | 363 | 1 |
| | | | | | | | |
| Quintozene (sum) | DK | 5471 | 5459 | 12 | Carrots | 340 | 6 |
| | | | | | Courgettes | 20 | 1 |
| | | | | | Parsnips | 51 | 2 |
| | | | | | Potatoes | 424 | 2 |
| | | | | | Spinach | 81 | 1 |
| Quintozene (sum) | F | 11309 | 11300 | 9 | Carrots | 158 | 1 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Cucumbers | 218 | 1 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Parsnips | 10 | 2 |
| Quizalofop | F | 3975 | 3972 | 3 | Peppers, sweet | 385 | 3 |
| | | | | | Pears | 123 | 1 |
| | | | | | Spinach | 66 | 2 |
| Spiroxamine | F | 4516 | 4505 | 11 | Table grapes | 193 | 11 |
| Tebuconazole | DK | 3881 | 3877 | 4 | Brussels sprouts | 18 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|--------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Tebuconazole | F | 10951 | 10776 | 175 | Leek | 38 | 2 |
| | | | | | Parsnips | 51 | 1 |
| | | | | | Apples | 394 | 1 |
| | | | | | Apricots | 31 | 5 |
| | | | | | Beans with pods | 305 | 1 |
| | | | | | Carrots | 158 | 4 |
| | | | | | Cherries | 24 | 3 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Cucumbers | 218 | 1 |
| | | | | | Currants, black | 17 | 1 |
| | | | | | Currants, red | 28 | 4 |
| | | | | | Figs, fresh | 13 | 1 |
| | | | | | Grapefruit | 388 | 2 |
| | | | | | Guava | 9 | 1 |
| | | | | | Kiwi | 363 | 1 |
| | | | | | Leek | 68 | 5 |
| | | | | | Lemons | 380 | 1 |
| | | | | | Lettuce | 208 | 1 |
| | | | | | Limes | 52 | 1 |
| | | | | | Mandarins, clementines | 411 | 4 |
| | | | | | Melons | 311 | 2 |
| | | | | | Nectarines | 212 | 40 |
| | | | | | Papayas | 89 | 1 |
| | | | | | Peaches | 158 | 27 |
| | | | | | Peas with pods | 40 | 13 |
| | | | | | Peppers, sweet | 385 | 6 |
| | | | | | Persimmon | 108 | 1 |
| | | | | | Plums | 319 | 15 |
| | | | | | Rhubarbs | 8 | 1 |
| | | | | | Table grapes | 459 | 21 |
| Tomatoes | 249 | 9 | | | | | |
| Rolled oat | 86 | 1 | | | | | |
| Tebufenozide | F | 2886 | 2882 | 4 | Apples | 85 | 2 |
| | | | | | Peppers, sweet | 117 | 1 |
| | | | | | Wine, red | 91 | 1 |
| Tebufenpyrad | F | 7145 | 7114 | 31 | Apples | 213 | 3 |
| | | | | | Grapefruit | 292 | 1 |
| | | | | | Lemons | 290 | 2 |
| | | | | | Mandarins, clementines | 272 | 7 |
| | | | | | Nectarines | 158 | 1 |
| | | | | | Oranges | 326 | 3 |
| | | | | | Peaches | 111 | 1 |
| | | | | | Pears | 202 | 1 |
| | | | | | Strawberries | 178 | 3 |
| Tecnazene | F | 10951 | 10949 | 2 | Table grapes | 348 | 9 |
| | | | | | Ginger | 28 | 1 |
| | | | | | Kiwi | 363 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|--------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| Tetrachlorvinphos | F | 11072 | 11071 | 1 | Oranges | 491 | 1 |
| Tetraconazole | F | 1689 | 1687 | 2 | Peaches | 18 | 1 |
| | | | | | Peas with pods | 16 | 1 |
| Tetradifon | F | 10951 | 10938 | 13 | Basil | 11 | 2 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Kumquates | 14 | 2 |
| | | | | | Lemons | 380 | 2 |
| | | | | | Mandarins, clementines | 411 | 1 |
| | | | | | Oranges | 491 | 2 |
| | | | | | Passion fruits | 38 | 1 |
| | | | | | Strawberries | 266 | 1 |
| Thiabendazole | DK | 3494 | 3492 | 2 | Cucumbers | 198 | 1 |
| | | | | | Potatoes | 424 | 1 |
| Thiabendazole | F | 10483 | 9449 | 1034 | Apples | 392 | 70 |
| | | | | | Avocados | 40 | 1 |
| | | | | | Bananas | 425 | 184 |
| | | | | | Beans with pods | 305 | 2 |
| | | | | | Grapefruit | 388 | 255 |
| | | | | | Kiwi | 362 | 8 |
| | | | | | Lemons | 381 | 91 |
| | | | | | Lettuce | 206 | 1 |
| | | | | | Limes | 52 | 20 |
| | | | | | Mandarins, clementines | 411 | 133 |
| | | | | | Mango | 58 | 18 |
| | | | | | Melons | 308 | 4 |
| | | | | | Mineola | 3 | 2 |
| | | | | | Nectarines | 210 | 2 |
| | | | | | Oranges | 491 | 137 |
| | | | | | Papayas | 89 | 54 |
| | | | | | Peaches | 158 | 2 |
| | | | | | Pears | 299 | 27 |
| | | | | | Peppers, sweet | 382 | 1 |
| | | | | | Pitaya | 19 | 1 |
| | | | | | Plums | 317 | 2 |
| | | | | | Pomelos | 61 | 15 |
| | | | | | Rambutan | 47 | 1 |
| | | | | | Raspberries | 101 | 1 |
| | | | | | Tamarillo | 20 | 1 |
| | | | | | Watermelon | 39 | 1 |
| Thiophanate-methyl | F | 8407 | 8342 | 65 | Apples | 277 | 7 |
| | | | | | Beans with pods | 285 | 2 |
| | | | | | Cucumbers | 177 | 1 |
| | | | | | Currants, red | 17 | 1 |
| | | | | | Gooseberries | 6 | 1 |
| | | | | | Head cabbage, spring | 13 | 1 |
| | | | | | Lemons | 348 | 1 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------|---|---------------------------------|---|---|------------------------|---------------------------------|---|
| | | | | | Lettuce | 184 | 1 |
| | | | | | Mandarins, clementines | 356 | 4 |
| | | | | | Melons | 273 | 2 |
| | | | | | Papayas | 87 | 6 |
| | | | | | Passion fruits | 23 | 2 |
| | | | | | Peaches | 136 | 7 |
| | | | | | Pears | 244 | 10 |
| | | | | | Quince | 2 | 1 |
| | | | | | Shallots | 11 | 1 |
| | | | | | Spinach | 149 | 1 |
| | | | | | Strawberries | 225 | 5 |
| | | | | | Table grapes | 423 | 1 |
| | | | | | Teas | 25 | 1 |
| | | | | | Tomatoes | 180 | 2 |
| | | | | | Wine, red | 265 | 6 |
| | | | | | Wine, white | 7 | 1 |
| Tolclofos-methyl | F | 11073 | 11064 | 9 | Carrots | 158 | 4 |
| | | | | | Lettuce | 208 | 5 |
| Tolyfluanid (sum) | DK | 3881 | 3823 | 58 | Apples | 213 | 17 |
| | | | | | Currants, black | 11 | 7 |
| | | | | | Currants, red | 9 | 3 |
| | | | | | Pears | 165 | 7 |
| | | | | | Raspberries | 8 | 3 |
| | | | | | Strawberries | 163 | 19 |
| | | | | | Tomatoes | 212 | 2 |
| Tolyfluanid (sum) | F | 10951 | 10837 | 114 | Apples | 394 | 27 |
| | | | | | Cucumbers | 218 | 1 |
| | | | | | Currants, black | 17 | 1 |
| | | | | | Currants, red | 28 | 14 |
| | | | | | Gooseberries | 7 | 1 |
| | | | | | Pears | 301 | 52 |
| | | | | | Raspberries | 101 | 11 |
| | | | | | Strawberries | 266 | 4 |
| | | | | | Tomatoes | 249 | 3 |
| Triadimefon (sum) | F | 11072 | 10952 | 120 | Aubergines | 87 | 2 |
| | | | | | Beans with pods | 305 | 1 |
| | | | | | Carambola | 103 | 13 |
| | | | | | Chilli peppers | 42 | 1 |
| | | | | | Coriander, leaves | 9 | 1 |
| | | | | | Courgettes | 63 | 1 |
| | | | | | Cucumbers | 218 | 2 |
| | | | | | Grapefruit | 388 | 1 |
| | | | | | Melons | 311 | 7 |
| | | | | | Peppers, sweet | 385 | 14 |
| | | | | | Persimmon | 109 | 1 |
| | | | | | Pineapples | 55 | 37 |
| | | | | | Pomelos | 61 | 2 |

| Pesticide | Origin of samples (DK=Domestic; F=Foreign) | Number of samples ana- lysed | No. of samples without detected pesticides | Number of samples with detected pesticides | Commodity | Number of samples ana- lysed | Number of samples with detected residues |
|-------------------|---|---------------------------------|---|---|--------------------|---------------------------------|---|
| Triallate | F | 3975 | 3973 | 2 | Spinach | 163 | 1 |
| | | | | | Strawberries | 266 | 12 |
| Triazophos | F | 11072 | 11066 | 6 | Table grapes | 458 | 13 |
| | | | | | Tomatoes | 249 | 9 |
| Trichlorfon | F | 9948 | 9945 | 3 | Gojiberries, dried | 1 | 1 |
| | | | | | Wine, red | 266 | 1 |
| Trichloronat | DK | 3881 | 3880 | 1 | Grapefruit | 155 | 1 |
| | | | | | Plums | 131 | 1 |
| Trifloxystrobin | F | 10951 | 10864 | 87 | Aubergines | 87 | 1 |
| | | | | | Pomelos | 61 | 5 |
| Triflumuron | F | 3975 | 3954 | 21 | Figs, fresh | 13 | 1 |
| | | | | | Persimmon | 109 | 1 |
| Vinclozolin (sum) | DK | 5441 | 5440 | 1 | Tomatoes | 249 | 1 |
| | | | | | Chinese radish | 1 | 1 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Apples | 394 | 13 |
| | | | | | Courgettes | 63 | 1 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Cucumbers | 218 | 1 |
| | | | | | Currants, red | 28 | 3 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Grapefruit | 388 | 4 |
| | | | | | Nectarines | 212 | 2 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Oranges | 491 | 5 |
| | | | | | Passion fruits | 38 | 1 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Peaches | 158 | 2 |
| | | | | | Pears | 301 | 7 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Peppers, sweet | 385 | 3 |
| | | | | | Strawberries | 266 | 4 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Table grapes | 459 | 41 |
| | | | | | Apples | 124 | 7 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Nectarines | 90 | 8 |
| | | | | | Peaches | 64 | 5 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Pears | 123 | 1 |
| | | | | | Peas without pods | 10 | 1 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Beans with pods | 305 | 16 |
| | | | | | Carrots | 158 | 2 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Kiwi | 363 | 27 |
| | | | | | Lettuce | 208 | 8 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Nectarines | 212 | 2 |
| | | | | | Peaches | 158 | 1 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Peas without pods | 43 | 11 |
| | | | | | Raspberries | 101 | 12 |
| Vinclozolin (sum) | F | 11309 | 11229 | 80 | Table grapes | 459 | 1 |

7.4 Consumption used for exposure calculations.

| Consumer group | Adults | Male | Female | Children | Male High F&V ^{a)} | Female High F&V ^{a)} | Assumed domestic consump- tion |
|-----------------------------------|----------|----------|----------|----------|--------------------------------|----------------------------------|---|
| Age (years) | 15-75 | 15-75 | 15-75 | 4-6 | 15-75 | 15-75 | |
| Avg. weight (kg) | 75.1 | 83.5 | 68.2 | 21.8 | 84.4 | 69 | |
| No. of individuals in group | 1599 | 721 | 878 | 106 | 118 | 258 | |
| Average consumption (g/kg bw/day) | | | | | | | |
| Apples | 0.957 | 0.811 | 1.08 | 2.54 | 2.0 | 1.9 | 35% |
| Apricot, dried | 0.00401 | 0.00224 | 0.00546 | 0.016 | 0.00575 | 0.00945 | 0% |
| Apricots | 0.000055 | 0.0 | 0.0001 | 0.0 | 0.0 | 0.00034 | 0% |
| Asparagus | 0.0138 | 0.012 | 0.0153 | 0.00485 | 0.0155 | 0.0198 | 5% |
| Aubergines | 0.0108 | 0.00984 | 0.0115 | 0.00272 | 0.0275 | 0.0205 | 0% |
| Avocados | 0.0298 | 0.014 | 0.0428 | 0.0165 | 0.0353 | 0.064 | 0% |
| Bananas | 0.454 | 0.347 | 0.541 | 1.27 | 0.69 | 0.883 | 0% |
| Barley grit | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 7% |
| Barley kernels | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 100% |
| Beans with pods | 0.018 | 0.0157 | 0.0199 | 0.0317 | 0.0283 | 0.0314 | 2% |
| Beans, dried | 0.00594 | 0.00685 | 0.00518 | 0.0141 | 0.00849 | 0.00835 | 0% |
| Berries, dried | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Berries, others | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 21% |
| Blueberries | 0.000057 | 0.000126 | 0.0 | 0.0 | 0.0 | 0.0 | 0% |
| Broccoli | 0.0344 | 0.0268 | 0.0407 | 0.0465 | 0.0727 | 0.0698 | 25% |
| Brussels sprouts | 0.0143 | 0.0113 | 0.0167 | 0.0194 | 0.0303 | 0.029 | 55% |
| Carrots | 0.498 | 0.349 | 0.62 | 1.48 | 0.558 | 1.04 | 68% |
| Cauliflower | 0.0374 | 0.0332 | 0.0409 | 0.0307 | 0.0588 | 0.0589 | 40% |
| Celeriac | 0.0138 | 0.013 | 0.0145 | 0.0337 | 0.0147 | 0.019 | 63% |
| Celery | 0.00681 | 0.00471 | 0.00854 | 0.00368 | 0.015 | 0.0179 | 34% |
| Cherries | 0.000489 | 0.000334 | 0.000616 | 0.00119 | 0.0 | 0.00102 | 11% |
| Chervil | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Chick pea | 0.000187 | 0.000077 | 0.000278 | 0.0 | 0.000239 | 0.000578 | 0% |
| Chilli peppers | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Chives | 0.00092 | 0.000658 | 0.00114 | 0.000733 | 0.00105 | 0.0014 | 0% |
| Coriander, leaves | 0.000013 | 0.0 | 0.000024 | 0.0 | 0.0 | 0.000025 | 0% |
| Corn, dried | 0.0053 | 0.00346 | 0.00681 | 0.043 | 0.00207 | 0.00873 | 2% |
| Cornflour | 0.000905 | 0.000836 | 0.000961 | 0.00605 | 0.000752 | 0.000572 | 0% |
| Courgettes | 0.0215 | 0.02 | 0.0227 | 0.0052 | 0.0647 | 0.0419 | 24% |
| Cucumbers | 0.222 | 0.162 | 0.271 | 1.35 | 0.324 | 0.406 | 48% |
| Currants | 0.000054 | 0.0 | 0.000098 | 0.0 | 0.0 | 0.000166 | 31% |
| Dill | 0.000339 | 0.000187 | 0.000463 | 0.000147 | 0.000621 | 0.000625 | 25% |
| Fennel | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Figs, fresh | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Fruit (other), dried | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Fruit, exotic | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Fruit, mixed | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Garlics | 0.0012 | 0.00104 | 0.00134 | 0.000271 | 0.00176 | 0.00169 | 0% |
| Gherkin | 0.0212 | 0.0245 | 0.0184 | 0.0295 | 0.0242 | 0.0188 | 100% |
| Gooseberries | 0.000076 | 0.000169 | 0.0 | 0.0 | 0.0 | 0.0 | 36% |
| Grapefruit | 0.0189 | 0.0122 | 0.0245 | 0.0254 | 0.0327 | 0.047 | 0% |
| Head cabbage | 0.0649 | 0.0505 | 0.0768 | 0.0745 | 0.073 | 0.118 | 69% |
| Head cabbage, red | 0.0188 | 0.0242 | 0.0144 | 0.0233 | 0.0163 | 0.0172 | 82% |
| Herbs | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 3% |

| Consumer group | Adults | Male | Female | Children | Male | Female | Assumed domestic consumption |
|-----------------------------------|----------|----------|----------|----------|------------------------|------------------------|------------------------------|
| | | | | | High F&V ^{a)} | High F&V ^{a)} | |
| Age (years) | 15-75 | 15-75 | 15-75 | 4-6 | 15-75 | 15-75 | |
| Avg. weight (kg) | 75.1 | 83.5 | 68.2 | 21.8 | 84.4 | 69 | |
| No. of individuals in group | 1599 | 721 | 878 | 106 | 118 | 258 | |
| Average consumption (g/kg bw/day) | | | | | | | |
| Kale | 0.0048 | 0.00466 | 0.00492 | 0.00843 | 0.00311 | 0.00806 | 82% |
| Kiwi | 0.0426 | 0.0122 | 0.0676 | 0.129 | 0.03 | 0.138 | 0% |
| Leek | 0.0269 | 0.0231 | 0.03 | 0.0318 | 0.0308 | 0.0421 | 36% |
| Lemons, lime | 0.00803 | 0.00515 | 0.0104 | 0.0128 | 0.00461 | 0.0151 | 0% |
| Lentils | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Lettuce | 0.0903 | 0.0746 | 0.103 | 0.066 | 0.111 | 0.126 | 54% |
| Lettuce, iceberg | 0.000016 | 0.000012 | 0.000019 | 0.000355 | 0.0 | 0.000028 | 29% |
| Mandarins, clementines | 0.108 | 0.0698 | 0.14 | 0.145 | 0.187 | 0.269 | 0% |
| Mango | 0.00574 | 0.00196 | 0.00884 | 0.00685 | 0.00298 | 0.0161 | 0% |
| Marmelade, orange | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Melons | 0.139 | 0.0652 | 0.2 | 0.773 | 0.166 | 0.423 | 0% |
| Mineola | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Mushroom | 0.0421 | 0.042 | 0.0421 | 0.0448 | 0.0531 | 0.0587 | 50% |
| Nectarines | 0.0668 | 0.0483 | 0.0821 | 0.188 | 0.131 | 0.142 | 0% |
| Oat kernels | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 33% |
| Okra (Lady's fingers) | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Onions | 0.149 | 0.156 | 0.144 | 0.21 | 0.184 | 0.176 | 79% |
| Oranges | 0.143 | 0.0924 | 0.185 | 0.192 | 0.248 | 0.356 | 0% |
| Oranges, juice | 0.647 | 0.575 | 0.706 | 1.79 | 0.992 | 0.892 | 0% |
| Papayas | 0.000941 | 0.000322 | 0.00145 | 0.00112 | 0.000489 | 0.00265 | 0% |
| Parsley | 0.000869 | 0.000572 | 0.00111 | 0.000648 | 0.00129 | 0.00147 | 38% |
| Parsley Root | 0.00287 | 0.00256 | 0.00312 | 0.00204 | 0.00933 | 0.00734 | 85% |
| Parsnips | 0.00568 | 0.00512 | 0.00614 | 0.00409 | 0.0187 | 0.0145 | 84% |
| Pasta | 0.077 | 0.0814 | 0.0734 | 0.291 | 0.105 | 0.0719 | 0% |
| Peaches | 0.069 | 0.0499 | 0.0846 | 0.192 | 0.134 | 0.146 | 0% |
| Pear, canned | 0.00196 | 0.00151 | 0.00233 | 0.00358 | 0.00283 | 0.0038 | 0% |
| Pears | 0.323 | 0.229 | 0.4 | 0.76 | 0.632 | 0.794 | 35% |
| Peas with pods | 0.000403 | 0.000164 | 0.000599 | 0.00587 | 0.0 | 0.00104 | 13% |
| Peas without pods | 0.119 | 0.106 | 0.131 | 0.135 | 0.185 | 0.191 | 19% |
| Peppers, sweet | 0.0995 | 0.074 | 0.12 | 0.248 | 0.122 | 0.19 | 1% |
| Persimmon | 0.00273 | 0.000933 | 0.0042 | 0.00326 | 0.00142 | 0.00767 | 0% |
| Pineapples | 0.0394 | 0.0203 | 0.0551 | 0.0624 | 0.044 | 0.0969 | 0% |
| Plums | 0.0416 | 0.0257 | 0.0546 | 0.0887 | 0.0803 | 0.0899 | 18% |
| Pomegranate | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Pomelos | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Potatoes | 1.27 | 1.49 | 1.09 | 2.27 | 1.51 | 1.04 | 84% |
| Radish | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 27% |
| Raisin | 0.0219 | 0.0135 | 0.0287 | 0.103 | 0.0283 | 0.0489 | 0% |
| Raspberries, blackberries | 0.0103 | 0.00507 | 0.0145 | 0.0205 | 0.0121 | 0.0315 | 6% |
| Rhubarbs | 0.00251 | 0.00232 | 0.00266 | 0.00582 | 0.00531 | 0.00427 | 11% |
| Rice | 0.0821 | 0.0829 | 0.0814 | 0.162 | 0.111 | 0.0927 | 0% |
| Rice, short grained | 0.00217 | 0.00163 | 0.00262 | 0.0263 | 0.00194 | 0.00288 | 0% |
| Rolled oat | 0.109 | 0.123 | 0.0977 | 0.34 | 0.137 | 0.104 | 39% |
| Ruccola | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 40% |
| Rye flour | 0.527 | 0.599 | 0.468 | 1.63 | 0.665 | 0.527 | 57% |
| Sesame seed | 0.000247 | 0.0 | 0.000449 | 0.000299 | 0.0 | 0.000384 | 0% |
| Spinach | 0.0114 | 0.0109 | 0.0118 | 0.0225 | 0.00804 | 0.0177 | 33% |
| Spring onions | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 9% |
| Strawberries | 0.0707 | 0.0398 | 0.096 | 0.137 | 0.0952 | 0.199 | 38% |

| Consumer group | Adults | Male | Female | Children | Male High F&V ^{a)} | Female High F&V ^{a)} | Assumed domestic consump- tion |
|-----------------------------------|----------|---------|----------|----------|--------------------------------|----------------------------------|---|
| Age (years) | 15-75 | 15-75 | 15-75 | 4-6 | 15-75 | 15-75 | |
| Avg. weight (kg) | 75.1 | 83.5 | 68.2 | 21.8 | 84.4 | 69 | |
| No. of individuals in group | 1599 | 721 | 878 | 106 | 118 | 258 | |
| Average consumption (g/kg bw/day) | | | | | | | |
| Sweet corn (small) | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Table grapes | 0.173 | 0.0995 | 0.234 | 0.372 | 0.269 | 0.401 | 0% |
| Teas | 0.022 | 0.014 | 0.0286 | 0.0025 | 0.024 | 0.0335 | 0% |
| Tomatoes | 0.518 | 0.464 | 0.563 | 0.971 | 0.74 | 0.756 | 46% |
| Vegetables, unspecified | 0.000133 | 0.00012 | 0.000147 | 0.000458 | 0.000118 | 0.000145 | 0% |
| Watermelon | 0.0392 | 0.0184 | 0.0563 | 0.218 | 0.0468 | 0.119 | 0% |
| Wheat bran | 0.00364 | 0.00369 | 0.00361 | 0.0101 | 0.00382 | 0.00321 | 13% |
| Wheat flour | 1.12 | 1.13 | 1.11 | 3.12 | 1.11 | 1.06 | 49% |
| Wine, red | 0.979 | 1.02 | 0.947 | 0.0 | 1.25 | 0.987 | 1% |
| Wine, white | 0.261 | 0.222 | 0.294 | 0.0 | 0.197 | 0.246 | 0% |

^{a)} Consumption over 550 g of fruits and vegetables (excluding potatoes)

7.5 ADIs for pesticides included in the risk assessment.

The information concerning ADIs are mainly from the EU Pesticide database (http://ec.europa.eu/sanco_pesticides/public/index.cfm?event=homepage&CFID=9015839&CFTOKEN=26271605&jsessionid=08049d004dc4b5e4b7b2TR)

| Pesticide | ADI mg/kg bw/day | Source | Pesticide | ADI mg/kg bw/day | Source |
|-------------------------|------------------------|--------------------|---------------------------------------|------------------------|----------------------------|
| 2,4-D (sum) | 0.05 | COM 2001 | Dichlorvos | 0.00008 | EFSA 2006, tentative |
| Acephate | 0.03 | JMPR 2005 | Dicloran | 0.005 | EFSA 2010 |
| Acetamiprid | 0.07 | COM 2004 | Dicofol (sum) | 0.002 | JMPR 1994 |
| Aclonifen | 0.07 | COM 2008 | Diethofencarb | 0.43 | EFSA 2010 |
| Aldicarb (sum) | 0.003 | JMPR 2001 | Difenoconazole | 0.01 | COM 2008 |
| Aldrin and Dieldrin | 0.0001 | JMPR 2004 | Diffufenican | 0.2 | COM 2008 |
| Atrazine | 0.02 | JMPR 2007 | Dimethoate | 0.001 | COM 2007 |
| Azinphos-methyl | 0.005 | Com 2006 | Dimethomorph | 0.05 | COM 2007 |
| Azoxystrobin | 0.2 | EFSA 2010 | Diniconazole | 0.02 | DAR, FR |
| Benalaxyl (sum) | 0.04 | EFSA 2007 | Diphenylamine | 0.075 | EFSA 2008 |
| Benfuracarb | 0.01 | COM 2009 | Dithiocarbamates | 0.05 | For mancozeb, COM 2005 |
| Bifenthrin | 0.015 | EFSA 2011 | Endosulfan (sum) | 0.006 | JMPR 2006 |
| Binapacryl | | No ADI (JMPR 1985) | Epoxiconazole | 0.008 | EFSA 2008 |
| Biphenyl | 0.125 | JMPR 1967 | Ethion | 0.002 | JMPR 1990 |
| Bitertanol | 0.003 | COM 2011 | Ethoxyquin | 0.005 | JMPR 2005 |
| Bromopropylate | 0.03 | JMPR 1993 | Famoxadone | 0.012 | COM 2002 |
| Bupirimate | 0.05 | COM 2011 | Fenarimol | 0.01 | COM 2006 |
| Buprofezin | 0.01 | COM 2010 | Fenazaquin | 0.005 | COM 2011 |
| Captafol | | No ADI (JMPR 1985) | Fenbuconazole | 0.006 | COM 2010 |
| Captan/Folpet (sum) | 0.1 | COM 2008 | Fenhexamid | 0.2 | COM 2001 |
| Carbaryl | 0.0075 | EFSA 2006 | Fenitrothion | 0.005 | EFSA 2006 |
| Carbendazim and benomyl | 0.02 | COM 2006 | Fenoxaprop-P-Ethyl | 0.01 | COM 2008 |
| Carbofuran (sum) | 0.00015 | EFSA 2009 | Fenpropathrin | 0.03 | JMPR 1993 |
| Carbosulfan | 0.005 | EFSA 2009 | Fenpropimorph | 0.003 | COM 2008 |
| Chlorfenvinphos | 0.0005 | JMPR 1994 | Fenson | | No information (COM 2002) |
| Chlormequat | 0.04 | EFSA 2008 | Fenthion (sum) | 0.007 | JMPR 2000 |
| Chlorothalonil | 0.015 | COM 2006 | Fenvalerat, esfenvalerat, RR- and SS- | 0.02 | COM 2000 |
| Chlorpropham | 0.05 | COM 2003 | Fenvalerat, esfenvalerat, RS- and SR- | 0.02 | JMPR 1986 |
| Chlorpropham (sum) | 0.05 | COM 2004 | Flucythrinate | 0.02 | JMPR 1985 |
| Chlorpyrifos | 0.01 | COM 2005 | Fludioxonil | 0.37 | COM 2007 |
| Chlorpyrifos-methyl | 0.01 | COM 2005 | Fluoxastrobin | 0.015 | COM 2008 |
| Chlorthal-dimethyl | 0.01 | DAR 2007, EL | Fluroxypyr (sum) | 0.8 | COM 2010 |
| Clofentezine | 0.02 | COM 2010 | Flusilazole | 0.002 | COM 2007 |
| Cyfluthrin (sum) | 0.003 | COM 2003 | Flutriafol | 0.01 | EFSA 2011 |
| Cyhalothrin, lambda- | 0.005 | COM 2001 | Fluvalinate, tau- | 0.005 | COM 2010 |
| Cypermethrin (sum) | 0.05 | COM 2005 | Glyphosate | 0.3 | COM 2001 |
| Cyprodinil | 0.03 | COM 2006 | HCH (sum) | | No information (JMPR 1973) |
| Cyromazine | 0.06 | COM 2009 | Hexachlorobenzene | | ADI withdrawn (JMPR 1978) |
| DDT (sum) | 0.01 | JMPR 2000 | Hexaconazole | 0.005 | JMPR 1990 |
| Deltamethrin | 0.01 | Com 2003 | Hexythiazox | 0.03 | COM 2011 |
| Diazinon | 0.0002 | EFSA 2006 | Imazalil | 0.025 | EFSA 2010 |
| Dichlofluanid | 0.3 | JMPR 1983 | Iprodione | 0.06 | COM 2002 |
| Dichlorprop | 0.06 | COM 2007 | Iprovalicarb | 0.015 | COM 2002 |

| Pesticide | ADI mg/kg bw/day | Source | Pesticide | ADI mg/kg bw/day | Source |
|-------------------------|------------------------|------------------------------|--------------------|------------------------|---|
| Kresoxim-methyl | 0.4 | COM 2011 | Propamocarb (sum) | 0.29 | COM 2007 |
| Lindane | 0.005 | COM 2000 | Propargite | 0.01 | JMPR 1991 EFSA: Data not sufficient to derive a MRL |
| Linuron | 0.003 | COM 2003 | Propiconazole | 0.04 | COM 2003 |
| Malathion (sum) | 0.03 | COM 2010 | Propoxur | 0.02 | JMPR 1989 |
| Mecoprop (sum) | 0.01 | COM 2003 | Propyzamide | 0.02 | COM 2003 |
| Mepiquat | 0.2 | COM 2008 | Prothiofos | | No information (COM 2002) |
| Metalaxyl (sum) | 0.08 | COM 2002 | Pymetrozine | 0.03 | COM 2001 |
| Methacrifos | 0.006 | JMPR 1990 | Pyraclostrobin | 0.03 | COM 2004 |
| Methamidophos | 0.001 | COM 2007 | Pyrazophos | 0.004 | JMPR 1992 |
| Methidathion | 0.001 | JMPR 1997 | Pyridaben | 0.01 | EFSA 2010 |
| Methiocarb (sum) | 0.013 | COM 2007 | Pyridate (sum) | 0.036 | COM 2001 |
| Methomyl and Thiodicarb | 0.0025 | COM 2009 | Pyrimethanil | 0.17 | COM 2006 |
| Metribuzin | 0.013 | COM 2007 | Pyriproxyfen | 0.1 | COM 2008 |
| Mevinphos | 0.0008 | JMPR 2000 | Quinalphos | | No information (COM 2002) |
| Monocrotophos | 0.0006 | JMPR 1995 | Quinoxifen | 0.2 | COM 2004 |
| Myclobutanil | 0.025 | EFSA 2010 | Quintozene (sum) | 0.01 | COM 2000 |
| Ofurace | | No information (COM 2002) | Quizalofop | | No information (COM 2002) |
| Omethoate | 0.0003 | COM 2007 | Spiroxamine | 0.025 | COM 1999 |
| Orthophenylphenol | 0.4 | EFSA 2008 | Tebuconazole | 0.03 | EFSA 2008 |
| Oxamyl | 0.001 | COM 2006 | Tebufenozide | 0.02 | COM 2010 |
| Oxydemeton-methyl (sum) | 0.0003 | COM 2006 | Tebufenpyrad | 0.01 | COM 2009 |
| Parathion-methyl (sum) | 0.003 | JMPR 2003 | Tecnazene | 0.02 | JMPR 1994 |
| Penconazole | 0.03 | COM 2009 | Tetrachlorvinphos | | No information (COM 2002) |
| Pendimethalin | 0.125 | COM 2003 | Tetraconazole | 0.004 | EFSA 2008 |
| Pentachlorobenzene | | No information | Tetradifon | 0.015 | DE 2001 |
| Pentachlorophenol | | No information | Thiabendazole | 0.1 | COM 2001 |
| Permethrin (sum) | 0.05 | JMPR 2002 | Thiophanate-methyl | 0.08 | COM 2005 |
| Phenthoate | 0.003 | JMPR 1984 | Tolclofos-methyl | 0.064 | COM 2006 |
| Phorate (sum) | 0.0007 | JMPR 2005 | Tolyfluanid (sum) | 0.1 | COM 2006 |
| Phosalone | 0.01 | COM 2006 | Triadimefon (sum) | 0.05 | JMPR 2004 |
| Phosmet (sum) | 0.003 | COM 2007 | Triallate | 0.025 | EFSA 2008 |
| Pirimicarb (sum) | 0.035 | COM 2006 | Triazophos | 0.001 | JMPR 2002 |
| Pirimiphos-methyl | 0.004 | EFSA 05 | Trichlorfon | 0.045 | DAR, ES EFSA: No agreed ADI available |
| Prochloraz (sum) | 0.01 | EFSA 2011 | Trifloxystrobin | 0.1 | COM 2003 |
| Procymidone | 0.0028 | DAR FR 2007 | Triflumuron | 0.014 | COM 2011 |
| Profenofos | 0.03 | JMPR 2007 | Vinclozolin (sum) | 0.005 | COM 2006 |

7.6 Reduction factors.

Reductions factors for the pesticides have been found from the literature in citrus fruits, melons and bananas. Reduction factors in citrus fruits from Rapport 7/98 Statens Livsmedelverk, Sweden is shown below.

Reduction factors in citrus fruits found in Sweden:

| Commodity/pesticide | Content of total residue in peel (%) |
|---------------------|--------------------------------------|
| Orange | |
| Azinfosmethyl | ≥90 |
| Bromopropylat | ≥90 |
| Dicofol | 97 |
| Dimethoate | ≥90 |
| Fenithrothion | ≥95 |
| Phosmet | ≥90 |
| Chlorfenvinphos | ≥90 |
| Chlorpyrifos | ≥90 |
| Quinalphos | ≥90 |
| Mecarbam | ≥95 |
| Methidathion | ≥95 |
| Parathion | ≥95 |
| Parathion-methyl | ≥95 |
| Tetradifon | ≥90 |
| Lemon | |
| Mecarbam | ≥95 |
| Grapefruit | |
| Ethion | ≥97 |
| Small citrus fruits | |
| Ethion | ≥95 |
| Chlorfenvinphos | ≥90 |
| Malathion | ≥97 |
| Methidathion | ≥95 |

Livsmedelverket has found that for thiabendazol about 15-25% of the pesticide is in the pulp from oranges.

The reduction factors found in the JMPR reports are shown below.

Reduction factors found in JMPR reports

| Pesticide | Commodity | Reduction | Source |
|---------------|-----------|--|-----------|
| Thiabendazole | Oranges | <3% in pulp; >97% in peel | JMPR 2000 |
| Imazalil | Melon | About 10% in pulp; about 90% in peel | JMPR 1994 |
| Phenyl-phenol | Oranges | 2-4% in pulp; 96-98 % in peel | JMPR 1999 |
| Benomyl | Oranges | From oranges to orange juice the reduction is 17-98% | JMPR 1998 |
| Procymidon | Kiwi | In pulp about 1%; in peel about 99% | JMPR 1998 |

Conclusion

Most results for reduction factors are for citrus fruits. As bananas and melons also have a thick peel it is estimated that the results for citrus fruits can be transferred to these two commodities. Therefore overall a reduction factor of 90% is used for, both citrus fruits, melons and bananas except for thiabendazole and pesticides from the benomyl group (carbendazim,

thiophanat-methyl and benomyl). For these substances a reduction of 75% is used, as it is the lowest reduction found and the worst-case situation.

Processing factors used in present report:

| Commodity | Processing factor |
|------------------------|---|
| Bananas | |
| Grapefruit | |
| Lemons, lime | |
| Mandarins, clementines | 75% for, thiabendazole, thiophanate-methyl, carbendazim and benomyl |
| Melons | |
| Mineola | 90% for all other pesticides |
| Oranges | |
| Pomelos | |
| Watermelon | |

7.7 Exposure and sum of Hazard Quotients (Hazard Index (HI)) for individual commodities (consumer group “Adults”).

Model 1: No correction for non-detects

Model 3: Non-detects assumed to be ½LOR; correction factor limited to 25.

Corrected for peeling unless stated otherwise.

| Commodity | Exposure (µg/kg bw/day) | | Exposure (µg/day) | | Hazard Index | |
|--------------------------------|-------------------------|--------------|-------------------|------------|--------------|--------------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Apples | 0.27 | 0.57 | 20 | 43 | 1.3% | 6.9% |
| Apricot, dried | 0.000015 | 0.000025 | 0.0012 | 0.0019 | 0.0% | 0.0001% |
| Apricots | 0.000015 | 0.000028 | 0.0011 | 0.0021 | 0.0001% | 0.0001% |
| Asparagus | 0.000064 | 0.00018 | 0.0048 | 0.013 | 0.0006% | 0.0013% |
| Aubergines | 0.00078 | 0.0027 | 0.059 | 0.21 | 0.0025% | 0.018% |
| Avocados | 0.0014 | 0.0021 | 0.1 | 0.16 | 0.011% | 0.017% |
| Bananas | 0.012 | 0.016 | 0.92 | 1.2 | 0.031% | 0.049% |
| <i>Bananas (not peeled)</i> | <i>0.087</i> | <i>0.11</i> | <i>6.5</i> | <i>8.6</i> | <i>0.27%</i> | <i>0.44%</i> |
| Barley grit | 0.0000097 | 0.000018 | 0.00073 | 0.0013 | 0.0% | 0.0% |
| Barley kernels | 0.000021 | 0.000029 | 0.0016 | 0.0022 | 0.0% | 0.0% |
| Beans with pods | 0.0013 | 0.0052 | 0.1 | 0.39 | 0.025% | 0.12% |
| Beans, dried | 0.0000081 | 0.000031 | 0.00061 | 0.0023 | 0.0001% | 0.0005% |
| Berries, dried | 0.000062 | 0.000083 | 0.0046 | 0.0062 | 0.0005% | 0.0009% |
| Berries, others | 0.0000024 | 0.0000091 | 0.00018 | 0.00068 | 0.0% | 0.0001% |
| Blueberries | 0.0000052 | 0.000011 | 0.00039 | 0.00084 | 0.0% | 0.0% |
| Broccoli | 0.00003 | 0.00033 | 0.0022 | 0.025 | 0.0002% | 0.0019% |
| Brussels sprouts | 0.000079 | 0.00023 | 0.006 | 0.017 | 0.0005% | 0.0017% |
| Carrots | 0.0055 | 0.037 | 0.41 | 2.7 | 0.15% | 1.0% |
| Cauliflower | 0.000012 | 0.00016 | 0.00093 | 0.012 | 0.0004% | 0.0057% |
| Celeriac | 0.00014 | 0.00033 | 0.01 | 0.025 | 0.0034% | 0.0072% |
| Celery | 0.0015 | 0.0021 | 0.11 | 0.16 | 0.01% | 0.016% |
| Cherries | 0.000028 | 0.000057 | 0.0021 | 0.0043 | 0.001% | 0.0042% |
| Chervil | 0.000018 | 0.00002 | 0.0013 | 0.0015 | 0.0002% | 0.0002% |
| Chick pea | 0.00029 | 0.00031 | 0.022 | 0.023 | 0.0001% | 0.0001% |
| Chilli peppers | 0.000065 | 0.0001 | 0.0049 | 0.0078 | 0.0017% | 0.0031% |
| Chives | 0.000028 | 0.00004 | 0.0021 | 0.003 | 0.0001% | 0.0002% |
| Coriander, leaves | 0.000034 | 0.000035 | 0.0025 | 0.0026 | 0.0003% | 0.0003% |
| Corn, dried | 0.00028 | 0.00038 | 0.021 | 0.029 | 0.004% | 0.0064% |
| Cornflour | 0.000065 | 0.000088 | 0.0049 | 0.0066 | 0.0005% | 0.0007% |
| Courgettes | 0.00034 | 0.0022 | 0.025 | 0.16 | 0.011% | 0.03% |
| Cucumbers | 0.024 | 0.053 | 1.8 | 4 | 0.16% | 0.69% |
| Currants | 0.00004 | 0.000052 | 0.003 | 0.0039 | 0.0001% | 0.0002% |
| Dill | 0.000044 | 0.000045 | 0.0033 | 0.0034 | 0.0004% | 0.0005% |
| Fennel | 0.0000015 | 0.0000022 | 0.00011 | 0.00016 | 0.0% | 0.0% |
| Figs, fresh | 0.000049 | 0.00006 | 0.0037 | 0.0045 | 0.0003% | 0.0004% |
| Fruit (other), dried | 0.00000027 | 0.0000023 | 0.00002 | 0.00018 | 0.0% | 0.0% |
| Fruit, exotic | 0.000021 | 0.000052 | 0.0015 | 0.0039 | 0.0001% | 0.0009% |
| Fruit, mixed | 0.0000026 | 0.0000049 | 0.0002 | 0.00037 | 0.0% | 0.0% |
| Garlics | 0.00000026 | 0.0000036 | 0.00002 | 0.00027 | 0.0% | 0.0% |
| Gherkin | 0.0028 | 0.0035 | 0.21 | 0.26 | 0.0056% | 0.0071% |
| Gooseberries | 0.0000069 | 0.0000096 | 0.00052 | 0.00072 | 0.0% | 0.0% |
| Grapefruit | 0.0046 | 0.0051 | 0.35 | 0.38 | 0.016% | 0.034% |
| <i>Grapefruit (not peeled)</i> | <i>0.032</i> | <i>0.037</i> | <i>2.4</i> | <i>2.7</i> | <i>0.14%</i> | <i>0.33%</i> |
| Head cabbage | 0.000023 | 0.00036 | 0.0017 | 0.027 | 0.0002% | 0.0051% |
| Head cabbage, red | 0.000017 | 0.000063 | 0.0013 | 0.0048 | 0.0% | 0.0001% |
| Herbs | 0.000014 | 0.000036 | 0.001 | 0.0027 | 0.0001% | 0.0007% |
| Kale | 0.00081 | 0.00087 | 0.061 | 0.065 | 0.0011% | 0.0013% |

| Commodity | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | | Exposure ($\mu\text{g}/\text{day}$) | | Hazard Index | |
|--|--|---------------|---------------------------------------|--------------|----------------|----------------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Kiwi | 0.039 | 0.044 | 2.9 | 3.3 | 0.12% | 0.27% |
| Leek | 0.0023 | 0.0034 | 0.17 | 0.25 | 0.0051% | 0.0079% |
| Lemons, lime | 0.0014 | 0.0016 | 0.11 | 0.12 | 0.0071% | 0.012% |
| <i>Lemons, lime (not peeled)</i> | <i>0.012</i> | <i>0.014</i> | <i>0.93</i> | <i>1.1</i> | <i>0.069%</i> | <i>0.12%</i> |
| Lentils | 0.00021 | 0.00022 | 0.016 | 0.016 | 0.0001% | 0.0001% |
| Lettuce | 0.052 | 0.063 | 3.9 | 4.8 | 0.18% | 0.26% |
| Lettuce, iceberg | 0.00000038 | 0.0000019 | 0.000029 | 0.00014 | 0.0% | 0.0% |
| Mandarins, clementines | 0.026 | 0.029 | 2 | 2.2 | 0.11% | 0.21% |
| <i>Mandarins, clementines (not peeled)</i> | <i>0.21</i> | <i>0.24</i> | <i>16</i> | <i>18</i> | <i>1.1%</i> | <i>2.0%</i> |
| Mango | 0.002 | 0.0025 | 0.15 | 0.19 | 0.012% | 0.014% |
| Marmelade, orange | 0.0000046 | 0.000014 | 0.00035 | 0.0011 | 0.0% | 0.0002% |
| Melons | 0.0015 | 0.0051 | 0.11 | 0.38 | 0.011% | 0.08% |
| <i>Melons (not peeled)</i> | <i>0.014</i> | <i>0.046</i> | <i>1</i> | <i>3.4</i> | <i>0.1%</i> | <i>0.79%</i> |
| Mineola | 0.000069 | 0.00007 | 0.0052 | 0.0052 | 0.0001% | 0.0001% |
| Mushroom | 0.0022 | 0.0033 | 0.17 | 0.25 | 0.013% | 0.022% |
| Nectarines | 0.011 | 0.03 | 0.84 | 2.3 | 0.039% | 0.3% |
| Oat kernels | 0.000056 | 0.000059 | 0.0042 | 0.0044 | 0.0001% | 0.0001% |
| Okra (Lady's fingers) | 0.000027 | 0.000038 | 0.0021 | 0.0029 | 0.0002% | 0.0002% |
| Onions | 0.00018 | 0.0019 | 0.014 | 0.14 | 0.0015% | 0.016% |
| Oranges | 0.025 | 0.03 | 1.9 | 2.2 | 0.098% | 0.28% |
| <i>Oranges (not peeled)</i> | <i>0.2</i> | <i>0.24</i> | <i>15</i> | <i>18</i> | <i>0.92%</i> | <i>2.7%</i> |
| Oranges, juice | 0.001 | 0.016 | 0.078 | 1.2 | 0.0038% | 0.068% |
| Papayas | 0.00043 | 0.00051 | 0.032 | 0.038 | 0.0015% | 0.0026% |
| Parsley | 0.0001 | 0.00011 | 0.0076 | 0.008 | 0.0033% | 0.0034% |
| Parsley Root | 0.000035 | 0.0001 | 0.0026 | 0.0076 | 0.0007% | 0.0019% |
| Parsnips | 0.000047 | 0.00023 | 0.0035 | 0.017 | 0.0007% | 0.0035% |
| Pasta | 0.00034 | 0.0025 | 0.025 | 0.19 | 0.0068% | 0.05% |
| Peaches | 0.013 | 0.032 | 0.95 | 2.4 | 0.055% | 0.28% |
| Pear, canned | 0.000086 | 0.00012 | 0.0065 | 0.0088 | 0.0002% | 0.0002% |
| Pears | 0.073 | 0.14 | 5.5 | 11 | 0.29% | 1.0% |
| Peas with pods | 0.00014 | 0.00019 | 0.01 | 0.015 | 0.0022% | 0.0034% |
| Peas without pods | 0.0024 | 0.0084 | 0.18 | 0.63 | 0.04% | 0.24% |
| Peppers, sweet | 0.0056 | 0.033 | 0.42 | 2.5 | 0.067% | 0.31% |
| Persimmon | 0.000056 | 0.00034 | 0.0042 | 0.026 | 0.0007% | 0.0038% |
| Pineapples | 0.015 | 0.02 | 1.1 | 1.5 | 0.052% | 0.24% |
| Plums | 0.0088 | 0.017 | 0.66 | 1.3 | 0.024% | 0.12% |
| Pomegranate | 0.000013 | 0.000024 | 0.001 | 0.0018 | 0.0001% | 0.0001% |
| Pomelos | 0.000028 | 0.000031 | 0.0021 | 0.0023 | 0.0001% | 0.0002% |
| <i>Pomelos (not peeled)</i> | <i>0.00018</i> | <i>0.0002</i> | <i>0.013</i> | <i>0.015</i> | <i>0.0007%</i> | <i>0.0015%</i> |
| Potatoes | 0.033 | 0.055 | 2.5 | 4.1 | 0.071% | 0.17% |
| Radish | 0.0000036 | 0.000006 | 0.00027 | 0.00045 | 0.0% | 0.0% |
| Raisin | 0.00025 | 0.00032 | 0.019 | 0.024 | 0.0012% | 0.0016% |
| Raspberries, blackberries | 0.0017 | 0.004 | 0.13 | 0.3 | 0.008% | 0.02% |
| Rhubarbs | 0.000015 | 0.000034 | 0.0012 | 0.0025 | 0.0% | 0.0001% |
| Rice | 0.00073 | 0.005 | 0.055 | 0.38 | 0.0089% | 0.067% |
| Rice, short grained | 0.0000092 | 0.000046 | 0.00069 | 0.0034 | 0.0002% | 0.0011% |
| Rolled oat | 0.0077 | 0.016 | 0.58 | 1.2 | 0.02% | 0.056% |
| Ruccola | 0.00017 | 0.00017 | 0.013 | 0.013 | 0.0003% | 0.0003% |
| Rye flour | 0.016 | 0.032 | 1.2 | 2.4 | 0.24% | 0.44% |
| Sesame seed | 0.0000065 | 0.000014 | 0.00049 | 0.0011 | 0.0002% | 0.0004% |
| Spinach | 0.00062 | 0.0023 | 0.046 | 0.17 | 0.0069% | 0.026% |
| Spring onions | 0.0000089 | 0.000019 | 0.00067 | 0.0014 | 0.0002% | 0.0005% |
| Strawberries | 0.01 | 0.023 | 0.76 | 1.8 | 0.033% | 0.11% |
| Sweet corn (small) | 0.00000017 | 0.0000013 | 0.000013 | 0.000097 | 0.0% | 0.0% |
| Table grapes | 0.056 | 0.11 | 4.2 | 8.4 | 0.21% | 0.74% |

| Commodity | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | | Exposure ($\mu\text{g}/\text{day}$) | | Hazard Index | |
|--------------------------------|--|---------------|---------------------------------------|-------------|---------------|---------------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Teas | 0.0021 | 0.0047 | 0.15 | 0.35 | 0.051% | 0.12% |
| Tomatoes | 0.045 | 0.15 | 3.4 | 11 | 0.29% | 1.2% |
| Vegetables, unspecified | 0.000026 | 0.00003 | 0.002 | 0.0022 | 0.0016% | 0.0018% |
| Watermelon | 0.00014 | 0.0006 | 0.01 | 0.045 | 0.0027% | 0.0078% |
| <i>Watermelon (not peeled)</i> | <i>0.0011</i> | <i>0.0051</i> | <i>0.086</i> | <i>0.38</i> | <i>0.027%</i> | <i>0.076%</i> |
| Wheat bran | 0.00073 | 0.00076 | 0.055 | 0.057 | 0.0026% | 0.0029% |
| Wheat flour | 0.061 | 0.22 | 4.6 | 16 | 0.3% | 1.3% |
| Wine, red | 0.037 | 0.11 | 2.8 | 8.4 | 0.13% | 0.48% |
| Wine, white | 0.02 | 0.022 | 1.5 | 1.6 | 0.045% | 0.05% |
| Total | 0.90 | 1.94 | 68 | 146 | 4% | 18% |

7.8 Exposure and sum of Hazard Quotients (Hazard Index (HI)) for individual pesticides (consumer group “Children”).

Model 1: No correction for non-detects

Model 3: Non-detects assumed to be ½LOR; correction factor limited to 25.

Corrected for peeling unless stated otherwise.

| Commodity | Exposure (µg/kg bw/day) | | Exposure (µg/day) | | Hazard Index | |
|--------------------------------|-------------------------|--------------|-------------------|------------|--------------|--------------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Apples | 0.72 | 1.5 | 16 | 33 | 3.6% | 18.0% |
| Apricot, dried | 0.000062 | 0.0001 | 0.0013 | 0.0022 | 0.00012% | 0.0002% |
| Apricots | | | | | | |
| Asparagus | 0.000023 | 0.000062 | 0.00049 | 0.0013 | 0.00022% | 0.00044% |
| Aubergines | 0.0002 | 0.00069 | 0.0043 | 0.015 | 0.00062% | 0.0045% |
| Avocados | 0.00075 | 0.0011 | 0.016 | 0.025 | 0.0063% | 0.0093% |
| Bananas | 0.034 | 0.043 | 0.75 | 0.95 | 0.086% | 0.14% |
| <i>Bananas (not peeled)</i> | <i>0.24</i> | <i>0.32</i> | <i>5.3</i> | <i>7</i> | <i>0.76%</i> | <i>1.2%</i> |
| Barley grit | 0.000033 | 0.00006 | 0.00073 | 0.0013 | 0.000011% | 0.00002% |
| Barley kernels | 0.000072 | 0.0001 | 0.0016 | 0.0022 | 0.000024% | 0.000034% |
| Beans with pods | 0.0023 | 0.0093 | 0.051 | 0.2 | 0.045% | 0.21% |
| Beans, dried | 0.000019 | 0.000073 | 0.00042 | 0.0016 | 0.00032% | 0.0012% |
| Berries, dried | 0.00021 | 0.00029 | 0.0046 | 0.0062 | 0.0017% | 0.0032% |
| Berries, others | 0.0000084 | 0.000031 | 0.00018 | 0.00068 | 0.000067% | 0.00026% |
| Blueberries | | | | | | |
| Broccoli | 0.00004 | 0.00044 | 0.00088 | 0.0096 | 0.00024% | 0.0026% |
| Brussels sprouts | 0.00011 | 0.00031 | 0.0024 | 0.0067 | 0.00063% | 0.0023% |
| Carrots | 0.016 | 0.11 | 0.35 | 2.4 | 0.46% | 3.0% |
| Cauliflower | 0.00001 | 0.00013 | 0.00022 | 0.0028 | 0.00036% | 0.0047% |
| Celeriac | 0.00033 | 0.00081 | 0.0073 | 0.018 | 0.0083% | 0.018% |
| Celery | 0.00079 | 0.0011 | 0.017 | 0.025 | 0.0056% | 0.0087% |
| Cherries | 0.000067 | 0.00014 | 0.0015 | 0.003 | 0.0023% | 0.01% |
| Chervil | 0.000061 | 0.00007 | 0.0013 | 0.0015 | 0.00057% | 0.00062% |
| Chick pea | | | | | | |
| Chilli peppers | 0.00022 | 0.00036 | 0.0049 | 0.0078 | 0.0059% | 0.011% |
| Chives | 0.000022 | 0.000032 | 0.00049 | 0.0007 | 0.00011% | 0.00018% |
| Coriander, leaves | | | | | | |
| Corn, dried | 0.0023 | 0.0031 | 0.049 | 0.068 | 0.032% | 0.052% |
| Cornflour | 0.00043 | 0.00058 | 0.0094 | 0.013 | 0.0035% | 0.0048% |
| Courgettes | 0.000082 | 0.00053 | 0.0018 | 0.012 | 0.0027% | 0.0072% |
| Cucumbers | 0.15 | 0.32 | 3.2 | 7 | 0.96% | 4.2% |
| Currants | | | | | | |
| Dill | 0.000019 | 0.00002 | 0.00042 | 0.00043 | 0.00019% | 0.0002% |
| Fennel | 0.0000051 | 0.0000074 | 0.00011 | 0.00016 | 0.000034% | 0.000047% |
| Figs, fresh | 0.00017 | 0.00021 | 0.0037 | 0.0045 | 0.00086% | 0.0012% |
| Fruit (other), dried | 0.00000092 | 0.000008 | 0.00002 | 0.00018 | 0.000005% | 0.00004% |
| Fruit, exotic | 0.000071 | 0.00018 | 0.0015 | 0.0039 | 0.00043% | 0.0031% |
| Fruit, mixed | 0.0000091 | 0.000017 | 0.0002 | 0.00037 | 0.000035% | 0.000067% |
| Garlics | 0.00000006 | 0.00000081 | 0.0000013 | 0.000018 | 0.0000004% | 0.000005% |
| Gherkin | 0.0039 | 0.0049 | 0.086 | 0.11 | 0.0079% | 0.0098% |
| Gooseberries | | | | | | |
| Grapefruit | 0.0062 | 0.0068 | 0.14 | 0.15 | 0.021% | 0.046% |
| <i>Grapefruit (not peeled)</i> | <i>0.043</i> | <i>0.049</i> | <i>0.94</i> | <i>1.1</i> | <i>0.19%</i> | <i>0.44%</i> |
| Head cabbage | 0.000026 | 0.00042 | 0.00058 | 0.0091 | 0.00025% | 0.0059% |
| Head cabbage, red | 0.000021 | 0.000078 | 0.00047 | 0.0017 | 0.000036% | 0.00013% |
| Herbs | 0.000047 | 0.00012 | 0.001 | 0.0027 | 0.00046% | 0.0025% |
| Kale | 0.0014 | 0.0015 | 0.031 | 0.033 | 0.002% | 0.0023% |

| Commodity | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | | Exposure ($\mu\text{g}/\text{day}$) | | Hazard Index | |
|--|--|---------------|---------------------------------------|-------------|----------------|----------------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Kiwi | 0.12 | 0.13 | 2.6 | 2.9 | 0.38% | 0.82% |
| Leek | 0.0027 | 0.004 | 0.059 | 0.087 | 0.006% | 0.0094% |
| Lemons, lime | 0.0022 | 0.0025 | 0.049 | 0.055 | 0.011% | 0.019% |
| <i>Lemons, lime (not peeled)</i> | <i>0.02</i> | <i>0.022</i> | <i>0.43</i> | <i>0.49</i> | <i>0.11%</i> | <i>0.18%</i> |
| Lentils | 0.00072 | 0.00075 | 0.016 | 0.016 | 0.00025% | 0.00027% |
| Lettuce | 0.038 | 0.046 | 0.83 | 1 | 0.13% | 0.19% |
| Lettuce, iceberg | 0.0000085 | 0.000043 | 0.00018 | 0.00093 | 0.00016% | 0.00083% |
| Mandarins, clementines | 0.035 | 0.04 | 0.77 | 0.86 | 0.15% | 0.28% |
| <i>Mandarins, clementines (not peeled)</i> | <i>0.29</i> | <i>0.33</i> | <i>6.3</i> | <i>7.2</i> | <i>1.5%</i> | <i>2.7%</i> |
| Mango | 0.0024 | 0.003 | 0.052 | 0.064 | 0.014% | 0.016% |
| Marmelade, orange | 0.000016 | 0.000049 | 0.00035 | 0.0011 | 0.00011% | 0.00063% |
| Melons | 0.0083 | 0.029 | 0.18 | 0.62 | 0.06% | 0.45% |
| <i>Melons (not peeled)</i> | <i>0.076</i> | <i>0.26</i> | <i>1.7</i> | <i>5.6</i> | <i>0.58%</i> | <i>4.4%</i> |
| Mineola | 0.00024 | 0.00024 | 0.0052 | 0.0052 | 0.00038% | 0.00038% |
| Mushroom | 0.0024 | 0.0035 | 0.052 | 0.077 | 0.014% | 0.023% |
| Nectarines | 0.032 | 0.086 | 0.69 | 1.9 | 0.11% | 0.84% |
| Oat kernels | 0.00019 | 0.0002 | 0.0042 | 0.0044 | 0.00043% | 0.00044% |
| Okra (Lady's fingers) | 0.000094 | 0.00013 | 0.0021 | 0.0029 | 0.00058% | 0.00084% |
| Onions | 0.00026 | 0.0027 | 0.0057 | 0.059 | 0.0021% | 0.022% |
| Oranges | 0.034 | 0.04 | 0.74 | 0.87 | 0.13% | 0.37% |
| <i>Oranges (not peeled)</i> | <i>0.27</i> | <i>0.33</i> | <i>6</i> | <i>7.2</i> | <i>1.2%</i> | <i>3.6%</i> |
| Oranges, juice | 0.0029 | 0.045 | 0.062 | 0.98 | 0.011% | 0.19% |
| Papayas | 0.00051 | 0.00061 | 0.011 | 0.013 | 0.0018% | 0.0032% |
| Parsley | 0.000075 | 0.000079 | 0.0016 | 0.0017 | 0.0024% | 0.0025% |
| Parsley Root | 0.000025 | 0.000072 | 0.00054 | 0.0016 | 0.00051% | 0.0013% |
| Parsnips | 0.000034 | 0.00017 | 0.00074 | 0.0036 | 0.00052% | 0.0025% |
| Pasta | 0.0013 | 0.0094 | 0.028 | 0.2 | 0.026% | 0.19% |
| Peaches | 0.035 | 0.089 | 0.77 | 1.9 | 0.15% | 0.79% |
| Pear, canned | 0.00016 | 0.00022 | 0.0034 | 0.0047 | 0.00039% | 0.00042% |
| Pears | 0.17 | 0.33 | 3.7 | 7.3 | 0.68% | 2.5% |
| Peas with pods | 0.002 | 0.0028 | 0.043 | 0.061 | 0.032% | 0.049% |
| Peas without pods | 0.0027 | 0.0095 | 0.059 | 0.21 | 0.045% | 0.27% |
| Peppers, sweet | 0.014 | 0.082 | 0.3 | 1.8 | 0.17% | 0.76% |
| Persimmon | 0.000067 | 0.00041 | 0.0015 | 0.0089 | 0.00083% | 0.0045% |
| Pineapples | 0.023 | 0.031 | 0.51 | 0.69 | 0.082% | 0.38% |
| Plums | 0.019 | 0.036 | 0.41 | 0.79 | 0.05% | 0.26% |
| Pomegranate | 0.000046 | 0.000082 | 0.001 | 0.0018 | 0.0002% | 0.00041% |
| Pomelos | 0.000097 | 0.00011 | 0.0021 | 0.0023 | 0.00027% | 0.00054% |
| <i>Pomelos (not peeled)</i> | <i>0.00062</i> | <i>0.0007</i> | <i>0.013</i> | <i>1.5%</i> | <i>0.0023%</i> | <i>0.00005</i> |
| Potatoes | 0.059 | 0.098 | 1.3 | 2.1 | 0.13% | 0.31% |
| Radish | 0.000013 | 0.000021 | 0.00027 | 0.00045 | 0.000059% | 0.000092% |
| Raisin | 0.0012 | 0.0015 | 0.026 | 0.033 | 0.0059% | 0.0076% |
| Raspberries, blackberries | 0.0034 | 0.0079 | 0.073 | 0.17 | 0.016% | 0.04% |
| Rhubarbs | 0.000036 | 0.000079 | 0.00078 | 0.0017 | 0.00011% | 0.00023% |
| Rice | 0.0014 | 0.01 | 0.032 | 0.22 | 0.018% | 0.13% |
| Rice, short grained | 0.00011 | 0.00055 | 0.0024 | 0.012 | 0.0028% | 0.014% |
| Rolled oat | 0.024 | 0.049 | 0.53 | 1.1 | 0.062% | 0.17% |
| Ruccola | 0.00058 | 0.00059 | 0.013 | 0.013 | 0.00099% | 0.0012% |
| Rye flour | 0.049 | 0.098 | 1.1 | 2.1 | 0.76% | 1.4% |
| Sesame seed | 0.0000079 | 0.000017 | 0.00017 | 0.00037 | 0.00021% | 0.00045% |
| Spinach | 0.0012 | 0.0045 | 0.027 | 0.098 | 0.014% | 0.052% |
| Spring onions | 0.000031 | 0.000064 | 0.00067 | 0.0014 | 0.00083% | 0.0019% |
| Strawberries | 0.019 | 0.045 | 0.42 | 0.99 | 0.063% | 0.2% |
| Sweet corn (small) | 0.00000058 | 0.0000044 | 0.000013 | 0.000097 | 0.00001% | 0.000095% |
| Table grapes | 0.12 | 0.24 | 2.6 | 5.2 | 0.44% | 1.6% |

| Commodity | Exposure ($\mu\text{g}/\text{kg bw}/\text{day}$) | | Exposure ($\mu\text{g}/\text{day}$) | | Hazard Index | |
|--------------------------------|--|--------------|---------------------------------------|-------------|--------------|--------------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Teas | 0.00023 | 0.00053 | 0.0051 | 0.012 | 0.0058% | 0.013% |
| Tomatoes | 0.084 | 0.27 | 1.8 | 6 | 0.55% | 2.3% |
| Vegetables, unspecified | 0.00009 | 0.0001 | 0.002 | 0.0022 | 0.0055% | 0.0061% |
| Watermelon | 0.00077 | 0.0033 | 0.017 | 0.072 | 0.015% | 0.043% |
| <i>Watermelon (not peeled)</i> | <i>0.0064</i> | <i>0.029</i> | <i>0.14</i> | <i>0.62</i> | <i>0.15%</i> | <i>0.42%</i> |
| Wheat bran | 0.002 | 0.0021 | 0.044 | 0.045 | 0.0072% | 0.008% |
| Wheat flour | 0.17 | 0.6 | 3.7 | 13 | 0.82% | 3.5% |
| Wine, red | | | | | | |
| Wine, white | | | | | | |
| Total | 2.0 | 4.5 | 44 | 98 | 10% | 44% |

7.9 Exposure and Hazard Quotients for individual pesticides (consumer group “Adults”).

Model 1: No correction for non-detects

Model 3: Non-detects assumed to be ½LOR; correction factor limited to 25.

Corrected for peeling.

| Pesticide | Exposure (µg/kg bw/day) | | Exposure (µg/day) | | Hazard Quotient | |
|-------------------------|-------------------------|----------|-------------------|---------|-----------------|----------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| 2,4-D (sum) | 0.00035 | 0.0023 | 0.026 | 0.17 | 0.0007% | 0.0046% |
| Acephate | 0.000091 | 0.0012 | 0.0068 | 0.088 | 0.0003% | 0.0039% |
| Acetamiprid | 0.0017 | 0.011 | 0.13 | 0.81 | 0.0024% | 0.015% |
| Aclonifen | 0.00001 | 0.00003 | 0.00077 | 0.0022 | 0.00002% | 0.00004% |
| Aldicarb (sum) | 0.000053 | 0.0013 | 0.004 | 0.097 | 0.0018% | 0.043% |
| Aldrin and Dieldrin | 0.000009 | 0.00023 | 0.00068 | 0.017 | 0.009% | 0.23% |
| Atrazine | 0.0000016 | 0.000041 | 0.00012 | 0.0031 | 0.00001% | 0.0002% |
| Azinphos-methyl | 0.0058 | 0.022 | 0.43 | 1.7 | 0.12% | 0.44% |
| Azoxystrobin | 0.013 | 0.021 | 0.99 | 1.6 | 0.0066% | 0.011% |
| Benalaxyl (sum) | 0.000035 | 0.00014 | 0.0026 | 0.011 | 0.00009% | 0.00036% |
| Benfuracarb | 0.00014 | 0.0017 | 0.011 | 0.13 | 0.0014% | 0.017% |
| Bifenthrin | 0.0014 | 0.0064 | 0.1 | 0.48 | 0.0092% | 0.043% |
| Binapacryl | 0.00000087 | 0.000022 | 0.000065 | 0.0016 | | |
| Biphenyl | 0.000044 | 0.0011 | 0.0033 | 0.083 | 0.00004% | 0.00089% |
| Bitertanol | 0.0058 | 0.009 | 0.43 | 0.67 | 0.19% | 0.3% |
| Bromopropylate | 0.0022 | 0.011 | 0.17 | 0.84 | 0.0075% | 0.037% |
| Bupirimate | 0.00051 | 0.012 | 0.038 | 0.9 | 0.001% | 0.024% |
| Buprofezin | 0.00014 | 0.0022 | 0.011 | 0.17 | 0.0014% | 0.022% |
| Captafol | 0.0000011 | 0.000027 | 0.000081 | 0.002 | | |
| Captan/Folpet (sum) | 0.0047 | 0.053 | 0.35 | 4 | 0.0047% | 0.053% |
| Carbaryl | 0.017 | 0.043 | 1.3 | 3.2 | 0.23% | 0.57% |
| Carbendazim and benomyl | 0.052 | 0.087 | 3.9 | 6.5 | 0.26% | 0.43% |
| Carbofuran (sum) | 0.00001 | 0.00017 | 0.00076 | 0.013 | 0.0067% | 0.11% |
| Carbosulfan | 0.00021 | 0.0048 | 0.016 | 0.36 | 0.0042% | 0.097% |
| Chlorfenvinphos | 0.00026 | 0.0028 | 0.019 | 0.21 | 0.051% | 0.57% |
| Chlormequat | 0.066 | 0.076 | 5 | 5.7 | 0.17% | 0.19% |
| Chlorothalonil | 0.0058 | 0.012 | 0.44 | 0.92 | 0.039% | 0.082% |
| Chlorpropham | 0.031 | 0.031 | 2.3 | 2.4 | 0.062% | 0.063% |
| Chlorpropham (sum) | 0.00026 | 0.0051 | 0.02 | 0.38 | 0.00052% | 0.01% |
| Chlorpyrifos | 0.014 | 0.023 | 1.1 | 1.8 | 0.14% | 0.23% |
| Chlorpyrifos-methyl | 0.0013 | 0.02 | 0.095 | 1.5 | 0.013% | 0.2% |
| Chlorthal-dimethyl | 0.000012 | 0.00031 | 0.00092 | 0.023 | 0.00012% | 0.0031% |
| Clofentezine | 0.00028 | 0.00096 | 0.021 | 0.072 | 0.0014% | 0.0048% |
| Cyfluthrin (sum) | 0.00012 | 0.0029 | 0.0093 | 0.21 | 0.0041% | 0.095% |
| Cyhalothrin, lambda- | 0.0011 | 0.0087 | 0.081 | 0.65 | 0.022% | 0.17% |
| Cypermethrin (sum) | 0.0022 | 0.01 | 0.17 | 0.78 | 0.0045% | 0.021% |
| Cyprodinil | 0.014 | 0.019 | 1 | 1.4 | 0.045% | 0.062% |
| Cyromazine | 0.0016 | 0.0084 | 0.12 | 0.63 | 0.0027% | 0.014% |
| DDT (sum) | 0.000068 | 0.0017 | 0.0051 | 0.13 | 0.00068% | 0.017% |
| Deltamethrin | 0.0013 | 0.0091 | 0.095 | 0.68 | 0.013% | 0.091% |
| Diazinon | 0.0002 | 0.0047 | 0.015 | 0.35 | 0.1% | 2.4% |
| Dichlofluanid | 0.00012 | 0.0018 | 0.0094 | 0.14 | 0.00004% | 0.00061% |
| Dichlorprop | 0.0000057 | 0.00014 | 0.00043 | 0.011 | 0.00001% | 0.00024% |
| Dichlorvos | 0.0000038 | 0.000096 | 0.00029 | 0.0072 | 0.0048% | 0.12% |
| Dicloran | 0.00086 | 0.0027 | 0.065 | 0.21 | 0.017% | 0.055% |
| Dicofol (sum) | 0.0025 | 0.017 | 0.19 | 1.3 | 0.12% | 0.86% |
| Diethofencarb | 0.00026 | 0.0035 | 0.02 | 0.27 | 0.00006% | 0.00082% |
| Difenoconazole | 0.0011 | 0.0066 | 0.081 | 0.5 | 0.011% | 0.066% |

| Pesticide | Exposure (µg/kg bw/day) | | Exposure (µg/day) | | Hazard Quotient | |
|---------------------------------------|-------------------------|------------|-------------------|----------|-----------------|----------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Diflufenican | 0.000055 | 0.0012 | 0.0042 | 0.092 | 0.00003% | 0.00061% |
| Dimethoate | 0.00049 | 0.0063 | 0.037 | 0.48 | 0.049% | 0.63% |
| Dimethomorph | 0.0064 | 0.014 | 0.48 | 1.1 | 0.013% | 0.028% |
| Diniconazole | 0.000051 | 0.0009 | 0.0038 | 0.068 | 0.00025% | 0.0045% |
| Diphenylamine | 0.061 | 0.065 | 4.6 | 4.9 | 0.081% | 0.086% |
| Dithiocarbamates | 0.098 | 0.21 | 7.4 | 16 | 0.2% | 0.42% |
| Endosulfan (sum) | 0.0027 | 0.0085 | 0.21 | 0.63 | 0.046% | 0.14% |
| Epoxiconazole | 0.000014 | 0.000077 | 0.0011 | 0.0058 | 0.00018% | 0.00097% |
| Ethion | 0.000067 | 0.00083 | 0.005 | 0.062 | 0.0034% | 0.041% |
| Ethoxyquin | 0.00023 | 0.0057 | 0.017 | 0.43 | 0.0046% | 0.11% |
| Famoxadone | 0.00063 | 0.0031 | 0.047 | 0.23 | 0.0052% | 0.026% |
| Fenarimol | 0.00012 | 0.0021 | 0.0089 | 0.16 | 0.0012% | 0.021% |
| Fenazaquin | 0.00043 | 0.0045 | 0.032 | 0.34 | 0.0085% | 0.091% |
| Fenbuconazole | 0.000025 | 0.00036 | 0.0018 | 0.027 | 0.00041% | 0.0059% |
| Fenhexamid | 0.061 | 0.072 | 4.6 | 5.4 | 0.031% | 0.036% |
| Fenitrothion | 0.00059 | 0.0054 | 0.044 | 0.41 | 0.012% | 0.11% |
| Fenoxaprop-P-Ethyl | 0.0000031 | 0.000077 | 0.00023 | 0.0058 | 0.00003% | 0.00077% |
| Fenpropathrin | 0.000041 | 0.001 | 0.003 | 0.076 | 0.00014% | 0.0034% |
| Fenpropimorph | 0.000063 | 0.0011 | 0.0048 | 0.08 | 0.0021% | 0.035% |
| Fenson | 0.00000002 | 0.0000006 | 0.0000018 | 0.000045 | | |
| Fenthion (sum) | 0.00019 | 0.00074 | 0.014 | 0.055 | 0.0027% | 0.011% |
| Fenvalerat, esfenvalerat, RR- and SS- | 0.00033 | 0.0044 | 0.025 | 0.33 | 0.0017% | 0.022% |
| Fenvalerat, esfenvalerat, RS- and SR- | 0.00015 | 0.0007 | 0.012 | 0.053 | 0.00077% | 0.0035% |
| Flucythrinate | 0.00001 | 0.00026 | 0.00078 | 0.019 | 0.00005% | 0.0013% |
| Fludioxonil | 0.0086 | 0.025 | 0.64 | 1.9 | 0.0023% | 0.0069% |
| Fluoxastrobin | 0.000002 | 0.000051 | 0.00015 | 0.0038 | 0.00001% | 0.00034% |
| Fluroxypyr (sum) | 0.00005 | 0.0013 | 0.0038 | 0.094 | 0.00001% | 0.00016% |
| Flusilazole | 0.000016 | 0.00039 | 0.0012 | 0.029 | 0.00079% | 0.02% |
| Flutriafol | 0.00083 | 0.001 | 0.063 | 0.078 | 0.0083% | 0.01% |
| Fluvalinate, tau- | 0.00036 | 0.0038 | 0.027 | 0.28 | 0.0072% | 0.076% |
| Glyphosate | 0.013 | 0.11 | 0.95 | 8.4 | 0.0042% | 0.037% |
| HCH (sum) | 0.000014 | 0.00035 | 0.0011 | 0.026 | | |
| Hexachlorobenzene | 0.000017 | 0.00032 | 0.0013 | 0.024 | | |
| Hexaconazole | 0.00011 | 0.00037 | 0.0086 | 0.028 | 0.0023% | 0.0075% |
| Hexythiazox | 0.00015 | 0.0022 | 0.011 | 0.17 | 0.00049% | 0.0074% |
| Imazalil | 0.042 | 0.072 | 3.2 | 5.4 | 0.17% | 0.29% |
| Iprodione | 0.056 | 0.099 | 4.2 | 7.4 | 0.093% | 0.16% |
| Iprovalicarb | 0.0018 | 0.0086 | 0.14 | 0.64 | 0.012% | 0.057% |
| Kresoxim-methyl | 0.00043 | 0.0029 | 0.032 | 0.21 | 0.00011% | 0.00071% |
| Lindane | 0.00000003 | 0.00000066 | 0.000002 | 0.000049 | 0.0% | 0.00001% |
| Linuron | 0.0026 | 0.01 | 0.2 | 0.76 | 0.087% | 0.34% |
| Malathion (sum) | 0.0037 | 0.019 | 0.28 | 1.4 | 0.012% | 0.064% |
| Mecoprop (sum) | 0.000034 | 0.00085 | 0.0025 | 0.064 | 0.00034% | 0.0085% |
| Mepiquat | 0.00077 | 0.0054 | 0.058 | 0.41 | 0.00039% | 0.0027% |
| Metalaxyl (sum) | 0.0026 | 0.012 | 0.2 | 0.91 | 0.0033% | 0.015% |
| Methacrifos | 0.000013 | 0.00032 | 0.000095 | 0.0024 | 0.00002% | 0.00053% |
| Methamidophos | 0.00091 | 0.0029 | 0.068 | 0.22 | 0.091% | 0.29% |
| Methidathion | 0.00025 | 0.00097 | 0.019 | 0.072 | 0.025% | 0.097% |
| Methiocarb (sum) | 0.00035 | 0.0016 | 0.026 | 0.12 | 0.0027% | 0.012% |
| Methomyl and Thiodicarb | 0.0026 | 0.0083 | 0.19 | 0.62 | 0.1% | 0.33% |
| Metribuzin | 0.00023 | 0.005 | 0.018 | 0.37 | 0.0018% | 0.038% |
| Mevinphos | 0.000044 | 0.00019 | 0.0033 | 0.014 | 0.0055% | 0.023% |
| Monocrotophos | 0.000013 | 0.000052 | 0.000095 | 0.00039 | 0.00021% | 0.00087% |
| Myclobutanil | 0.0013 | 0.006 | 0.097 | 0.45 | 0.0052% | 0.024% |
| Ofurace | 0.000095 | 0.0024 | 0.0071 | 0.18 | | |

| Pesticide | Exposure (µg/kg bw/day) | | Exposure (µg/day) | | Hazard Quotient | |
|-------------------------|-------------------------|------------|-------------------|----------|-----------------|----------|
| | Model 1 | Model 3 | Model 1 | Model 3 | Model 1 | Model 3 |
| Omethoate | 0.00022 | 0.0039 | 0.017 | 0.29 | 0.074% | 1.3% |
| Orthophenylphenol | 0.0051 | 0.011 | 0.39 | 0.84 | 0.0013% | 0.0028% |
| Oxamyl | 0.0003 | 0.0026 | 0.022 | 0.19 | 0.03% | 0.26% |
| Oxydemeton-methyl (sum) | 0.000029 | 0.00074 | 0.0022 | 0.055 | 0.0098% | 0.25% |
| Parathion-methyl (sum) | 0.000068 | 0.0016 | 0.0051 | 0.12 | 0.0023% | 0.053% |
| Penconazole | 0.00028 | 0.0019 | 0.021 | 0.14 | 0.00092% | 0.0064% |
| Pendimethalin | 0.000088 | 0.0011 | 0.0066 | 0.082 | 0.00007% | 0.00088% |
| Pentachlorobenzene | 0.000006 | 0.00015 | 0.00045 | 0.011 | | |
| Pentachlorophenol | 0.00000005 | 0.0000013 | 0.000004 | 0.0001 | | |
| Permethrin (sum) | 0.0004 | 0.0017 | 0.03 | 0.12 | 0.0008% | 0.0033% |
| Phenthoate | 0.000021 | 0.000086 | 0.0016 | 0.0065 | 0.0007% | 0.0029% |
| Phorate (sum) | 0.0000065 | 0.00016 | 0.00049 | 0.012 | 0.00093% | 0.023% |
| Phosalone | 0.0066 | 0.019 | 0.5 | 1.4 | 0.066% | 0.19% |
| Phosmet (sum) | 0.00095 | 0.021 | 0.072 | 1.6 | 0.032% | 0.69% |
| Pirimicarb (sum) | 0.002 | 0.0082 | 0.15 | 0.61 | 0.0056% | 0.023% |
| Pirimiphos-methyl | 0.016 | 0.043 | 1.2 | 3.2 | 0.4% | 1.1% |
| Prochloraz (sum) | 0.0045 | 0.0064 | 0.34 | 0.48 | 0.045% | 0.064% |
| Procymidone | 0.0076 | 0.018 | 0.57 | 1.3 | 0.27% | 0.64% |
| Profenofos | 0.000056 | 0.00046 | 0.0042 | 0.034 | 0.00019% | 0.0015% |
| Propamocarb (sum) | 0.026 | 0.032 | 2 | 2.4 | 0.0091% | 0.011% |
| Propargite | 0.021 | 0.061 | 1.6 | 4.6 | 0.21% | 0.61% |
| Propiconazole | 0.000023 | 0.00035 | 0.0017 | 0.026 | 0.00006% | 0.00087% |
| Propoxur | 0.00000002 | 0.00000039 | 0.0000012 | 0.000029 | 0.0% | 0.0% |
| Propyzamide | 0.0003 | 0.0006 | 0.022 | 0.045 | 0.0015% | 0.003% |
| Prothiofos | 0.000082 | 0.00015 | 0.00062 | 0.011 | | |
| Pymetrozine | 0.003 | 0.0057 | 0.22 | 0.42 | 0.01% | 0.019% |
| Pyraclostrobin | 0.014 | 0.023 | 1 | 1.7 | 0.045% | 0.076% |
| Pyrazophos | 0.00004 | 0.001 | 0.003 | 0.075 | 0.001% | 0.025% |
| Pyridaben | 0.00042 | 0.0027 | 0.032 | 0.2 | 0.0042% | 0.027% |
| Pyridate (sum) | 0.0000025 | 0.000056 | 0.00019 | 0.0042 | 0.00001% | 0.00015% |
| Pyrimethanil | 0.013 | 0.043 | 1 | 3.2 | 0.0079% | 0.025% |
| Pyriproxyfen | 0.0011 | 0.003 | 0.085 | 0.22 | 0.0011% | 0.003% |
| Quinalphos | 0.000012 | 0.00029 | 0.00088 | 0.022 | | |
| Quinoxifen | 0.0000033 | 0.000037 | 0.00025 | 0.0028 | 0.0% | 0.00002% |
| Quintozene (sum) | 0.00026 | 0.0061 | 0.02 | 0.46 | 0.0026% | 0.061% |
| Quizalofop | 0.000057 | 0.00086 | 0.0043 | 0.064 | | |
| Spiroxamine | 0.00057 | 0.0014 | 0.043 | 0.1 | 0.0023% | 0.0056% |
| Tebuconazole | 0.004 | 0.013 | 0.3 | 0.95 | 0.013% | 0.042% |
| Tebufenozide | 0.0017 | 0.01 | 0.13 | 0.75 | 0.0086% | 0.05% |
| Tebufenpyrad | 0.0014 | 0.0064 | 0.1 | 0.48 | 0.014% | 0.064% |
| Tecnazene | 0.0000039 | 0.000095 | 0.00029 | 0.0071 | 0.00002% | 0.00048% |
| Tetrachlorvinphos | 0.00000038 | 0.0000095 | 0.000028 | 0.00071 | | |
| Tetraconazole | 0.000043 | 0.00037 | 0.0032 | 0.028 | 0.0011% | 0.0092% |
| Tetradifon | 0.000016 | 0.00038 | 0.0012 | 0.029 | 0.0001% | 0.0026% |
| Thiabendazole | 0.1 | 0.11 | 7.5 | 8.5 | 0.1% | 0.11% |
| Thiophanate-methyl | 0.017 | 0.031 | 1.3 | 2.4 | 0.022% | 0.039% |
| Tolclofos-methyl | 0.00059 | 0.0018 | 0.044 | 0.13 | 0.00093% | 0.0028% |
| Tolyfluanid (sum) | 0.0085 | 0.019 | 0.64 | 1.4 | 0.0085% | 0.019% |
| Triadimefon (sum) | 0.015 | 0.028 | 1.1 | 2.1 | 0.029% | 0.055% |
| Triallate | 0.0000032 | 0.00008 | 0.00024 | 0.006 | 0.00001% | 0.00032% |
| Triazophos | 0.0000036 | 0.000082 | 0.00027 | 0.0062 | 0.00036% | 0.0082% |
| Trichlorfon | 0.000031 | 0.00065 | 0.0023 | 0.049 | 0.00007% | 0.0014% |
| Trifloxystrobin | 0.0019 | 0.0095 | 0.14 | 0.72 | 0.0019% | 0.0095% |
| Triflumuron | 0.0024 | 0.007 | 0.18 | 0.52 | 0.017% | 0.05% |
| Vinclozolin (sum) | 0.0065 | 0.009 | 0.49 | 0.68 | 0.13% | 0.18% |
| Hazard Index | 0.90 | 1.94 | 68 | 146 | 4% | 18% |

Fødevareinstituttet
Danmarks Tekniske Universitet
Mørkhøj Bygade 19
DK - 2860 Søborg

T: 35 88 70 00
F: 35 88 70 01
www.food.dtu.dk

ISBN: 978-87-92763-78-5